

Review Article

How Bacteria Cause Food Contamination?

ABSTRACT

Aims: to revisit the route of microbial contamination of food which can cause morbidity and mortality to vulnerable individuals and how to prevent it

Discussion: Unsafe foods or food contamination contribute to morbidity and mortality and the most common causative agent is bacteria such as *Salmonella typhi*, *Escherichia coli*, *Campylobacter* spp, *Staphylococcus aureus*, *Listeria monocytogenes*, *Clostridium perfringens*, *Vibrio parahaemolyticus* and *Bacillus cereus*. Poor personal hygiene, improper food handling combined with the existence of these opportunistic organisms can facilitate entry for these bacteria to their host. By tightly controlling the environment and other supporting conditions, potentially harmful bacteria will not be able to survive, grow, and multiply, or in other words will never be able to cause definite illness. In the food processing, controlling potentially harmful and potentially dangerous microorganisms is crucial for ensuring food safety and quality. There are six factors that affect bacterial growth, which can be directed to by the mnemonic FATTOM, namely Food source, Acid, Temperature, Time, Oxygen and Moisture.

Keywords: cross contamination

1. INTRODUCTION

Bacteria are the most common cause of global foodborne diseases [1] which roughly estimates two-thirds of human major food-borne zoonotic diseases worldwide with towering burden in developing countries [2]. Food contamination during travelling to previously mentioned countries presents unique risks to those affected and recently become a trend in epidemiology. This food contamination can cause infection [3] of the gastrointestinal (GI) system or even worse, its toxin can cause intoxication of the GI tract [4].

Bacterial gastroenteritis is an infection restricted to digestive tract which solely caused by bacteria. Its common pathognomonic symptoms include nausea, vomiting, fever, diarrhea, abdominal cramping, and pain [1-4]. Some common bacteria that contaminate food include

- *Salmonella* spp- usually related to consumption of raw or undercooked meat, poultry, and eggs or egg products or by drinking unpasteurized milk [5],
- *Escherichia coli*- contaminate foods like raw or undercooked meat, raw milk, fruits, and vegetables. It can also contaminate foods during preparation if surfaces or utensils are contaminated [6],
- *Campylobacter* spp- prevalent in food animals like poultry, cattle, pigs, sheep, and ostriches, as well as pets like cats and dogs and humans can become infected with *Campylobacter* by consuming contaminated or undercooked food, or by touching animals or their belongings [7],

- *Staphylococcus aureus* which its relationship with the occurrence and development of diabetes mellitus [8] and in the context of Staphylococcal food poisoning (SFP) is a common foodborne illness caused by ingesting *S. aureus* enterotoxins, which are heat resistant and produced in contaminated food [9],
- *Listeria monocytogenes*- a rare but serious foodborne disease with a significant public health concern because of its high mortality rate [10],
- *Clostridium perfringens*- a common cause of food poisoning, and outbreaks are often associated with large groups of people being served food [11],
- *Vibrio parahaemolyticus*- related to the consumption of raw or undercooked shellfish or contaminated water [12], and
- *Bacillus cereus*- a facultative anaerobic, toxin-producing gram-positive bacterium found in soil, vegetation, and food such as beef, turkey, rice, beans, and vegetables [13].

The most common biological agent responsible for food contamination is due to bacteria. Some bacteria have predilection to raw or undercooked food, e.g., the rod shape gram-negative *Salmonella* spp [5], the motile gram negative *Campylobacter* spp [7], the motile gram- negative *Escherichia coli* [6]; while some have the tendency causing food poisoning, e.g., the gram-positive, rod-shaped, non-spore-forming *Listeria* [10], *Staphylococcus aureus* the Gram-positive, coccus-shaped bacterium that forms clusters that look like grapes [8,9], the rod-shaped, gram-positive, anaerobic toxin producer *Clostridium perfringens* [11], the gram-negative, rod-shaped, facultative anaerobe *Vibrio parahaemolyticus* [12] and the Gram-positive aerobic or facultative anaerobic, motile, spore-forming, rod-shaped *Bacillus cereus*.

Poor personal hygiene [14], improper food handling [15] combined with the existence of this opportunistic organisms [3] can facilitate the *port de entrée* for these bacteria enter their host. For example, People who carry the bacteria *Staphylococcus aureus* (Staph), which is commonly found on the skin as part normal human microbiota [16], can contaminate food if they do not wash their hand properly before eating [17].

The world is challenged with nourishing an enormous expanding population with paradoxically more and more limited resources, including the essential food chain [18]. The increasing complexity of food supply chains, its production, and distribution of food products may increase the risk for sustainable food supply.

Food contamination with microbial agents can take place at any stage of the food chain, from farm to fork [19]. It can have happened in a number of ways, including due to poor handwashing [20], cross-contamination which can took place in domestic or public kitchen [21], inappropriate storage which caused food spoilage and food waste [22], cooking temperatures in the danger zone [23] and eat raw food/ ingredients [24]. Foods that are high in water such as ready to reheat vegetable soup [25], starch that high in bread or cereal [26], or protein such as in biofilm formation of spoiled meat [27] are more likely to be contaminated because they facilitate an ideal growing media for bacteria overgrowth [22,28]. Cross-contamination can happen anywhere during processing, packaging, storage, and even transport [21,29]. Serving contaminated food [30] or at least unknown origin of comestibles [31] can lead to food-borne illness outbreaks; A confirmed foodborne disease outbreak is defined as an incident in which two or more persons experience a similar illness after ingestion of a common food or meal or drink and epidemiologic evaluation implicates the meal or food or drink as the source of illness. Confirmed outbreaks may or may not be laboratory-confirmed.

The aim of this mini review was to revisited the route of microbial contamination of food which can cause morbidity and mortality to vulnerable individuals.

2. ROUTE OF CONTAMINATION

The most common cause of food contamination is poor food handling [33]. This includes not washing hands at the appropriate time [34] – before eating [35] and preparing food [36] or after using the toilet [37], or after blowing nose, coughing or sneezing [33]. A study conducted by Mihalache et al [38] revealed that washing hands with warm water and soap for 20 second is the most effective method investigated when hands are either dirty or greasy. Even if not proper washing, rinsing under running water for 5 second is a cleaning procedure that may significantly reduce the probability of cross-contamination, as it removes 90% of the hands' dirt. Although less effective than water and soap, the usage of

antibacterial wipes was significantly more effective than wet wipes, indicating that they are a better choice when water and soap are not available.

In addition to the food handler contribution to food contamination, using dirty utensils, such as knives and grater [39], kitchen sponges [40], other cutlery such as dish, spoon, fork [41], poor practice of not properly washing fruits and vegetables with clean water prior to consumption- even when those fruits and vegetables look so fresh [42], and storing raw and cooked food in the same place which actually increased the risk of cross contamination [43].

Sick people should not handle food because he/she can contaminate food and spread illness to others. Especially for food handler, he/she should not come to work if he/she have a fever, sore throat, cough, runny nose, or other cold and flu symptoms. They should also not come to work if he/she have vomiting, diarrhea, jaundice, or a fever accompanied by a sore throat [44,45].

Consuming under-cooked foods, particularly meat, is definitely not recommended [46]. Raw meat can contain bacteria like Salmonella [5,47], Listeria [10, 48], Clostridium difficile [49], Campylobacter [7,50], and E. coli [6,51]. These bacteria can live on the meat and spread to other foods if it is not managed properly [52]

Poor farming practices can also allow bacteria to contaminate food. This includes the heavy use of pesticides and antibiotics which favor resistant bacteria strain [53], or watering and irrigating fruits and vegetables using contaminated soil and water [54], or during field cultivation [55]. The practice of implicating inadequately composted or raw animal manure or sewage is also potentially harmful, especially in long term [56].

Fresh foods can lead to a number of illnesses. In Kenya, for instance, the contamination of meat, fruits and vegetables with human waste is relatively common. This is attributed to the use of contaminated water to wash food. Flies carrying contaminants can also directly transfer fecal matter and bacteria onto plant leaves or fruits.

Another aspects of route of transmission is by street foods are another common source of food contamination [57]. These foods are widely consumed in low- and middle-income countries because their affordable price easily accessible [58]. The hygiene of street food depends heavily on how it is handled while it is raw, cooked, and consumed along with the eating utensils they use [59]. Unhealthy food handling practices applied by street food vendors are an important cause of food contamination [33,57-59]. These chain of issues include inappropriate locations for the production of street food [60], poor quality raw materials [58,61], inappropriate water quality and supplies. Besides that, inadequate waste management practice and its disposal facilities also led to potential hazards of contamination [63].

The general sources of microbiological contamination for these food street vendor include the location of food preparation and food showcase [64], the cooking and serving equipment used, the raw materials used [65], and the personal hygiene practice of the food handler [66]. Due to the ease contamination of microbes and the growth of foodborne diseases, street foods can be a source of these illnesses [57-66]. Food hygiene and safety all be negatively affected by street food vendors' lack of adequate education background [58] and insufficient understanding due to poor knowledge [57], and may contain contaminants that provide a higher risk due to physical, chemical, and biological elements, necessitating a comprehensive examination of food safety and hygiene conducted by the authority in order to keep the safety margin of this kind of food.

3. CAN IT BE PREVENTED?

By keeping proper food hygiene and hand hygiene practices, until the level of individual, can minimize the spread of foodborne illnesses [67]. Keeping food hygiene is key to preventing the escalate of food safety hazards and foodborne illnesses that can jeopardize human life. Appropriate and proper food handling and storage (e.g., separating between the raw and cooked food) can prevent most foodborne illnesses. In order for pathogens to grow in food, certain conditions must be present, one of them is temperature [23]. By controlling the environment and conditions, even if potentially harmful bacteria are present in the unprepared or raw food [24,46-49, 51], they will not be able to survive, grow, and multiply, or in other word will never be able to cause definite illness.

In the food processing, controlling potentially harmful and potentially dangerous microorganisms is crucial for ensuring food safety and quality. There are six factors that affect bacterial growth, which can be directed to by the mnemonic FATTOM, namely Food source, Acid, Temperature, Time, Oxygen and Moisture. Each of these six factors contributes to bacterial growth in the following ways:

- Food source: Bacteria require food to survive [68]. For this reason, moist [25], starch [26], and protein-rich foods [27] are good potential sources of bacterial growth. Plant-based extracts include several compounds endowed with antimicrobial activity [69], e.g., terpenes [70], phenolic compounds [71] and Allyl isothiocyanates [72]. These three compounds have shown efficacy against spoilage microorganisms (including bacteria, yeasts, and fungi) and foodborne pathogens [70-72]. Plant-based extracts have been extensively applied to minimize microbial growth in different types of foods and to prevent the distribution of foodborne pathogens and antibiotic resistance [73]. The antimicrobial action of plant extracts (e.g., control of microbial growth, reduction in biofilm production) depends on the chemical structure of the extract and the specific microbial targets [74]. Unfortunately, despite the antimicrobial properties of these plant extracts, their application as potent food preservatives are restricted by their stability during processing or storage ambience [75]. Several strategies have been suggested to amplify its stability by way of spray-drying, encapsulation, inclusion in biopolymers, etc.). However, plant extracts can be contaminated with several dangerous toxic substances or have the potency to accumulate unwanted dangerous substance such as mycotoxins, heavy metals, residues of crop protection products [76]. Due to this reason, safety matters regarding the application of plant extracts are of great interest.
- Acid: Bacteria do not grow in acidic environments [77]. This is why acidic foods like lemon juice and vinegar able to inhibit the growth of certain pathogenic bacteria and can be used as preservatives, as reported by Sengun on carrot [78], citric acid in red wine [79], etc.
- Temperature: Most bacteria will grow rapidly between 4°C and 60°C (40°F and 140°F). This is referred to as the danger zone (see the section below for more information on the danger zone).
- Time: Bacteria require time to multiply. When small amount of bacteria is already existing, the initial risk is usually low in the beginning, but extended time with the right conditions will allow the bacteria to multiply and increase the risk of contamination [80]. That is why optimization of food storage has become a central issue for food science and biotechnology, especially in the field of functional foods [80,81].
- Oxygen: due to the types of bacteria which divided based on their oxygen need, namely the aerobic and anaerobic bacteria. The aerobic necessitate oxygen to grow; without sufficient oxygen, this aerobic will fail to live multiply and also preserve the food better, e.g., in an oxygen-free environment such as a vacuum-packaged container [82]. Anaerobic bacteria will only manage to survive and grow in oxygen-free milieu. Food that has been improperly processed and then stored at room temperature can be at risk to be contaminated by anaerobic bacteria. A common example is a product containing harmful *Clostridium botulinum* (botulism-causing) bacteria [83] that has been erroneously processed during canning [84], and then is consumed without any further cooking or reheating. Recent advancements revealed that in non-thermal physical treatments namely ionizing irradiations present an alternative to thermal sterilization, although with some limitations, to reduce number of vegetative cells and spores of this bacteria [85].
- Moisture: Bacteria need moisture to survive and will grow rapidly in moist foods [86]. Water activity is the most important factors that should be considered in controlling the microbial growth in food products. This is the reason why dry or salted foods are at lower risk of being hazardous from the perspective of bacteria growth inhibition [87].

4. CONCLUSION

Unsafe foods or food contamination which contribute to morbidity and mortality commonly caused by bacteria such as *Salmonella thyphi*, *Eschericia coli*, *Campylobacter spp*, *Staphylococcus aureus*, *Listeria monocytogenes*, *Clostridium perfringens*, *Vibrio parahaemolyticus* and *Bacillus cereus*. Its underlying condition such as poor personal hygiene, improper food handling combined with the existence of this opportunistic organisms facilitate transmission and then entry to the host. By carefully controlling the environment and other supporting conditions, pathogen bacteria will fail to survive, grow, and multiply. In the food processing, controlling potentially harmful and potentially dangerous microorganisms is crucial for ensuring food safety and quality. There are six factors that restrict bacterial growth, which can be directed to by the mnemonic FATTOM, namely Food source, Acid, Temperature, Time, Oxygen and Moisture.

CONSENT (WHERE EVER APPLICABLE)

Not needed

ETHICAL APPROVAL (WHERE EVER APPLICABLE)

Not needed

REFERENCES

1. Bintsis T. Foodborne pathogens. *AIMS Microbiol.* 2017 Jun 29;3(3):529-563. <https://doi.org/10.3934/microbiol.2017.3.529>.
2. Abebe E, Gugsu G, Ahmed M. Review on Major Food-Borne Zoonotic Bacterial Pathogens. *J Trop Med.* 2020 Jun 29;2020:4674235. <https://doi.org/10.1155/2020/4674235>.
3. Sattar SBA, Singh S. Bacterial Gastroenteritis. [Updated 2023 Aug 8]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK513295/>
4. Hernández-Cortez C, Palma-Martínez I, Gonzalez-Avila LU, Guerrero-Mandujano A, Solís RC, Castro-Escarpulli G. Food Poisoning Caused by Bacteria (Food Toxins) [Internet]. Poisoning - From Specific Toxic Agents to Novel Rapid and Simplified Techniques for Analysis. InTech; 2017. Available from: <http://dx.doi.org/10.5772/intechopen.69953>
5. Ehuwa O, Jaiswal AK, Jaiswal S. Salmonella, Food Safety and Food Handling Practices. *Foods.* 2021 Apr 21;10(5):907. <https://doi.org/10.3390/foods10050907>.
6. Mueller M, Tainter CR. Escherichia coli Infection. [Updated 2023 Jul 13]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK564298/>
7. Fischer GH, Hashmi MF, Paterek E. Campylobacter Infection. [Updated 2024 Jan 10]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK537033/>
8. Liang T, Liang Z, Wu S, Ding Y, Wu Q, Gu B. Global prevalence of Staphylococcus aureus in food products and its relationship with the occurrence and development of diabetes mellitus. *Med Adv.* 2023; 1(1): 53–78. <https://doi.org/10.1002/med4.6>
9. Fisher EL, Otto M, Cheung GYC. Basis of Virulence in Enterotoxin-Mediated Staphylococcal Food Poisoning. *Front Microbiol.* 2018 Mar 13;9:436. <https://doi.org/10.3389/fmicb.2018.00436>.
10. Schlech WF. Epidemiology and Clinical Manifestations of Listeria monocytogenes Infection. *Microbiol Spectr.* 2019; 7:10. <https://doi.org/10.1128/microbiolspec.gpp3-0014-2018>
11. Yao PY, Annamaraju P. Clostridium perfringens Infection. [Updated 2023 Aug 8]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK559049/>
12. Rezny BR, Evans DS. Vibrio parahaemolyticus Infection. [Updated 2023 Jun 26]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK459164/>
13. McDowell RH, Sands EM, Friedman H. Bacillus Cereus. [Updated 2023 Jan 23]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK459121/>
14. Payment P. Transmission of gastrointestinal diseases: hygiene as the final barrier. *Am J Infect Control.* 2001 Aug;29(4):218-21. <https://doi.org/10.1067/mic.2001.115683>.

15. Putri MS, Susanna D. Food safety knowledge, attitudes, and practices of food handlers at kitchen premises in the Port 'X' area, North Jakarta, Indonesia 2018. *Ital J Food Saf.* 2021 Nov 22;10(4):9215. <https://doi.org/10.4081/ijfs.2021.9215>.
16. Liu A, Garrett S, Hong W, Zhang J. Staphylococcus aureus Infections and Human Intestinal Microbiota. *Pathogens.* 2024; 13(4):276. <https://doi.org/10.3390/pathogens13040276>
17. Ho J, Boost MV, O'Donoghue MM. Tracking sources of Staphylococcus aureus hand contamination in food handlers by spa typing. *Am J Infect Control.* 2015 Jul 1;43(7):759-61. <https://doi.org/10.1016/j.ajic.2015.03.022>.
18. Fróna D, Szenderák J, Harangi-Rákos M. The Challenge of Feeding the World. *Sustainability.* 2019; 11(20):5816. <https://doi.org/10.3390/su11205816>
19. Tropea A. Microbial Contamination and Public Health: An Overview. *Int J Environ Res Public Health.* 2022 Jun 17;19(12):7441. <https://doi.org/10.3390/ijerph19127441>.
20. Xun Y, Shi Q, Yang N, Yang N, Li Y, Si W, et al. Associations of hand washing frequency with the incidence of illness: a systematic review and meta-analysis. *Ann Transl Med.* 2021 Mar;9(5):395. <https://doi.org/10.21037/atm-20-6005>.
21. Gorman R, Bloomfield S, Adley CC. A Study of Cross-Contamination of Foodborne Pathogens in the Domestic Kitchen in the Republic of Ireland. *International Journal of Food Microbiology,* 2002;76: 143-150. [https://doi.org/10.1016/S0168-1605\(02\)00028-4](https://doi.org/10.1016/S0168-1605(02)00028-4)
22. Karanth S, Feng S, Patra D, Pradhan AK. Linking microbial contamination to food spoilage and food waste: the role of smart packaging, spoilage risk assessments, and date labeling. *Front Microbiol.* 2023 Jun 22;14:1198124. <https://doi.org/10.3389/fmicb.2023.1198124>.
23. Kim C, Alrefaei R, Bushlaibi M, Ndegwa E, Kaseloo P, Wynn C. Influence of growth temperature on thermal tolerance of leading foodborne pathogens. *Food Sci Nutr.* 2019 Nov 16;7(12):4027-4036. <https://doi.org/10.1002/fsn3.1268>.
24. Park MS, Moon JS, Todd EC, Bahk GJ. Bacterial contamination in Saeng-go-gi, a ready-to-eat fresh raw beef dish sold in restaurants in South Korea. *J Food Prot.* 2015 Mar;78(3):619-23. <https://doi.org/10.4315/0362-028X.JFP-14-446>.
25. Alberghini G, Fabbian A, Ferioli M, Miotti Scapin R, Catellani P, Giaccone V. The Growth Potential of *Bacillus cereus* in Ready-to-Reheat Vegetable Soups. *Hygiene.* 2023; 3(3):339-350. <https://doi.org/10.3390/hygiene3030025>
26. Caro I, Portales S, Gómez M. Microbial characterization of discarded breads. *LWT,* 2023; 173:114291 <https://doi.org/10.1016/j.lwt.2022.114291>.
27. Gong J, Liu S, Wang H, Shao L, Chen S, Xu X, et al. Investigating meat-borne bacterial profiles related to biofilm formation: An in situ and in vitro assessment, *Food Control,* 2024;157:110175 <https://doi.org/10.1016/j.foodcont.2023.110175>.
28. Alexa EA, Papadochristopoulos A, O'Brien T, Burgess CM. Chapter 1 - Microbial contamination of food. Editor(s): Jaiswal AK, Shankar S. *Food Packaging and Preservation,* Academic Press, 2024, pp 3-19, ISBN 9780323900447, <https://doi.org/10.1016/B978-0-323-90044-7.00001-X>.
29. Supian K. 16 - Cross-contamination in processing, packaging, storage, and transport in halal supply chain. Ed(s): Ali ME, Nizar NNA. In *Woodhead Publishing Series in Food Science, Technology and Nutrition, Preparation and Processing of Religious and Cultural Foods* Woodhead Publishing, 2018, pp 309-321. ISBN 9780081018927. <https://doi.org/10.1016/B978-0-08-101892-7.00016-X>.
30. Friesema IH, Slegers-Fitz-James IA, Wit B, Franz E. Surveillance and characteristics of food-borne outbreaks in the Netherlands, 2006 to 2019. *Euro Surveill.* 2022 Jan;27(3):2100071. <https://doi.org/10.2807/1560-7917.ES.2022.27.3.2100071>.

31. Gargiulo AH, Duarte SG, Campos GZ, Landgraf M, Franco BDGM, Pinto UM. Food Safety Issues Related to Eating In and Eating Out. *Microorganisms*. 2022 Oct 26;10(11):2118. <https://doi.org/10.3390/microorganisms10112118>.
32. Lee H, Yoon Y. Etiological Agents Implicated in Foodborne Illness World Wide. *Food Sci Anim Resour*. 2021 Jan;41(1):1-7. <https://doi.org/10.5851/kosfa.2020.e75>.
33. Augustin JC, Kooh P, Bayeux T, Guillier L, Meyer T, Jourdan-Da Silva N, et al. Contribution of Foods and Poor Food-Handling Practices to the Burden of Foodborne Infectious Diseases in France. *Foods*. 2020 Nov 11;9(11):1644. <https://doi.org/10.3390/foods9111644>.
34. Todd ECD, Greig JD, Michaels BS, Bartleson CA, Smith D, Holah J. Outbreaks Where Food Workers Have Been Implicated in the Spread of Foodborne Disease. Part 11. Use of Antiseptics and Sanitizers in Community Settings and Issues of Hand Hygiene Compliance in Health Care and Food Industries. *Journal of Food Protection*, 2010; 73 (12):2306-20 <https://doi.org/10.4315/0362-028X-73.12.2306>.
35. Ali MM, Verrill L, Zhang Y. Self-reported hand washing behaviors and foodborne illness: a propensity score matching approach. *J Food Prot*. 2014 Mar;77(3):352-8. <https://doi.org/10.4315/0362-028X.JFP-13-286>.
36. Mihalache OA, Teixeira P, Langsrud S, Nicolau AI. Hand hygiene practices during meal preparation-a ranking among ten European countries. *BMC Public Health*. 2023 Jul 10;23(1):1315. <https://doi.org/10.1186/s12889-023-16222-5>.
37. Khan KM, Chakraborty R, Brown S, Sultana R, Colon A, Toor D, et al. Association between Handwashing Behavior and Infectious Diseases among Low-Income Community Children in Urban New Delhi, India: A Cross-Sectional Study. *Int J Environ Res Public Health*. 2021 Nov 28;18(23):12535. <https://doi.org/10.3390/ijerph182312535>.
38. Mihalache OA, Borda D, Neagu C, Teixeira P, Langsrud S, Nicolau AI. Efficacy of Removing Bacteria and Organic Dirt from Hands-A Study Based on Bioluminescence Measurements for Evaluation of Hand Hygiene When Cooking. *Int J Environ Res Public Health*. 2021 Aug 21;18(16):8828. doi: <https://doi.org/10.3390/ijerph18168828>.
39. Erickson MC, Liao J, Cannon JL, Ortega YR. Contamination of knives and graters by bacterial foodborne pathogens during slicing and grating of produce. *Food Microbiol*. 2015 Dec;52:138-45. <https://doi.org/10.1016/j.fm.2015.07.008>.
40. Wolde T, Bacha K. Microbiological Safety of Kitchen Sponges Used in Food Establishments. *Int J Food Sci*. 2016;2016:1659784. <https://doi.org/10.1155/2016/1659784>.
41. Akwila D, Aminougroho AT, Wahyudi D. Bacterial contamination test on plates, spoons, and glass at a food stall (angkringan) Surakarta City, Central Java. *Bioedupat: Pattimura Journal of Biology and Learning*,2022; 2(1): 19-22. <https://org/10.30598/bioedupat.v2.i1>.
42. Leff JW, Fierer N. Bacterial communities associated with the surfaces of fresh fruits and vegetables. *PLoS One*. 2013;8(3):e59310. <https://doi.org/10.1371/journal.pone.0059310>.
43. Kirchner M, Goulter RM, Chapman BJ, Clayton J, Jaykus LA. Cross-Contamination on Atypical Surfaces and Venues in Food Service Environments. *J Food Prot*. 2021 Jul 1;84(7):1239-1251. <https://doi.org/10.4315/JFP-20-314>.
44. Carpenter LR, Green AL, Norton DM, Frick R, Tobin-D'Angelo M, Reimann DW, Blade H, Nicholas DC, Egan JS, Everstine K, Brown LG, Le B. Food worker experiences with and beliefs about working while ill. *J Food Prot*. 2013 Dec;76(12):2146-54. <https://doi.org/10.4315/0362-028X.JFP-13-128>.
45. Norton DM, Brown LG, Frick R, Carpenter LR, Green AL, Tobin-D'Angelo M, Reimann DW, Blade H, Nicholas DC, Egan JS, Everstine K. Managerial practices regarding workers working while ill. *J Food Prot*. 2015 Jan;78(1):187-95. <https://doi.org/10.4315/0362-028X.JFP-14-134>.

46. Ballout R, Toufeili I, Kharroubi SA, Kassem II. Raw Meat Consumption and Food Safety Challenges: A Survey of Knowledge, Attitudes, and Practices of Consumers in Lebanon. *Foods*. 2023 Dec 29;13(1):118. <https://doi.org/10.3390/foods13010118>.
47. Meyer C, Thiel S, Ullrich U, Stolle A. Salmonella in raw meat and by-products from pork and beef. *J Food Prot*. 2010 Oct;73(10):1780-4. <https://doi.org/10.4315/0362-028x-73.10.1780>.
48. Okorie-Kanu OJ, Anyanwu MU, Ezenduka EV, Mgbeahuruike AC, Okorie-Kanu CO, Ugwujiem EE, et al. Occurrence and antibiogram of *Listeria* species in raw pork, beef, and chicken meats marketed in Enugu State, Southeast Nigeria. *Vet World*. 2020 Feb;13(2):317-325. <https://doi.org/10.14202/vetworld.2020.317-325>.
49. Rahimi E, Jalali M, Weese JS. Prevalence of *Clostridium difficile* in raw beef, cow, sheep, goat, camel and buffalo meat in Iran. *BMC Public Health*. 2014 Feb 5;14:119. <https://doi.org/10.1186/1471-2458-14-119>.
50. Igwaran AI, Okoh A. *Campylobacteriosis* Agents in Meat Carcasses Collected from Two District Municipalities in the Eastern Cape Province, South Africa. *Foods*. 2020; 9(2):203. <https://doi.org/10.3390/foods9020203>
51. Bude SA, Abdu R. Isolation and identification of *Escherichia coli* from raw meat at selected export abattoirs and retail houses in Bishoftu town, Ethiopia. *Acad. J. Agric. Res*. 2022;10(3): 036-041. <https://doi.org/10.15413/ajar.2021.0130>
52. Rouger A, Tresse O, Zagorec M. Bacterial Contaminants of Poultry Meat: Sources, Species, and Dynamics. *Microorganisms*. 2017 Aug 25;5(3):50. <https://doi.org/10.3390/microorganisms5030050>.
53. Friedman M. Antibiotic-resistant bacteria: prevalence in food and inactivation by food-compatible compounds and plant extracts. *J Agric Food Chem*. 2015 Apr 22;63(15):3805-22. <https://doi.org/10.1021/acs.jafc.5b00778>.
54. Steele M, Odumeru J. Irrigation water as source of foodborne pathogens on fruit and vegetables. *J Food Prot*. 2004 Dec;67(12):2839-49. <https://doi.org/10.4315/0362-028x-67.12.2839>.
55. Alegbeleye OO, Singleton I, Sant'Ana AS. Sources and contamination routes of microbial pathogens to fresh produce during field cultivation: A review. *Food Microbiol*. 2018 Aug;73:177-208. <https://doi.org/10.1016/j.fm.2018.01.003>.
56. Stiborova H, Kracmarova M, Vesela T, Biesiekierska M, Cerny J, Balik J, Demnerova K. Impact of Long-Term Manure and Sewage Sludge Application to Soil as Organic Fertilizer on the Incidence of Pathogenic Microorganisms and Antibiotic Resistance Genes. *Agronomy*. 2021; 11(7):1423. <https://doi.org/10.3390/agronomy11071423>
57. Ruchi Verma R, and Sunita Mishra S. Street Food Vendor Handling Practices and Bacterial Contamination of Ready-To-Eat Foods. *Acta Scientific Agriculture*, 2023; 7(8): 50-61. <https://doi.org/10.31080/ASAG.2023.07.1288>
58. Salamandane A, Malfeito-Ferreira M, Brito L. The Socioeconomic Factors of Street Food Vending in Developing Countries and Its Implications for Public Health: A Systematic Review. *Foods*. 2023 Oct 14;12(20):3774. <https://doi.org/10.3390/foods12203774>.
59. Trafiałek J, Drosinos EH, Laskowski W, Jakubowska-Gawlik, K, Tzamalís P, Leksawasdi N, et al. Street food vendors' hygienic practices in some Asian and EU countries – A survey. *Food Control*, 2018; 85: 212-222. <https://doi.org/10.1016/J.FOODCONT.2017.09.030>
60. Ceyhun Sezgin A, Şanlıer N. Street food consumption in terms of the food safety and health. *Journal of Human Sciences*, 2016;13(3): 4072–4083.
61. Rane S. Street vended food in developing world: hazard analyses. *Indian J Microbiol*. 2011 Jan;51(1):100-6. <https://doi.org/10.1007/s12088-011-0154-x>.
62. Mulyodarsono NDAR, Kristopo H. The Water Hygiene of Street Food Vendors in Southeast Asia: A Review. in *IOP Conf. Ser.: Earth Environ. Sci*. 2024;1324: 012104. <https://doi.org/10.1088/1755-1315/1324/1/012104>
63. Grangxabe XS, Madonsela BS, Maphanga T, Gqomfa B, Phungela TT, Malakane KC. An overview of waste management practices of street vendors in sub-saharan africa: A meta-analysis. *J Environ Manage*. 2024 Jul;364:121464. <https://doi.org/10.1016/j.jenvman.2024.121464>.

64. Moloi M, Lenetha GG, Malebo NJ. Microbial levels on street foods and food preparation surfaces in Mangaung Metropolitan Municipality. *Health SA*. 2021 Jan 29;26:1407. <https://doi.org/10.4102/hsag.v26i0.1407>.
65. Nicolas B, Abdoul BK, Aly S, Amadou OC, Ilboudo JA, & Alfred T. Hygienic status assessment of dish washing waters, utensils, hands and pieces of money from street food processing sites in Ouagadougou (Burkina Faso). *African Journal of Biotechnology*, 2006; 5. <https://doi.org/10.4314/AJB.V5i11.42982>
66. Ghosh A. An Account of Hygienic Practices and Street Food Safety Around the Medical Colleges of Kolkata, India. *J Pure Appl Microbiol*. 2023;17(4):2502-2513. <https://doi.org/10.22207/JPAM.17.4.48>
67. Manafe M, Gordon RE, Ncube LJ. Food hygiene and food safety practices of households in a township north of Tshwane, Gauteng. *Health SA*. 2023 Oct 17;28:2346. <https://doi.org/10.4102/hsag.v28i0.2346>.
68. Marco ML, Hill C, Hutkins R, Slavin J, Tancredi DJ, Merenstein D, Sanders ME. Should There Be a Recommended Daily Intake of Microbes? *J Nutr*. 2020 Dec 10;150(12):3061-3067. <https://doi.org/10.1093/jn/nxaa323>.
69. Pinto L, Tapia-Rodríguez MR, Baruzzi F, Ayala-Zavala JF. Plant Antimicrobials for Food Quality and Safety: Recent Views and Future Challenges. *Foods*. 2023 Jun 8;12(12):2315. <https://doi.org/10.3390/foods12122315>.
70. Masyita A, Mustika Sari R, Dwi Astuti A, Yasir B, Rahma Rumata N, Bin Emran T et al. Terpenes and terpenoids as main bioactive compounds of essential oils, their roles in human health and potential application as natural food preservatives. *Food Chemistry* 2022; X: 13. <https://doi.org/10.1016/j.fochx.2022.100217>
71. Hochma E, Yarmolinsky L, Khalfin B, Nisnevitch M, Ben-Shabat S, Nakonechny F. Antimicrobial Effect of Phytochemicals from Edible Plants. *Processes*. 2021; 9(11):2089. <https://doi.org/10.3390/pr9112089>
72. Lin CM, Preston JF 3rd, Wei CI. Antibacterial mechanism of allyl isothiocyanate. *J Food Prot*. 2000 Jun;63(6):727-34. <https://doi.org/10.4315/0362-028x-63.6.727>.
73. Woo SH, Sung JM, Park H, Kim J, Kim YJ, Kim TK, et al. Inhibitory effect of natural extract mixtures on microbial growth and lipid oxidation of sausages during storage. *J Anim Sci Technol*. 2023 Jan;65(1):225-243. <https://doi.org/10.5187/jast.2022.e92>.
74. Vaou N, Stavropoulou E, Voidarou CC, Tsakris Z, Rozos G, Tsigalou C, Bezirtzoglou E. Interactions between Medical Plant-Derived Bioactive Compounds: Focus on Antimicrobial Combination Effects. *Antibiotics (Basel)*. 2022 Jul 28;11(8):1014. <https://doi.org/10.3390/antibiotics11081014>.
75. Kapcum C, Uriyapongson J. Effects of storage conditions on phytochemical and stability of purple corn cob extract powder. *Food Science and Technology*. 2018;38. <https://doi.org/10.1590/1678-457x.23217>.
76. Brima EI. Toxic Elements in Different Medicinal Plants and the Impact on Human Health. *Int J Environ Res Public Health*. 2017 Oct 11;14(10):1209. <https://doi.org/10.3390/ijerph14101209>.
77. McGlynn W. The Importance of Food pH in Commercial Canning Operations. Jul 2016. Retrieved from <https://extension.okstate.edu/fact-sheets/the-importance-of-food-ph-in-commercial-canning-operations.html>
78. Sengun IY, Karapinar M. Effectiveness of lemon juice, vinegar and their mixture in the elimination of *Salmonella typhimurium* on carrots (*Daucus carota* L.). *Int J Food Microbiol*. 2004 Nov 15;96(3):301-5. <https://doi.org/10.1016/j.ijfoodmicro.2004.04.010>.
79. Comuzzo P, Battistutta F. Chapter 2 - Acidification and pH Control in Red Wines. Editor(s): Antonio Morata in book *Red Wine Technology*. Academic Press, 2019, pp 17-34, ISBN 9780128143995, <https://doi.org/10.1016/B978-0-12-814399-5.00002-5>.
80. Cabello-Olmo M, Oneca M, Torre P, Díaz JV, Encio IJ, Barajas M, Araña M. Influence of Storage Temperature and Packaging on Bacteria and Yeast Viability in a Plant-Based Fermented Food. *Foods*. 2020 Mar 7;9(3):302. <https://doi.org/10.3390/foods9030302>.
81. Tarlak F. The Use of Predictive Microbiology for the Prediction of the Shelf Life of Food Products. *Foods*. 2023; 12(24):4461. <https://doi.org/10.3390/foods12244461>

82. Ząbek K, Miciński J, Milewski S, Sobczak A. Effect of modified atmosphere packaging and vacuum packaging on quality characteristics of lamb meat. *Arch Anim Breed*. 2021 Oct 15;64(2):437-445. <https://doi.org/10.5194/aab-64-437-2021>.
83. Tiwari A, Nagalli S. Clostridium botulinum Infection. [Updated 2024 Jul 2]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK553081/>
84. Jarosz A, Grenda T, Goldsztejn M, Kozak B, Kwiatek K. Potential Risk of Botulinum Neurotoxin -producing Clostridia Occurrence in Canned Fish. *J Vet Res*. 2022 Nov 7;66(4):605-611. <https://doi.org/10.2478/jvetres-2022-0060>.
85. Munir MT, Mtimet N, Guillier L, Meurens F, Fravalo P, Federighi M, Kooh P. Physical Treatments to Control Clostridium botulinum Hazards in Food. *Foods*. 2023 Apr 7;12(8):1580. <https://doi.org/10.3390/foods12081580>.
86. Cruz-Paredes C, Tajmel D, Rousk J. Can moisture affect temperature dependences of microbial growth and respiration? *Soil Biology and Biochemistry*. 2021 May 1;156:108223. <https://doi.org/10.1016/j.soilbio.2021.108223>
87. Rezaei F, Vandergheynst JS. Critical moisture content for microbial growth in dried food-processing residues. *J Sci Food Agric*. 2010 Sep;90(12):2000-5. <https://doi.org/10.1002/jsfa.4044>.

UNDER PEER REVIEW