

## Impact of Plant Spacing and Plant Growth Regulators on Flowering and Fruiting Attributes of Strawberry cv. Winter Dawn Under Polyhouse Condition

### Abstract

The experiment was carried out under the polyhouse of the College of Agriculture, Ummadganj Kota, Rajasthan during 2021-22 and 2022-23 to evaluate the performance of spacing and plant growth regulators on flowering and fruiting attributes of strawberry (*Fragaria×ananassa* Duch.) cv. Winter Dawn under polyhouse condition. The experiment was designed in a factorial randomized block design with involving eighteen treatment combinations comprising two plant spacing viz., 15 cm×30 cm & 30 cm×30 cm and three plant growth regulators viz., salicylic acid (50 ppm, 100 ppm & 150 ppm), Triaccontanol (5 ppm, 10 ppm & 15 ppm) and NAA (25 ppm, 50 ppm & 75 ppm) respectively. The experiment revealed that maximum number of flowers per plant (25.08), days taken to flower initiation (34.69), duration of flowering (45.27), number of fruits per plant (24.36) and fruit set percentage (81.52%) was recorded under treatment S<sub>2</sub> (30 cm ×30 cm) spacing. In terms of PGRs the highest number of flowers per plant (27.11), duration of flowering (50.13), the shortest time (32.78) to flower initiation, number of fruits per plant (22.57) and fruit set percentage was (84.07%) obtained under the treatment treated with SA at 100 ppm ~~where~~ as ~~whereas~~, the interaction effect showed that the maximum number of flowers per plant (27.89), duration of flowering (51.81) and minimum days to flower initiation (32.00), number of fruits per plant (26.55) and fruit set percentage (84.75%) was recorded under S<sub>2</sub>+SA<sub>100</sub> ppm (30 cm ×30cm + Salicylic@100 ppm) treatment combination. It can be concluded that treatment combination of 30 × 30 cm +SA @ 100 ppm exhibited superior performance in terms of the number of flowers, duration of flower, days taken to flower initiation, number of fruits per plant and fruit set% among other treatment combinations used.

**Keywords:** Spacing, Plant growth regulators, Salicylic acid, Strawberry

### 1. Introduction:

Strawberry (*Fragaria x ananassa*Duch.) is a natural hybrid species which belongs to the Rosaceae family that is cultivated all over the world for its aggregate accessory fruits. The modern cultivated variety is a hybrid of *Fragaria virginiana* and *Fragaria chiloensis* (Hancock *et al.*,2000). The fruit is extensively cultivated in the temperate as well as subtropical regions of the world. Strawberry is an accessory fruit defined as a modified receptacle with many seeded or achenes located on ~~its~~ outer surface. In India, strawberry is generally cultivated in the hills and the climatic variability that exists in these areas and the broad adaptability of the plant allow its harvesting for more than half part of the year. Strawberry, a non-climacteric, delicious fruit known for ~~its~~ flavour, taste, fresh use and a rich source of vitamin C. Strawberry plants produce stolons known as runners which are the primary means of propagating strawberries commercially. Strawberry inflorescence is a modified stem called cyme terminates by a primary blossom typically followed by two secondary, four tertiary and depending on the variety and time of season eight quaternary flowers (Handley *et al.*, 1998). Flower induction in strawberry is controlled primarily by the interaction of photoperiod temperature and genotype (Darnell *et al.*, 2010). An individual flower typically has 10 green sepals five white petals 20 to 30 stamens and 60 to 600 pistils. The greatest number of pistils occurs on the primary flower and decreases successively down the inflorescence. Each pistil contains a single ovary that develops into an achene (Darnell *et al.*, 2010). The achenes (the true fruits of strawberry) together with receptacle form an aggregate fruit which is referred to as a berry (the eterio of achenes) but it is not a true berry in the botanical sense (Hancock *et al.*, 2000).

Modern cultivated strawberry is herbaceous perennial that has a central stem (crown) from which leaves, roots, stolons (runners) and inflorescences emerge. The main stem of the strawberry plant is much shortened called '~~erown~~-Plantcrown'. Plant spacing is crucial for ensuring excellent growth, output and quality. It contributes to ensuring adequate room for air circulation around and between the plants which lessens the spread of various diseases particularly in wet and humid conditions. It assists in achieving high output per square meter. Marketable strawberry yields are higher with closer spacing than with wider spacing but quality in respect to size & other parameters are good in wider spacing. Plant spacing is increased to promote growth and yield per plant. The leaf area and leaf area index are also improved. The

Commented [U1]: ? Please insert words in english

greatest harvest index and yield per square meter are provided by close ~~spacing.Plant~~spacing. Plant growth hormones are thought to be essential for many activities in the life cycle of a plant under unfavorable environmental conditions where growth ingredients alter during the growing season with corresponding fluctuations in growth and ~~development.Salicylic~~development. Salicylic acid plays a crucial role in the growth and development of the plant for crucial physiological functions such as increasing the plant's response to stress conditions (biotic and abiotic) increasing the plant's resistance to system-acquired resistance (SAR) or both. Triacntanol has been demonstrated to improve hormone functionality and root enzymatic activity when administered to plants at their peak growth period. This increases the plant's overall performance. Triacntanol essentially works by improving a plant's fundamental processes such as speeding up photosynthesis and producing more sugar or glucose. NAA hormone has many beneficial effects on plants including the ability to initiate flowering speed up, chlorophyll production, prevent fruit falling, improve crop maturity and boost agricultural yields. It can encourage the development of adventitious roots and roots of plants in addition to the basic actions of general growth hormones. It is also used to encourage the germination of seeds, the rooting of cuttings and the production of fibrous roots in agricultural crops.

## **2. Material and Methods**

### **2.1 Experimental materials**

The runners of the Winter Dawn variety of strawberry were brought from the strawberry farm situated in Pune (Maharashtra) in the month of November 2021–22 and 202-23. The runners were kept for two days in shade for hardening before being transplanted into well-prepared beds under poly-house conditions which were distributed randomly in three replications. Standard cultural practices were followed during the period of the experiment for cultivation of crops.

### **2.2 Field preparation**

The experimental area was brought to a fine tilth by ploughing followed ~~bylevelingby~~leveling. Weeds and stones were removed from the field raised beds of 15-20 cm in height were prepared for planting the strawberry ~~plants.The~~plants. The required amount of fertilizers and manures, viz., urea (80kg/ha) muriate of potash(40

kg/ha), single superphosphate (40 kg/ha) and FYM (20 ton/ha) were calculated and applied on a plot basis before transplanting and thoroughly mixed in the beds.

### 2.3 Transplanting

For the experimentation, runners of strawberry cv. Winter Dawn planted on raised beds under poly-house at the different spacing of viz, 15 cm ×30 cm and 30 cm ×30 cm in the months of November (2021-22 and 2022-23) in 54 plots. Healthy tissue cultured planting materials along with ball of earth were planted carefully with a khurpi after treatment with Ridomil Gold @2 g per l of water on 30th November of the year 2021-22 and 2023-23 respectively. While planting, care was taken so that the crown portion of the plant do not come in contact with the soil and one-third portion of plant was above the soil. This experiment was laid out in factorial ~~randomized~~ randomized block design under 18 treatment combinations.

Commented [U2]: ?

### 2.4 Mulching

To protect the strawberry plants from frost, prevent fruits from touching the ground, minimize bird attacks, conserve moisture in the soil and control weeds in the plots mulching with rice husk was done.

### 2.5 Preparation of plant growth regulator's solution and method of spray

(1) Based on the treatment, the solutions of three plant growth regulators, namely, salicylic acid (SA), naphthalene acetic acid (NAA) and triacontanol solutions were prepared by dissolving in the prescribed solvents. The salicylic acid (50 ppm, 100 ppm and 150 ppm), triacontanol (5ppm, 10 ppm and 15 ppm) and ~~NAA~~NAA (25 ppm, 50 ppm and 75 ppm) were prepared by weighing 50 mg, 100 mg and 150 mg of salicylic acid, 5 mg, 10 mg and 15 mg of triacontanol and 25 mg, 50 mg and 75 mg of NAA. Which were dissolved in a small quantity of methanol and made desired volume using distilled water.

### 2.6 Harvesting

Fruit harvesting was done when the fruits attained more than 75% color. The picking was carried out for more than one month.

### 2.7 Statistical Analysis

The data recorded during field experimentation and data obtained from laboratory determination were subjected to the statistical analysis of variance using factorial randomized block design and completely randomized design respectively as described by Panse and Sukhatme (1985). The overall significance of difference

among the treatments was tested, using critical differences (C.D.) at 5% level of significance. The results were statistically analyzed with the help of a windows-based computer package OPSTAT (Sheoran *et al.* 1998) which calculated the standard error of the difference in mean (SEd) and Critical difference (CD) between the treatments at 5% level of significance. Graphs were prepared by using Microsoft Excel (MS office ver. 2007).

### **3. Results and Discussion**

#### **3.1 Number of flowers per plant**

##### **3.1.1 Effect of spacing**

In relation to different level of spacing at 90 days after transplanting, the leading number of flowers per plant measuring (24.25), (25.91) and (25.08) was documented in S<sub>2</sub> (30 cm × 30 cm) signifying a notable superiority over the alternative spacing levels. In contrast the minimum number of flowers per plant measuring (20.34), (21.28) and (20.81) was noted in S<sub>1</sub> (15 cm × 30 cm) for both individual years and in the combined data respectively. The results suggest that using a 30 cm × 30 cm spacing is more favorable than opting for a 15 cm × 30 cm. These findings are in line with the results reported by Ramamneh *et al.* (2013), Sonkar *et al.* (2012) and Sarrwy *et al.* (2012).

##### **3.1.2 Effect of plant growth regulators (PGRs)**

Among the different levels of plant growth regulators at 90 days after transplanting the highest number of flowers per plant measuring (26.92), (27.30) and (27.11) was observed in treatment treated with SA @ 100 ppm. In close succession, NAA @ 75 ppm displayed a considerable number of flowers per plant measuring (24.80), (26.18) and (25.49) for both individual years and in the combined data. However, the minimum number of flowers measuring (18.95), (20.21) and (19.58) were noted in treatment treated with TA @ 5 ppm for both individual years and in the pooled data ~~respectively. The~~ ~~respectively. The~~ findings showed that using SA 100 ppm was the most successful compared to other plant growth regulators, resulting in the most positive outcomes. The current research aligns with the findings reported by Qureshi *et al.* (2013), Kazemi (2013) and Jamali *et al.* (2011).

##### **3.1.3 Interaction effect of spacing and plant growth regulators (S × PGRs)**

The combined influence of different spacing levels and plant growth regulators on number of flowers per plant at 90 days after transplanting was

determined to be significant for both individual year as well as in pooled. The maximum number of flowers per plant measuring (27.66), (28.13) and (27.89) was observed in treatment combination of S<sub>2</sub>+SA100 ppm (30 cm ×30 cm + SA@100 ppm). In close succession, treatment combination of S<sub>2</sub>+NAA75 ppm (30 cm × 30 cm + NAA@75 ppm) displayed measurements of (26.49), (27.92) and (27.21). In contrast, the ~~least number~~ least number of flowers per plant measuring (15.65), (16.30) and (15.98) was recorded in treatment combination of S<sub>1</sub>+TA<sub>5</sub> ppm (15 cm × 30 cm +TA@5ppm) for the years 2021-22, 2022-23 and in the pooled data, respectively. The present study is also in consonance with the reports of Sharma *et al.* (2020) and Jamali *et al.* (2011).

### 3.2 Days Taken to Initiation of Flowering

#### 3.2.1 Effect of spacing

The initiation of flowering was significantly affected by the application of various level of spacing. Among the spacing levels early flowering occurred at (35.49), (33.88) and (34.69) days in S<sub>2</sub> (30 cm×30 cm). In contrast, a delayed onset of flowering at (38.40), (36.69) and (37.55) days was observed in specifically in S<sub>1</sub> (15 cm ×30 cm). The outcomes imply that choosing 30 cm ×30 cm spacing is better than selecting a 15 cm×30 cm spacing. The results of the present study are in conformity with the findings by Sarrwy *et al.* (2012) and Dwivedi *et al.* (2009) who reported early flowering and fruiting in wider spaced plants.

#### 3.2.2 Effect of plant growth regulators (PGRs)

Upon examination of the data, it was evident that the application of plant growth regulators had a significant impact hastening the initiation of the flowering for both years and in pooled. In the context of various levels of plant growth ~~regulators~~ regulators, the timing of flowering initiation was found to be noteworthy. Early flowering, occurring at (33.65 days), (31.89 days) and (32.78 days) was observed in treatment treated with SA @ 100 ppm which was followed by NAA @ 75 ppm with a flowering initiation period of (34.20 days), (32.44 days) and (33.32 days). ~~On~~ On the other hand, delayed flowering at (40.68 days), (38.96 days) and (39.82 days) was noted in plants treated with TA @ 5 ppm. The results indicated that application of SA 100 ppm was the most effective among various plant growth regulators, leading to the most favorable outcomes. Similar findings were reported by Eshghi *et al.* (2017)

### 3.2.3 Interaction effect of spacing and plant growth regulators (S × PGRs)

The duration to the initiation of flowering was found to be significantly affected by various treatment combinations for both individual year and in pooled. When considering the interactions between spacing and plant growth regulators treatment combination S<sub>2</sub>+SA100 ppm (30 cm ×30 cm + SA @ 100 ppm) exhibited the shortest period to flowering recording (32.88 days), (31.12 days) and (32.00 days). This was followed by treatment combination of S<sub>2</sub>+NAA75 ppm which exhibited (33.23 days), (31.47 days) and (32.35 days) to initiate flowering. In contrast, late flowering was observed in the treatment treated with S<sub>1</sub>+TA5 ppm (30 cm ×30 cm + TA @ 5 ppm) which took (41.28 days), (39.52 days) and (40.40 days) to initiate ~~floweringforflowering for~~ both individual year and in pooled data. The present study is also in consonance with the reports of Kazemi *et al.* (2013).

### 3.3 Duration of Flowering (Days)

#### 3.3.1 Effect of spacing

Among these spacing maximum duration of flowering was observed at (44.80 days), (45.74 days) and (45.27 days) in S<sub>2</sub> (30 cm ×30 cm). However, a minimum duration of flowering occurred at (41.95 days), (43.29 days) and (42.62 days) was noted specifically in S<sub>1</sub> (15 cm ×30 cm). The results suggest that using a 30 cm ×30 cm spacing is more favorable than opting for a 15 cm × 30 cm spacing. These results are in line with the findings of Hazarika *et al.* (2019) and Sarrwy *et al.* (2012).

#### 3.3.2 Effect of plant growth regulators (PGRs)

Upon scrutinizing the data, it became apparent that the application of plant growth regulators significantly increases the duration of flowering. Considering various levels of plant growth ~~regulatorsregulators~~, the timing of the duration of flowering was noteworthy. Maximum duration of flowering occurring at (49.44 days), (50.81 days) and (50.13 days) was observed in treatment treated with SA @ 100 ppm followed by NAA @ 75 ppm with a duration of flowering period of (46.81 days), (48.06 days) and (47.74 days). on the other hand, minimum duration of flowering at (38.26 days), (39.06 days) and (38.66 days) was noted in plants treated with TA @ 5 ppm for both individual year and in pooled ~~data.The data. The~~ findings suggested that utilizing SA at a concentration of 100 ppm was the most successful compared to other plant growth regulators, resulting in the most positive outcomes. These findings are consistent with the results reported by Ghaderi *et al.* (2015).

### 3.3.3 Interaction effect of spacing and plant growth regulators (S × PGRs)

The time taken for the duration of flowering was found to be significantly influenced by various treatment combinations for both years and in pooled. When examining the interactions between spacing and plant growth regulators, treatment combination of S<sub>2</sub>+SA100 ppm (30 cm ×30 cm + Salicylic@100 ppm) demonstrated the maximum period to flowering recording (50.92 days), (52.70 days) and (51.81 days). Following closely, treatment combination of S<sub>2</sub>+NAA75 ppm exhibited (47.59 days), (48.53 days) and (48.06 days) to duration of flowering. In contrast, minimum duration of flowering was observed in the treatment treated with S<sub>2</sub>+TA5 ppm (15 cm ×30 cm + TA @ 5 ppm) taking (36.13 days), (36.64 days) and (36.39 days) to duration of flowering for both individual years and in the pooled data. These findings are consistent with the results reported by Ghaderi *et al.* (2015).

### 3.4 Number of fruits per plant

#### 3.4.1 Effect of spacing

Upon analyzing the data, it is apparent that different level of spacing have a distinct impact on number of fruits per plant, the highest number of fruits per plant measured at (23.03), (25.68) and (24.36) were observed in S<sub>2</sub> (30 cm ×30 cm) for the years 2021-22, 2022-23 and in the overall combined data. In contrast, the number of fruits per plant measured at (15.55), (17.18) and (16.37) was noted at the minimum spacing level, S<sub>1</sub> (15 cm ×30 cm) for the years 2021-22, 2022-23 and in the combined data, respectively. The findings indicate that a spacing of 30 cm × 30 cm is more effective than a spacing of 15 cm×30 cm. This conforms with the views of Dogar *et al.* (2020) and Shahzad *et al.* (2018).

#### 3.4.2 Effect of plant growth regulators (PGRs)

In the assessment of different levels of plant growth regulators (PGRs) on number of fruits per plant a superior number of fruits per plant recorded at (21.07), (24.06) and (22.57) was observed with the foliar application of SA at 100 ppm in the years 2021-22, 2022-23 and the combined data respectively. Following closely, number of fruits per plant measurements of (20.91), (23.94) and (22.43) were noted with NAA at 75 ppm. In contrast, the minimum number of fruits per plant measuring (17.06), (19.47) and (18.26) was recorded with TA at 5 ppm for both individual years and in the combined data. The findings indicated that utilizing salicylic acid (SA) at a concentration of 100 ppm was more effective than employing other plant growth

regulators, leading to the most favorable results in terms of plant growth and development. These findings align with the results observed by Kumar *et al.* (2019) Singh *et al.* (2001)

### **3.4.3 Interaction effect of spacing and plant growth regulators (S × PGRs)**

The association between different spacing levels and plant growth regulators resulted significant for both individual year and in pooled. The maximum number of fruits per plant recorded at (25.09), (28.00) and (26.55) were observed in treatment combination of S<sub>2</sub>+SA100 ppm (30 cm ×30 cm + Salicylic acid@100 ppm). Following this trend, S<sub>2</sub>+NAA75ppm (30 cm ×30 cm + NAA at 75 ppm) exhibited number of fruits per plant (24.93), (27.87) and (26.40) for both individual years and in the combined data. On the other hand, the minimum number of fruits per plant, measuring (13.00), (14.87) and (13.93) was noted in treatment combination of S<sub>1</sub>+TA5ppm (15 cm ×30 cm spacing + TA at 5 ppm) for the years 2021-22, 2022-23 and in the combined data, respectively. These findings are consistent with the results reported by Mohamed *et al.* (2018)

## **3.5 Fruit Set %**

### **3.5.1 Effect of spacing**

Concerning examining the data, it is apparent that different level of spacing have a distinct impact on fruits set percent. The highest levels of fruit set percent, measured at 80.62%, 82.42% and 81.52% were observed in S<sub>2</sub> (30 cm ×30 cm) for the individual years 2021-22, 2022-23 and in pooled respectively. In contrast, the fruit set percent, measured at 78.73%, (79.81%) and (79.27%) was noted at the minimum spacing level S<sub>1</sub> (15 cm ×30 cm) for the years 2021-22, 2022-23 and in the combined data, respectively. The results suggest that using a 30 cm ×30 centimeter spacing is more favorable than opting for a 15 cm ×30 centimeter spacing. Research results are in accordance with the findings of Singh *et al.* (2018), Tripathi *et al.* (2000) and Sharma *et al.* (2020)

### **3.5.2 Effect of plant growth regulators (PGRs)**

In evaluating various levels of plant growth regulators (PGRs) on the fruits set percent, a higher fruits set percent, recorded at (83.54%), (84.59%) and (84.07%) was observed with the foliar application of SA at 100 ppm in the year of 2021-22, 2022-23 and the combined data, respectively. Following closely, the fruits set percent, measuring (82.56%), (84.22%) and (83.39%) were noted with NAA at 75 ppm. In

contrast, the minimum fruits set percent, measuring (75.21%), (76.33%) and (75.77%) was recorded with TA a@5 ppm for both individual years and in the combined data. The study results suggested that using salicylic acid (SA) at a concentration of 100 ppm was more successful than the use of other plant growth regulators. This resulted in the best outcomes for the growth and development of the plants. These findings align with the results observed by Kazemi (2013) in strawberries.

### **3.5.3 Interaction effect of spacing and plant growth regulators (S × PGRs)**

The correlation between different spacing levels and plant growth regulators significantly influenced the fruit set percent for both individual year and in pooled. The highest fruit set percent, recorded at (84.16%), (85.34%) and (84.75%) were observed in treatment combination of S<sub>2</sub>+SA 100 ppm (30 cm ×30 cm + SA at 100 ppm). In line with this pattern, treatment combination of S<sub>2</sub>+NAA75ppm (30 cm ×30 cm + NAA at 75 ppm) exhibited fruit set percent (83.77%), (85.07%) and (84.43%) for both individual years and in the combined data. However, the minimum fruits set percent, measuring (74.60%), (75.37%) and (74.98%) was noted in treatment combination of S<sub>1</sub>+TA5ppm (15 cm ×30 cm spacing + TA at 5 ppm) for the years 2021-22, 2022-23 and in the combined data, respectively. These findings are consistent with the results reported by Ghaderi *et al.* (2015).

## **4. Conclusion**

A plant spacing of 30 cm × 30 cm combined with the application of salicylic acid at a concentration of 100 ppm demonstrated the best results regarding the number of flowers per plant, duration of flowering, time taken for flower initiation, number of fruits per plant and ~~and~~ fruit set %. This research can benefit farmers by showing that adjusting plant spacing and using plant growth regulators (PGRs) can effectively improve flowering characteristics, leading to more successful and profitable strawberry cultivation.

Table 1-Effect of Spacing and PGRs on Number of flowers per plant, Days taken to initiation of flowering, Duration of flowering , Number of fruits per plant and fruit set of Strawberry cv. Winter Dawn

Treatments	Number of Flowers per plant			Days taken to initiation of flowering (Days)			Duration of flowering (Days)			Number of fruits per plant			Fruit set (%)		
	2021-22	2022-23	Pooled	2021-22	2022-23	Pooled	2021-22	2022-23	Pooled	2021-22	2022-23	Pooled	2021-22	2022-23	Pooled
<b>Spacing(S)</b>															
S <sub>1</sub> 15 × 30cm	20.34	21.28	20.81	38.40	36.69	37.55	41.95	43.29	42.62	15.55	17.18	16.37	78.73	79.81	79.27
S <sub>2</sub> 30 × 30 cm	24.25	25.91	25.08	35.49	33.89	34.69	44.80	45.74	45.27	23.03	25.68	24.36	80.62	82.42	81.52
SE(m)±	0.48	0.46	0.46	0.01	0.43	0.22	0.63	0.52	0.40	0.18	0.22	0.08	0.34	0.09	0.18
CD(P=0.05)	1.39	1.32	1.34	0.03	1.25	0.63	1.81	1.49	1.17	0.51	0.64	0.23	0.98	0.26	0.53
<b>Plant Growth Regulators (PGRs)</b>															
SA 50 ppm	23.23	24.17	23.70	36.16	34.69	35.43	44.59	45.33	44.96	19.87	21.94	20.90	80.97	82.27	81.62
SA 100 ppm	26.92	27.30	27.11	33.65	31.89	32.78	49.44	50.81	50.13	21.07	24.06	22.57	83.54	84.59	84.07
SA 150 ppm	24.00	24.93	24.47	34.40	32.99	33.70	45.83	47.01	46.42	20.19	23.18	21.69	81.14	82.83	81.99
NAA 25 ppm	20.85	22.91	21.88	38.05	36.29	37.17	41.53	43.08	42.31	18.70	20.41	19.56	78.99	80.82	79.91
NAA 50 ppm	21.17	23.13	22.15	37.01	35.49	36.25	43.56	44.72	44.14	19.51	20.57	20.04	79.68	81.67	80.68
NAA 75 ppm	24.80	26.18	25.49	34.20	32.44	33.32	46.81	48.06	47.44	20.91	23.94	22.43	82.56	84.22	83.39
TA 5 ppm	18.95	20.21	19.58	40.68	38.96	39.82	38.26	39.06	38.66	17.06	19.47	18.26	75.21	76.33	75.77
TA 10 ppm	20.07	21.28	20.67	39.69	37.93	38.81	39.31	40.57	39.94	17.89	19.57	18.73	76.73	77.81	77.28
TA 15 ppm	20.66	22.27	21.46	38.68	36.91	37.80	41.02	42.03	41.53	18.46	19.76	19.11	78.23	79.49	78.86
SE(m)±	0.23	0.22	0.22	0.01	0.20	0.10	0.30	0.24	0.19	0.08	0.10	0.04	0.16	0.04	0.09
CD(p=0.05)	0.66	0.62	0.63	0.02	0.59	0.30	0.85	0.70	0.55	0.24	0.30	0.11	0.46	0.12	0.25
<b>Interaction effect of Spacing and PGRs</b>															
Treatment Combinations															
S <sub>1</sub> +SA 50 ppm	22.12	22.40	22.26	38.19	36.43	37.32	44.25	45.03	44.64	16.00	18.34	17.17	80.28	81.19	80.74
S <sub>2</sub> +SA 50 ppm	24.35	25.93	25.14	34.13	32.95	33.54	44.92	45.64	45.28	23.73	25.53	24.63	81.67	83.34	82.51
S <sub>1</sub> + SA 100 ppm	26.18	26.48	26.33	34.42	32.66	33.55	47.96	48.92	48.44	17.05	20.12	18.59	82.92	83.84	83.38
S <sub>2</sub> +SA 100 ppm	27.66	28.13	27.89	32.88	31.12	32.00	50.92	52.70	51.81	25.09	28.00	26.55	84.16	85.34	84.75
S <sub>1</sub> +SA 150 ppm	22.55	22.69	22.62	35.36	33.60	34.49	45.76	46.44	46.10	16.40	19.24	17.82	80.41	81.44	80.93
S <sub>2</sub> +SA 150 ppm	25.45	27.17	26.31	33.43	32.37	32.90	45.91	47.57	46.74	23.97	27.12	25.55	81.87	84.21	83.04
S <sub>1</sub> +NAA 25 ppm	18.55	20.52	19.54	40.06	38.30	39.19	38.83	41.23	40.03	15.31	15.87	15.59	78.21	79.32	78.77
S <sub>2</sub> +NAA 25 ppm	23.15	25.30	24.23	36.03	34.27	35.15	44.23	44.92	44.58	22.09	24.94	23.52	79.78	82.31	81.05
S <sub>1</sub> + NAA 50 ppm	18.72	20.78	19.76	39.31	37.99	38.65	42.52	44.16	43.34	15.89	16.02	15.96	79.42	80.02	79.72
S <sub>2</sub> +NAA 50 ppm	23.61	25.48	24.55	34.71	33.00	33.86	44.60	45.27	44.94	23.12	25.12	24.12	79.95	83.31	81.63
S <sub>1</sub> +NAA 75 ppm	23.11	24.43	23.77	35.17	33.41	34.29	46.03	47.59	46.81	16.90	20.00	18.45	81.34	83.37	82.36
S <sub>2</sub> + NAA 75 ppm	26.49	27.92	27.21	33.23	31.47	32.35	47.59	48.53	48.06	24.93	27.87	26.40	83.77	85.08	84.43
S <sub>1</sub> +TA 5 ppm	15.65	16.30	15.98	41.28	39.52	40.40	36.13	36.64	36.39	13.00	14.87	13.93	74.60	75.37	74.98
S <sub>2</sub> +TA 5 ppm	22.26	24.12	23.19	40.07	38.40	39.24	40.38	41.49	40.93	21.11	24.07	22.59	75.83	77.29	76.56
S <sub>1</sub> +TA 10 ppm	17.64	18.30	17.97	41.13	39.37	40.26	37.54	39.54	38.54	14.43	15.00	14.72	74.69	76.28	75.49
S <sub>2</sub> +TA 10 ppm	22.49	24.25	23.37	38.24	36.48	37.37	41.08	41.60	41.34	21.34	24.15	22.75	78.77	79.34	79.06
S <sub>1</sub> +TA 15 ppm	18.51	19.60	19.05	40.67	38.91	39.79	38.49	40.06	39.28	15.00	15.17	15.09	76.70	77.44	77.07
S <sub>2</sub> +TA 15 ppm	22.81	24.93	23.87	36.68	34.92	35.80	43.55	43.99	43.78	21.92	24.34	23.13	79.76	81.53	80.65
SE(m)±	0.68	0.65	0.65	0.02	0.61	0.31	0.88	0.73	0.57	0.25	0.31	0.11	0.48	0.13	0.26
CD(p=0.05)	1.97	1.87	1.89	0.04	1.77	0.89	2.55	2.11	1.65	0.72	0.90	0.32	1.38	0.37	0.75

## References:

1. A.O.A.C. 1980. Official methods of analysis. Association of Official Analytical Chemist. Washington, D.C. USA.
2. Awasthi, RP. and Badiyala, SD. (1983). Spacing and performance studies in some strawberry-cultivars. *Indian Journal of Horticulture*, **40**(1 and 2): 29-34.
3. Baba, T.R., Ali, A., Kumar, A. and Husain, M (2017). Effect of exogenous application of salicylic acid and triacontanol on growth characters and yield of strawberry. *The Pharma Innovation Journal*, **6**(11): 274-279.
4. Bhatia, S.K., Sharma, R., Kumar, R (2017). Effect of different planting time and spacing on growth, yield and quality of strawberry (*Fragaria × ananassa*) cv. Ofra. *International Journal of Pure and Applied Bioscience*, **5**(5): 207-211.
5. Camacaro, ME., Camacaro, GJ., Hadley, P., Dennett, MD. and carew, JG. (2004). Effect of plant –spacing and initial crown size on growth, development and yield in strawberry cultivar Elsanta and Bolero. *The Journal of Horticultural Sciences and Biotechnology*, **79**(5): 739-746.
6. Darnell, R. L., Cantliffe, D. J., Kirschbaum, D. S., Chandler, C. K., &Janick, J. (2010). The physiology of flowering in strawberry. *Horticultural Reviews*, **28**: 325-345.
7. Diwivedi, D., Maurya, D., Ram, RB. and Chaturvedi, SK. (2009). Effect of spacing and susem of planting on biological parameters of strawberry cv. Chandler. *Progressive Horticulture*, **41**(1): 40-42.
8. Dogar, W. A., Ahmad, T., Umar, M., Nadeem, S., Bukhari, M. A., & Shah, A. N. (2020). Study to determine the effect of spacing and varieties on fruit quality of citrus. *JPAA*, **5**(4): 74-82.
9. Eshghi, S., Moharami, S., &Jamali, B. (2017). Effect of salicylic acid on growth, yield and fruit quality of strawberry cv. 'Paros' under salinity conditions. *Journal of Soil and Plant Interactions*-Isfahan University of Technology, **7**(4): 163-174.
10. Ghaderi, N., Normohammadi, S., &Javadi, T. (2015). Morpho-physiological responses of strawberry (*Fragaria× ananassa*) to exogenous salicylic acid application under drought stress.*Journl of Agricultural Science and Tecgnology*, **17**(1): 167-168.

11. Goulart, B. L., &Funt, R. C. (1986). Influence of raised beds and plant spacing on growth and yield of strawberry. *Journal of the American Society for Horticultural Science*, **111**(2): 176-181.
12. Hancock, J., &Erez, A. (2000). Strawberries. Temperate Fruit Crops in Warm Climate.*Springer Science and Business Media*.
13. Handley, D. T., Schupp, M. A., &Dill, J. F. (1998). Strawberry cultivar performance and susceptibility to tarnished plant bug injury. *HortScience*, **33**(3): 460-460.
14. Hazarika, TK., Lalchhanmawia, J.and Nautiyal, BP (2019). Effect of Planting density on growth, yield and quality of strawberry (*Fragaria×ananassa*) cv. Camarosa. *Indian Journal of Agricultural Sciences*,**89**(3): 489-493.
15. Jamali, B., Eshghi, S. and Tafazoli, E (2011). Vegetative and reproductive growth of strawberry plants cv. Pajaro affected by salicylic acid and nickel. *Journal of Agriculture Sciences and Technology*,**13**: 895-904.
16. Kazemi, M (2013). Foliar application of salicylic acid and calcium on yield, yield component and chemical properties of strawberry. *Bulletin of Environment, Pharmacology and Life Sciences*,**2**(11): 19-23.
17. Kumar, S., & Kaur, G. (2019). Effect of pre and post harvest applications of salicylic acid on quality attributes and storage behaviour of strawberry cv. Chandler. *Journal of Pharmacognosy and Phytochemistry*, **8**(4): 516-522.
18. Kumar, S., & Kaur, G. (2019). Effect of pre and post harvest applications of salicylic acid on quality attributes and storage behaviour of strawberry cv. Chandler. *Journal of Pharmacognosy and Phytochemistry*, **8**(4): 516-522.
19. Metwally, A.A., Youssef, S.M., Miniawy-El S.M., Ragab, M.E (2013). Effect of foliar spraying of salicylic acid on growth, yield and quality of cold stored strawberry plant. *Journal of Biological Chemistry and Environmental Sciences*, **8**(1): 1-17.
20. Mohamed, R.A., Abdelbaset, A.K., Elkader, A.D.Y (2018). Salicylic acid effects on growth, yield, and fruit quality of strawberry cultivars. *Journal of Medicinally Active Plants*, **6** : 1-4.
21. Muhi, H. M., &Abd, A. M. (2022). Effect Of foliar spray with tricacontanol, boron and low biuret urea on flowering and fruiting of apple trees cv. Sharaby. *Biochemical & Cellular Archives*, **22**(1): 2963-2968

22. Panse, V.G. and Shukhatme, P.V. (1967). Statistical Methods for Agricultural Workers, 2nd eds. Indian Council of Agriculture Research, New Delhi.
23. Ramamneh, E., Rawashdeh, Z., Karajeh, M. and Romman, S (2013). Plant response of strawberry to intra-row spacing and growing conditions in south of Jordan. *Asia Journal of Plant Sciences*, **12**(5): 201-207.
24. Sarrwy, S.M.A.; Mostafa, E.A.M. and Hassan, H.S.A. (2012). Growth, yield and fruit quality of Williams Banana affected by different planting distances. *International Journal of Agricultural Sciences*, **7**(5): 266-275.
25. Shahzad, U., Muhammad, I., Noor, N., Muhammad, S., Hassan, Z., Khan, A. A (2018). Variation in growing media and plant spacing for the improved production of Strawberry (*Fragaria × ananassa*) cv. Chandler. *Philippine Journal of Sciences*, **147**(4): 711-719.
26. Sharma, K. and Negi, M (2020). Impact of different spacing and combination with organic manures on growth and yield attributes of strawberry (*Fragaria × ananassa*) cv. Jutogh special under mid-hill conditions of Uttarakhand. *Journal of Pharmacognosy and phytochemical*, **9**(2): 1014-1018.
27. Singh, A., Kumar, S. S., Sharma, M. and Singh, K. S (2018). Impact of mulching and planting density on phenology, yield and fruiting characters of strawberry (*Fragaria × ananassa* Duch) cv. Winter Dawn. *Plant Archives*, **18**(2): 1899-1904
28. Singh, V. K., Saini, J. P., and Misra, A. K. (2001). Response of salicylic acid on flowering, floral malformation, fruit set, yield and associated bio-physical and biochemical characters of mango. *Indian journal of Horticulture*, **58**(3): 196-201.
29. Sonkar, P., Ram, R.B. and Meena, M.L (2012). Effect of various mulch materials and spacing on growth, yield and quality of strawberry. *Hort Flora Research Spectrum*, **1**(4): 323-327.
30. Tripathi, V.K.; Dwivedi, M.P.; Sharma, R.M. and Agrahari, P.R. (2000). Effect of planting date and spacing on yield and quality of Chandler strawberry (*Fragaria x Ananassa* Duch.). *Haryana Journal of Horticultural Sciences*, **29**(3/4): 185-186.