

Original Research Article

Detection and Analysis of Land Use and Land Cover Dynamics in the Highlands of Northern Ethiopia

Abstract

In the highlands of northern Ethiopia, many land use and land cover changes (LULCC) have occurred during different activities of the populations. Despite its coverage and unwise use of natural resources, these land uses and land covers are facing increasing or decreasing their area coverage. Thus, this study is designed to identify the basis of the above-mentioned problems and to set real solutions for the problems. Hence, a field survey was carried out in 2017 to identify the major land cover types in the watershed. Detection Analysis of land use and land cover dynamics were calculated from the selected watershed in Southern Tigray, Northern Ethiopia. These land use types were cultivated, forest, bush, shrub, and homestead lands. The land use and land cover dynamics were estimated using landsat7 and landsat8 for the years 1998 and 2017, respectively, using Arc GIS 10.1. Furthermore, a normalized difference vegetation index (NDVI) was developed for both years. The land use and land cover change results indicated that forest, cultivated land, and bush land decreased by 4.4%, 2.2%, and 2.4%, respectively. However, shrub land and homestead increased by 5.5% and 3.6%, respectively. Therefore, the change detection analysis using GIS and remote sensing could deliver useful information to understand the seasonal patterns of land use dynamics for planners and decision makers; consequently, sustainable land management planning is possible.

Keyword: Satellite image, Change detection, Land use and land cover, Watershed,

1. INTRODUCTION

As Deng (2011) and Bin et al. (2006) reported, the global environment is facing severe challenges such as changes in land use and land cover (LULC), which is the prominent ecological symbol on the earth's surface. Moreover, Turner et al. (2007) reported that LULC change has been a key research priority with multi-directional impacts on both human and natural systems yet also a challenging research theme in the field of land use change science. Land use affects land cover and land changes in land cover (Rawat and Kumar, 2015). This articulates social, environmental, institutional, and economic processes (Mas et al., 2014). Human impacts on the global LULC change are operating at unprecedented magnitudes, rates, and spatial scales (Turner et al., 1994). Scientists recognize that the magnitude of this change is large (Lambin et al., 2001).

The LULC pattern of a region is an outcome of natural and socio-economic factors and their utilization by man in time and space. These driving forces, which are also referred to as factors that can be categorized as natural and human-induced, were recognized in the study Watershed (Yifru A et al., 2011). The natural factors in the study area include the high intensity of rainfall and steep relief (Yimer, F. et al., 2007) and soil types. Among the straightforward identified human causative factors are population growth and density, over-intensification of land use, farm size, land tenure status, and lack of policies on land use planning (Yimer, F. et al., 2007).

As rightly noted by Yifru A et al. (2011) land use and land cover are an endlessly changing process taking place on the surface of the earth. Major land cover changes have also occurred at a local level for all land types. For instance, a significant increase in cultivated land at the expense of forestland was found to have occurred between 1957 and 1995 in the Dembecha area, northwest Ethiopia (Gete, 2001). Kebrom and Hedlund (2000) reported settlements increased at the expense of shrublands and forests between 1958 and 1986 in the Kalu area, north-central Ethiopia. On the other hand, the deforestation trend was reduced through appropriate interventions by promoting the planting of local tree species in the Chemoga Watershed, Blue Nile basin, Ethiopia (Woldeamlak, 2003) between 1957 and 1998. Similarly, Alemayehu (2007) concluded that in the Beressa Watershed (central Ethiopia), there were substantial land use changes in the area during the second half of the 20th century. The most important changes were the destruction of the natural vegetation, increased plantations, and expansion of grazing land.

To understand the interaction effect of LULCC on earth systems (e.g., hydrosphere, biosphere, and atmosphere), accurate information on what type of change occurs, when and where it occurs, and the rate of changes that occur is needed (Lambin, 2001). Wejic watershed requires Detection and Analysis of Land Use and Land Cover Dynamics study, which necessarily leads to an improved understanding of the process of land use and land cover change, which researchers and policymakers would pursue to establish effective conservation and management strategies for land resources.

2. MATERIALS AND METHODS

2.1. Area description

The study was undertaken at Wejic Watershed in Raya Azebo district, southern Tigray (Northern Ethiopia) (Figure 1). It is about 5 km west of Moheni town, the district capital. It lies between 39°16'30" to 39°37'30" E and 12°40'30" to 12°49'30" N, with an altitude ranging from 1828 to 3450 m.a.s.l.




Figure 1 Map of the study watershed

The study watershed is located in three districts, Emba Alaje (11.5 %), Enda Mohoni (10.9 %), and Raya Azebo (77.6 %). Based on the regional government of Tigray geological map, the study watershed is classified into Alaje formation.

2.2. Climate

A large part of this district is in the mid highland zone (80% of area), with the lowland (500-1500 masl) and highland (2300-3200 m.a.s.l.) areas covering 18.6% and 1.4% of the district, respectively. Rainfall is bimodal, but potential evaporate-transpiration is high and exceeds rainfall amount in most of the years (Weldeamlak et al. 2015). According to the metrological data of Tigray regional state, the study district's mean annual rainfall is 690 mm, and the mean monthly temperature varies from 12 to 28.8 oC (Figure 2).




Figure 2 metrological data of the study watershed

According to FAO (2006) land slope classification, the watershed was classified into seven slope types: flat (<3%), gentle sloping (3–8%), rolling or sloping (8–15%), hilly (15–30%), mountainous (30–40%), steep mountainous (40–60%), and very steep mountainous (>60%). Therefore, the slope in the study watershed varies from flat to very steep mountainous (Figure 3).




Figure 3 Land Slope classification of the study watershed

2.3. Study methods

2.3.1. Site selection

The study watershed was selected in consultation with the district watershed experts and on field survey following the pre-developed criteria, such as the existence of different land uses, various slope classes/landforms, and the existence of different agro-ecologies and watersheds with an age of two decades or more. As shown in the photo below, the left side and the right side indicate the lower and upper parts of the watershed, respectively (Figure 4).



Figure 4 Photo showing the study Watershed in part (photo: Abadi, 2017)

According to FAO's (2006) landform classification, the Watershed is classified into three parts (landforms) which are the upper part, middle part, and lower part. The entire Watershed contains four land use types.

2.3.2. Data collection and analysis

Land use and land cover data were obtained from Landsat images for the years 1998 TM/Landsat 5 & 8 and 2017 ETM+/Landsat 7 for December. In this month all crops are harvested, and the average monthly rainfall of those years is almost the same. The images were acquired from the United States Geological Survey (USGS) via <http://earthexplorer.usgs.gov/>. The year 1998 was selected because it is the year when an integrated watershed approach was introduced to the Tigray region (Irish Aid, 2006). Land use and land cover types were described based on the land use and land cover description guide proposed by Teka et al. (2014) (Table 1).

Table 1 Land use and land cover description

Land use and land cover classes	Each land use class description
Forest land	Patches with a canopy cover over 50%. These can be remnants of high natural forest, found in small patches around churches, on steep slopes and less populated areas, or recent eucalyptus plantations.
Cultivated land	Annual and perennial crops (>70% of the land), frequently observed on level lands (plains, plateaus, foot slopes, and valley floors)

Bushland	Dominant vegetation lower than 3 m but higher than 1 m with a canopy cover above 15%, or dominant vegetation below 1 m with a canopy cover above 50%
Shrub land	Dominant vegetation lower than 1 m, with a canopy cover below 50% but higher than 5%
Homestead	Residences, administrative buildings, and small industrial, and trade enterprises.

Image pre-processing and post-classification were done to correct errors (Figure 3). A total of 430 ground control points (GCPs) were collected with the help of a global positioning system (GPS). These (GCPs) were grouped into two, in which 50% was used for classification by the earth resources data analysis system (ERDAS-imagine v 2014) and the remaining for validation. Land cover maps of the different years were created with the help of ArcGIS software 10.2, and change detection was also done to see the temporal and spatial variations.

The rate of land use and land cover change was calculated for each land use and land cover using the following formula.

$$\text{Rate of change} \left(\frac{\text{ha}}{\text{yr}} \right) = \frac{\text{final study} - \text{initial study}}{\text{time interval}} \text{-----Equation 1}$$

Based on this formula rate of land use land cover change between 1998 and 2017 year was calculated. This method was used by several authors like Abate S. and K.L. Singh (2011).




Figure 5 Representation diagrams for analyzing land use and land cover dynamics in the years 1998 and 2017

2.3.3. Accuracy Assessment

For this study, the producer's accuracy, user's accuracy, omission error, commission error, overall accuracy, and kappa coefficient were assessed to validate the reliability of the land use land cover classification based on pixel-to-pixel comparison for the year 2017 LULC categories. The overall accuracy, kappa coefficient, user's accuracy, and producer's accuracy were mathematically estimated as the following questions:

$$\text{User's accuracy} = \frac{A_{ii}}{(A_{i+})} \dots \dots \dots \text{Equation 2}$$

$$\text{Producer's accuracy} = \frac{A_{ii}}{(A_{i+})} \dots \dots \dots \text{Equation 3}$$

$$\text{Overall accuracy} = \frac{C}{D} \dots \dots \dots \text{Equation 4}$$

Where A_{ii} = total number of observations in row i and column i , A_{i+} and A_{+i} are the marginal totals of row i and column i , respectively. N = total number of observations, C = total number of correct pixels, and D = total number of pixels in the error matrix.

In this study, the land use and land cover change matrix was calculated from the years 1998 to 2017. The overall accuracy value was expressed by the sum of the diagonal pixels converted into a percentage of the area (Paegelow et al. 2014). And also Fan et al. (2008) also express how many measurements have been proposed to improve the interpretation of the change matrix. However, the kappa coefficient is one of the universally used to find the accuracy assessment techniques. In this study, overall accuracy, kappa coefficient, omission error, and producer's accuracy results were calculated.

Table 2 Level of the kappa coefficient

Kappa coefficient (k_{hat})	Level of agreement/ accuracy
>0.80	Strong
0.40-0.80	Medium
<0.40	Poor

Source (Lillesand and Kiefer, 1994)

3. RESULTS AND DISCUSSIONS

3.1. Spatial and temporal land use and land cover detection

In the study Watershed, five major land use and land cover types were identified (Figure 8). The classified land use and land cover types are forests, bushland, shrubland, cultivated land, and homesteads. Those land use and land cover types are mostly dominant in the study watershed.




Figure 6 Land cover map of the study periods

Table 3 Land covers classes of coverage in each date of study image

Land cover type	Land cover in each year image of 1998 and 2017			
	1998		2017	
	ha	%	ha	%
Forest	819	32.3	730.3	28.8
Cultivated	509	20.1	464.5	18.3
Bushland	776	30.6	727.7	28.7
Shrub land	272	10.7	381.1	15
Homestead	160	6.3	232.4	9.2

Total	2536	2536
-------	------	------

According to the above table, only about 20% and 18% of the land cover were under cultivation in the years 1998 and 2017, respectively. This reduction occurred due to the implementation of exclosures and land rehabilitation programs. Therefore, massive areas were protected to improve the vegetation coverage during this time. Eskinder et al. (2017) reported lined with this result, which is cropland coverage declined for the year between 1984 and 1995 by 21.4%. As a result of the unwise use of forest and illegal construction of resettlement, land use coverage of forest, bush, and cultivated showed a decline of 3.5%, 1.9%, and 1.8%, respectively. A rapid reduction in woodland from 8.35% to 0.28% and grassland from 7.75% to 0.91% cover in the landscape took place between 1972 and 2007 (D. Tsegaye et al., 2010).

3.2. The extent and rate of land use and land cover change

As shown in Table 4, shrub and homestead land cover increased from 10.7% to 15% and 6.3% to 9.2%, respectively. The increment of shrubland is from bushland and forest land directly and indirectly. Because the bushland and forest land are used by the population for economic aspects, construction, and agricultural instruments. On the other hand, the increment of homestead changed directly from the cultivated land due to new resettlement built after the single marriage. Most of the resettlement construction of the study area is done on the cultivated land. Due to this reason, the size of cultivated land and forest land decreased from 1998 to 2017 by 8.7% and 10.8% respectively (Table 4). Also, bushland cover type decreased from the year 1998 to 2017 by 6.2%. Terefe et al. (2016) reported that, similar to this finding, forest cover decreased by 14.7% between 1973 and 1986 and by 38.5% between 1986 and 2001. But, contradictory to the result of shrub land, which is shrub lands were also reduced by 25.9% from 2001 to 2015 because, during the study period, shrub land was mainly converted to cultivation and settlement. The land use and land cover change were also mapped (Figure 6).

Table 4 Changes in land use and land cover from 1998 to 2017

LULC	1998		2017		LULC changes b/n 1998 and 2017		Rate of change Ha/yrs.
	Area (ha)	%	Area (ha)	%	Area (ha)	%	
Forest	819	32.3	730.3	28.8	-88.7	-10.8	-4.4
Cultivated	509	20.1	464.5	18.3	-44.5	-8.7	-2.23
Bushland	776	30.6	727.7	28.7	-48.3	-6.2	-2.42
Shrubland	272	10.7	381.1	15	109.1	40.1	5.46
Homestead	160	6.3	232.4	9.2	72.4	45.3	3.62
Total	2536	100	2536	100	-	-	-

Where he stands for hectares and yrs. is also for years.

The land use and land cover change matrix results show the direction of change and its remains. Therefore, the change matrix of the land use land cover is depicted in (Table 5) which is columns that represent the initial study year categories and rows that contain the final study year categories.




Figure 7 Map of land use and land cover change

Table 5 Summary of LULC change matrix in ha from 1998 to 2017.

From TO	Forest land	Cultivated land	Bushland	Shrub land	Homestead	Total (2017) ^b
Forest land	730	0	0	0	0	730
Cultivated land	0	432.5	20	0	12	464.5
bushland	59.2	61.5	572	15	20	727.7
shrub land	14	15	89	253.1	10	381.1
Homestead	15.8	0	95	4	118	232.4
total(1998) ^a	819	509	776	272	160.3	2536

The total sum of the rows is the amount of land cover for each LULC in the initial study year (1998). The total sum of the column is also the amount of land cover for each LULC in the final study year (2017). The amount of changed land from one land use and land cover into another LULC is listed in each cell of the above table. Similar to this finding, Abate et al. (2011) reported that the forest land changed into shrubland and bushland. While shrubland changed into bushland and cultivated land.

3.3. Level of Accuracy assessment

The Overall accuracy, kappa coefficient, user accuracy, and producer's accuracy for the year 2017 were estimated in the following table.

Table 6 Level of kappa coefficient of agreement

LULC 2017	Reference data						Classification accuracy (%)	
	FL	CL	BL	SL	HS	Total	User's accuracy	Errors of commission
FL	46	0	3	1	0	50	92	8
CL	0	51	0	2	1	54	94.4	5.6
BL	3	4	45	3	0	55	81.8	18.2
SL	1	5	2	33	1	42	78.6	21.4
HS	0	0	0	1	13	14	92.9	7.1
Total	50	60	50	40	15	215	-	60.3
PA	92	85	90	82.5	86.7			
EO (%)	8	15	10	18	13.3			
OA (%)	87.4							
KC	0.84							

Where PA = producers' accuracy, EO = error of omission, OA= overall accuracy and KC = kappa coefficient

4. CONCLUSIONS AND RECOMMENDATIONS

In this study, LULC detection of Woji watershed over the last two decades was analyzed. The result of the study showed that significant change detection had been observed during the study period. Shrub land and residential/homestead areas showed an increasing trend of 5.46 and 3.62%, respectively, while forest, bush land, and cultivated areas showed a decreasing trend of 4.4, 2.42, and 2.233%, respectively. This result discovered that the change of forest, bush land, and cultivated land to shrub and residential/homestead areas may cause problems, including changes in stream flow, soil degradation, and hydrological systems in the watershed. These changes could have implications for sustainable resource management and the livelihood of the local society. Therefore, developed land management practices (soil and water conservation), reinforced agricultural inputs, integrated watershed management (land use planning and management), and active participation of the local community should be advanced to prevent undesirable LULC dynamics in the watershed. In this study, the change detection analysis using GIS and remote sensing could deliver useful information to understand the seasonal patterns of land use dynamics for planners and decision makers; consequently, sustainable land management planning is possible.

Based on the study the following points are recommended:

- Forest land is highly decreased while the correspondent shrub land and homestead is increased in the last two-decade years (1998-2017) and this has to be needed preserve the forest area through protection and afforestation.
- Increasing forest cover through tree plantation and natural regeneration should be encouraged and deforestation must be prohibited in all watersheds of Tigray region.

Reference

- Abate Shiferaw and K.L. Singh, (2011). Evaluating Land Use and Land Cover Dynamics in Borena *Woreda*, South Wollo Highlands, Ethiopia. *EJBE* 2(1); 64-104.
- Alemayehu Kiflu, (2007). Effects of different land use systems and topography on soil properties at Delbo watershed, Wolaita zone. MSc Thesis, Awassa University, Ethiopia. 64p.
- Bin Q, Jian-Fei CHEN, Hong-Lie QIU, Römken MJM, Xiao-Qi YANG, Jiang SF, Bi-Cheng LI (2006) Spatial-temporal pattern and driving forces of land use changes in xiamen11project supported by the Fujian Provincial Natural Science Foundation of China (No. D0210010). *Pedosphere* 16(4):477–488
- Deng X (2011). Modeling the dynamics and consequences of land system change. Springer, Berlin, pp 257–261
- Diress Tsegaye, Stein R. Moe, Paul Vedeld and Ermias Aynekulu, (2010). Land-use/cover dynamics in Northern Afar rangelands, Ethiopia, Norwegian University of Life Sciences, 174-180
- Food and Agriculture Organization (FAO) (Eds) (2006). Guidelines for soil description. The United Nations, Rome.
- Gete Zeleke and H. Hurni (2001). Implications of land use and land cover dynamics for mountain resources degradation in the north-western Ethiopian highlands. *Mountain Research and Development* 21(2): 184-191.
- Gidey, E., Oagile D., Reuben S. Eagilwe S. and Amauel Z., (2017). Modeling the spatio-temporal dynamics and evaluation of land use and land cover (1984-2015) using remote sensing and GIS in Raya, Northern Ethiopia. p. 1-17
- IRIS AID. (2006). Evaluation of the Irish Aid/Tigray Regional support program (TRSP).
- Kebrom Tekle and L. Hodlund. (2000). Land cover changes between 1958 and 1986 in Kalu district, southern Wollo, Ethiopia. *Mountain Research and Development* 20(1): 42-51.
- Kassa T., Anton V., Jean P., Simon V., Jozet D., Kassa A. (2014). Spatial Analysis of Land Cover Changes in Eastern Tigray (Ethiopia) from 1965 to 2007, *Earth and Environmental Sciences*, KU Leuven, Heverlee, Belgium
- Lambin, F.E., Turner, B.L., Geist, J.H. (2001). The causes of land-use and land-cover change: moving beyond the myths. *Global Environmental Change*, 11: 261–269.
- Mas JF, Kolb M, Paegelow M, Olmedo MTC, Houet T (2014) Inductive pattern-based land use/cover change models: a comparison of four software packages. *Environ Model Software* 51:94–111
- Rawat JS and Kumar M (2015) Monitoring land use/cover change using remote sensing and GIS techniques: a case study of Hawalbagh block, district Almora, Uttarakhand, India. *Egypt J Remote Sens Space Sci* 18(1):77–84
- Tefera T., Feyera S. and Mogos K. (2016). Landscape composition and configuration in the central highlands of Ethiopia.
- Turner, B.L, D. Skole and S. Sanderson (1995). Land-use and land-cover change: Science/Research plan. IGBP Report 35 / HDP Report 7. IGBP Press Stockholm and Geneva. 165.
- Turner, B.L., Lambin, E.F., Reenberg, A. (2007). The emergence of land change science for global environmental change and sustainability. *Proc. Natl. Acad. Sci. U.S.A.* 104, 20666–20671.
- Woldeamlak Bewket and L. Stroosnijder (2003). Effects of agro-ecological land use succession on soil properties in the Chemoga watershed, Blue Nile basin, Ethiopia. *Geoderma*. 111: 85-98.
- Yifru A, Taye, B. (2011). Effects of land use on soil organic carbon and nitrogen in soils of Bale, South-eastern Ethiopia. *Tropical and Subtropical Agro ecosystems*, 14:229 – 235.
- Yimer, F., Ledin, S., Abdulakdir, A. (2007). Changes in soil organic carbon and total nitrogen contents in three adjacent land use types in the Bale Mountains, southeastern highlands of Ethiopia. *For Ecol. Manag.*, 242: 337–342.