

## Original Research Article

### Evaluation of Nutritional Composition of *Ocimum gratissimum* Leaf Extract

#### ABSTRACT

Vegetables are known to be vital sources of nutritive foods. This study evaluates the nutritional composition of ~~Ocimum~~ *Ocimum gratissimum* leaf extract vis-a-vis proximate minerals and vitamins analyses. Fresh leaves of *Ocimum gratissimum* were purchased from a local market in Awka North Local Government Area of Anambra State, Nigeria and were washed, destalked, oven-dried at 40 °C, pulverized and sieved through 2.0 mm mesh prior to analysis. Proximate, minerals, and vitamin analyses were carried out, using standard methods. Results indicated that *Ocimum gratissimum* contained high levels of carbohydrate (51.78 ± 0.50 %), protein (10.50 ± 0.04 %), and fiber (9.38 ± 0.05 %), with minute quantity of crude fat (4.94 ± 0.03 %). Analysis of the mineral compositions showed that the vegetable contained considerable quantity of minerals that is sufficient to improve general well-being, while the vitamin analysis revealed that the leaves of *Ocimum gratissimum* was sufficiently rich in Vitamin E and could therefore act as antioxidant and consequently help to improve health status of its consumers.

**Keywords:** Green leafy vegetables, *Ocimum gratissimum*, nutrition, health,

Formatted: Font: Italic

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

## INTRODUCTION

Green leafy vegetables are widely succulent plants parts grown in gardens and consumed as a side dish or soup with starchy staples among the tribes in Nigeria [1].

The importance of dietary components of leafy vegetables is significant in African

population since they are comparatively rich in fiber, ~~whereas~~ cereals, root vegetables, and other food ~~stuff~~ are relatively poor sources [2]. ~~In Nigeria, just as in~~

~~most~~ Similar to other African countries, a typical Nigerian ~~where~~ diet is dominated by starchy staple foods, ~~and~~ vegetables are considered ~~the~~ cheap source of energy. ~~They~~

~~Vegetables~~ are ~~very~~ major ~~rich~~ sources of nutrients such as carotene, protein, vitamins, calcium, iron, ~~and~~ ascorbic acid, ~~and~~ they also contain tangible concentration of trace minerals [3].

Many species of vegetables abound in the world, and they are ~~characterized~~ utilized in ~~with~~ addition of flavor, taste, and color to diet which would ~~otherwise~~ have otherwise been ~~monotonous~~ be monotonous. The invaluable food nutrients contained in vegetables can be used for body ~~building~~ growth, protective, and regulatory ~~material~~ roles as well as sources of energy for the body.

Vegetables also contain plants secondary metabolites which act as anti-nutrients [4].

These secondary metabolites include: tannins, phytate, oxalate, and cyanide; all of which have ~~different~~ various detrimental ~~al~~ effects to nutrition and health [5]. A good leafy vegetable is the one that contains ~~a~~ ~~great~~ ~~high~~ ~~deal~~ amount of these beneficial nutrients and a very minute amount of anti-nutrients. One of the promising vegetables that has been acclaimed 'good' is *Ocimum gratissimum*.

*O. ~~cinum~~ gratissimum* Linn (Lamiaceae) is a ~~perennial~~ herbaceous shrub notably found in tropical countries, including Nigeria [6], where it is commonly called Clove

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

basil, Sweet basil, tea-bush, Scent leaf or fever plant, ~~but~~ it is also popularly known with different local names in Nigeria as “*Daidoya*” (hausa), “*Effirin*” (yoruba), and “*Nchuanwu*” (igbo).

~~*Ocimum gratissimum* is a widespread and commercially viable perennial herbaceous plant with a~~Due to its very strong aromatic smell, ~~it~~ is used as a natural ~~flavouring~~flavoring agent, condiment, or vegetable in the preparation of fish, meat, soup, and stew.

Scientific reports have shown that *O. gratissimum* has a wide range of bio-active compounds such as flavonoids, ~~and~~ polyphenols, [7] and essential oils with several beneficial effects [8]. Since the ~~concentrations of the~~ nutrient ~~contents~~ s vary from one leafy vegetable species to another, there ~~is a~~ is a need to investigate these ~~variations~~ ~~se~~ ~~varying concentrations in within~~ the vegetables. Hence, this study ~~aimed to evaluate~~ ~~evaluate~~ the nutritional composition of ~~*O. eimum*~~ *gratissimum*.

## MATERIALS AND METHODS

### Plant Materials

Fresh leaves of ~~*O. eimum*~~ *gratissimum* were purchased from ~~a the~~ daily market, Mgbakwu in Awka North Local Government Area of Anambra State, Nigeria and were authenticated by a Taxonomist, Mr Iroka Finan, ~~of from~~ the Department of Botany, Nnamdi Azikiwe University, Awka. ~~A, and a~~ voucher specimen was ~~also~~ deposited with Herbarium no NAUH 35B. The leaves were then detached from the stalk, rinsed with distilled water, ~~and~~ ~~were~~ oven-dried at 40 °C. The dried samples were ground into fine powder and sieved through 2.0 mm mesh prior to analysis.

### Proximate Analysis

Moisture, ash, crude fat, and crude fiber were analyzed in the fresh samples in accordance with the methods described by Association of Official Analytical Chemist

[9], while [the percentage of](#) nitrogen was determined by the micro-Kjeldahl method (1976) and ~~the percentage of was nitrogen was~~ converted to crude protein by multiplying by 6.25. Carbohydrate was determined by difference [10].

### Vitamin Analysis

Vitamin A and C were determined by the Fütter-mayer colorimetric method of Kirk and Sawyer [11], while Vitamin E was determined using the Emmerie-Engel reaction for Tocopherol estimation as reported by Rosenberg [12].

### Mineral Analysis

Wet ashing method was used ~~in releasing to release~~ the investigated metals in the samples from biological complexes before evaluating them with atomic absorption spectrophotometer (PinAAcle™ 900H) and flame photometer (Gulfex FP6410). ~~The~~ ~~W~~et ashing was achieved by heating the sample with a digestion mixture containing concentrated nitric, perchloric, and sulphuric acids in a Kjeldahl flask [9]. Calcium, Iron, Manganese, Magnesium, ~~and~~ Zinc were determined by atomic absorption spectrophotometer; Sodium and Potassium by Flame photometer while Phosphorus was determined by Molybdate method using hydroquinone as a reducing agent [9].

### Data Analysis

~~The~~ ~~mean?~~ values obtained and the significance between the treated and control group was analyzed by one-way ANOVA using the SPSS version 17 and  $P < 0.05$  was considered to be statistically significant.

Formatted: Highlight

Formatted: Font color: Red, Highlight

Formatted: Highlight

## RESULTS AND DISCUSSION

Table 1.0 showed the result for phytochemical analysis of the leaf extract of *O. cinereum gratissimum*. Result showed that the leaf extract contained high levels of Steroids ( $81.91 \pm 14.16$  ug/g), Flavol-3-ol ( $81.88 \pm 9.14$  ug/g), Anthocyanin ( $67.13 \pm 6.63$  ug/g), and Phenol ( $50.72 \pm 6.07$  ug/ml), with considerable amount of Sapogenin ( $26.36 \pm 5.68$

ug/g), Quinine (19.95±5.11 ug/g), and Lunamarin (14.64±0.15 ug/g), while Proanthocyanin 0.42±0.03 ug/g, Naringenin (0.70±0.03 ug/g) and Epihedrine (1.55±0.51 ug/g) are were found in trace quantities.

**Table 1.0: Table showing a average phytochemical composition of the leaf extract of *O. cimum-gratissimum***

Parameters	Quantity
Proanthocyanin <del>*(ug/g)</del>	0.42±0.03
Rutin <del>*(ug/g)</del>	6.02±1.56
Ribalidine <del>*(ug/g)</del>	4.90±0.00
Quinine <del>*(ug/g)</del>	19.95±5.11
Flavol-3-ol <del>*(ug/g)</del>	81.88±9.14
Anthocyanin <del>*(ug/g)</del>	67.13±6.63
Lunamarin <del>*(ug/g)</del>	14.64±0.15
Sapogenin <del>*(ug/g)</del>	26.36±5.68
Epihedrine <del>*(ug/g)</del>	1.55±0.51
Phenol <del>^(ug/ml)</del>	50.72±6.07
Flavonones <del>*(ug/g)</del>	6.04±1.14
Steroids <del>*(ug/g)</del>	81.91±14.16
Epicatechin <del>*(ug/g)</del>	21.02±3.12
Kaepferol <del>*(ug/g)</del>	2.97±0.07
Phytate <del>*(ug/g)</del>	1.15±0.17
Oxalate <del>*(ug/g)</del>	5.61±0.62
Resveratol <del>*(ug/g)</del>	5.53±0.54
Flavones <del>*(ug/g)</del>	3.32±0.22
Naringenin <del>*(ug/g)</del>	0.70±0.03

Tannin  $\mu\text{g/g}$  12.39 $\pm$ 2.85

Values are expressed in mean $\pm$ SEM: \* ( $\mu\text{g/g}$ ), ^ ( $\mu\text{g/mL}$ )

Table 2.0 showed the result for proximate analysis of the leaf extract of *O. eimum gratissimum*. Result indicated highest content of Carbohydrate (51.78  $\pm$  0.50%), with minimal content of crude fat (4.94  $\pm$  0.03%).

**Table 2.0: The percentage proximate composition of the leaf extract of *O. eimum gratissimum***

Formatted: Font: Bold

Proximate components	Percentage composition (%)
Moisture	11.92 $\pm$ 0.02
Ash	11.48 $\pm$ 0.10
Fiber	9.38 $\pm$ 0.05
Crude fat	4.94 $\pm$ 0.03
Protein	10.50 $\pm$ 0.04
Carbohydrates	51.78 $\pm$ 0.50

Values are expressed in mean $\pm$ SEM

Values are mean $\pm$ SEM

Table 3.0 showed the result for vitamin analysis of the leaf extract of *O. eimum gratissimum*. Result indicated that the sample was rich in vitamin E (35.85 $\pm$  1.56  $\mu\text{g/g}$ ), with minimal content of vitamin A (1.86 $\pm$  0.07 $\mu\text{g/g}$ ).

**Table 3: The vitamin composition of the leaf extract of *O. eimum gratissimum***

Vitamins	Concentration ( $\mu\text{g/g}$ )
A	1.86 $\pm$ 0.07

C	3.40 ± 0.00
E	35.85 ± 1.56

Values are expressed in mean±SEM

~~Values are mean±SEM~~

Table 4.0 showed the result for mineral composition of the leaf extract of *O. eimum gratissimum*. Result indicated that the sample contained considerable quantity of iron (1.40 ± 0.50 ppm) with minute quantity of copper (0.29 ± 0.17ppm).

**Table 4:** ~~0: The~~ **Mineral composition of the leaf extract of *O. eimum gratissimum***

Minerals	Quantity (ppm)
Copper (ppm)	0.29 ± 0.17
Iron (ppm)	1.40 ± 0.50
Manganese (ppm)	0.31 ± 0.08
Zinc (ppm)	0.79 ± 0.10
Selenium (ppm)	0.37 ± 0.02

Values are expressed in mean±SEM

~~Values are mean±SEM~~

The results of the proximate analysis of the leafy vegetable ~~is was~~ represented in Table 2.0. Generally, the results indicated that the vegetable ~~was~~ a good source of carbohydrates protein, and crude fiber. Carbohydrate content was much higher (51.78 ± 0.50 %) compared to other macronutrients. The high Carbohydrate content of *O. eimum gratissimum* ~~is was~~ in accordance with the reports of Hussain *et al.* [13] and Oladejo [14]. *O. eimum gratissimum* ~~meets~~ the requirement of providing by

Formatted: Font: Not Italic

providing more than 12% of its calorific value from protein and could, thus, be considered a good source of protein according to Ajayi *et al.*; [1]. However, The high protein content of *O.cimum gratissimum* however, does not present it as a good sourcegood source of protein as the quality of a protein is dependent on its ability to supply all the essential amino acids in the required amount [15];hHence, protein quality assessment is necessary to ascertain the ability of *O.cimum gratissimum* to meet dietary protein needrequirements.

Formatted: Font: Italic

Formatted: Font: Italic

The microelements: copper, iron, manganese, and zincand zincare were also well represented in the vegetable. Although the Dietary Reference Intake (DRI) of 20.5 mg of iron cannot be met by the vegetable at 100g per day serving, the vegetable is nevertheless a good source of iron as supplementation with other food components will make up for the iron need. Epidemiological studies have indicated that diets high in iron can help to fight infections, boost brain health, and also promote systemic circulation of oxygen [14]. Manganese is an essential metal because it is required for proper immune function, regulation of blood sugar and cellular energy, reproduction, digestion, bone growth, blood coagulation, and homeostasis as well as defense against reactive oxygen species [16]. The beneficial effects of manganese are due to the incorporation of the metal into metalloproteins, thus aiding biochemical functions. The minute quantity of manganese ( $0.31 \pm 0.08$  ppm) present in this leafy vegetable indicates that it could provide the body with the minimum concentrations of manganese required for basic enzyme activities.

Similarly, the zinc concentrations present in the vegetable as observed in this study were found to be comparable with previous works [1, 14]. Zinc is highly recommended in diets as it is required for proper sexual organ functioning and

enzyme activity. High levels of zinc are usually associated with high-protein food stuff, whereas low levels are obtained from food rich in carbohydrates [17].

Vitamins are organic compounds that are essential to an organism in small quantities for proper metabolic function. As depicted in this study, *O. eimum gratissimum* contained considerable quantity of vitamins, especially vitamins A, C, and E. Vitamin A is a fat-soluble vitamin that is naturally present in many foods. It is important for normal vision, immune system, reproduction, growth, and development. It has also been reported to help in the proper functioning of the heart and lungs [16].

The vitamin A content as reported in this study ~~is~~ was consistent with the findings of Oladejo [14] and Enemor *et al.*, [17]. In the same vein, the researched vegetable ~~is~~ was found to be rich source of vitamin E ( $35.85 \pm 1.56 \mu\text{g/g}$ ). Vitamin E is a fat-soluble vitamin that act as antioxidant and consequently help to protect the cells from damages occasioned by free radicals. By implication, consumption of this vegetable could aid the body's antioxidant defense mechanism and consequently improve health.

## CONCLUSION

This study indicated ~~s~~ that the investigated vegetable ~~is~~ was a rich source of carbohydrate, protein, fiber, and essential minerals, as well as vitamins A, C, and E. *O. eimum gratissimum* presented a nutritive profile capable of promoting health and well-being. The anti-nutrients in the leafy vegetable ~~are~~ were much lower than the safe limits and the reports for other plant sourced foods, including nuts, roots, beans and berries. A serving of 100 g per day of the investigated vegetables, complemented by other nutrient sources, will provide the RDA of the essential minerals investigated. While a serving of up to 2 kg per day of the vegetables does not contribute anti-nutrients higher than the safe limits.

Formatted: Font: Italic

Formatted: Font: Italic

### REFERENCES

1. Ajayi, O. B., Bamidele, T. J., Malachi, O. I. and Oladejo, A. A. (2018). Comparative Proximate, Minerals and Antinutrient Analysis of Selected Nigerian Leafy Vegetables. *Journal of Applied Life Sciences International*, 16(1): 1-8.
2. Okonwu, K., Egerton, N. A. Ajayi, A. E. and Agobua, J. U. (2022). Nutritional Composition of Some of the Most Consumed Cereals and Legumes in Nigeria. *European Journal of Nutrition and Food Safety*, 14 (2):53-58.
3. Nakano M. R., Umar B. D., Nuhu A. L. and Adamu I. T. (2023). Proximate and mineral composition of some wild leafy vegetables consumed in Katsina State Nigeria. *World Journal of Advanced Engineering Technology and Sciences*, 09(01):084–088.
4. Baiyeri, S.O., Chimaluka, C.A., and Okorie, O. N. (2023). Proximate, Mineral, Vitamin, and Anti-Nutrient Contents of the Leaves of *Senecio biafrae*. *Journal of Agricultural Sciences*, 68(1):67-79.

5. Ojmelukwe, P.C. and Amaechi, N. (2019). Composition of *Vernonia amygdalina* and its potential health benefits. *International Journal of Environment, Agriculture and Biotechnology*, 4 (6):1836-1848.
6. Akara, E.U., Okezie E., Ude V.C., Uche-Ikonne C., Eke G., Ugbogu A.E. (2021). *Ocimum gratissimum* leaf extract ameliorates phenylhydrazine-induced anaemia and toxicity in Wistar rats. *Drug Metab. Pers. Ther.* 15(1):20-35.
7. Irondi E.A., Agboola S.O., Oboh G. and Boligon A.A. (2016). Inhibitory effect of leaves extracts of *Ocimum basilicum* and *Ocimum gratissimum* on two key enzymes involved in obesity and hypertension *in vitro*. *J. Intercult. Ethnopharmacol.* 5:396–402.
8. Melo R.S., Azevedo A.M.A., Pereira A.M.G., Rocha R.R., Cavalcante R.M.B., Matos M.N.C., Lopes P.H.R., Gomes G.A., Rodrigues T.H.S., Dos Santos H.S., Ponte I.L., Costa R.A., Brito G.S., Júnior F.E.A.C., Carneiro V.A. (2019). Chemical composition and antimicrobial effectiveness of *ocimum gratissimum* L. Essential oil against multidrug-resistant isolates of *Staphylococcus aureus* and *Escherichia coli*. *Molecules.*3(8)64:2–17.
9. A.O.A.C. Official method of analysis. Association of Official Analytical Chemist (1990);15:910-28.
10. McDonald P, Edwards RA, Greenhalgh JFD. Animal Nutrition. Edinburgh: T and A Constable Ltd. 1973: 2-5.
11. Kirk, R.S and Sawyer, (1991). Pearsons Food Composition and Analysis. 4<sup>th</sup> Edition. Macmillan Publisher Company, United Kingdom.
12. Rosenberg, H.R. (1992). Chemistry and physiology of the vitamins. Interscience Publisher, New York. P 452-3.

13. Hussain, J., Rehman, N.U., Khan, A.L., Hamayun, M., Hussain, S.M. and Shinwari, Z.K. (2010). Proximate and essential nutrients evaluation of selected vegetables species from Kohat region, Pakistan. *Pak. J. Bot.* ;42(4):2847-2855.
14. Oladejo A.A. (2019). Comparative Vitamin Analysis of Some Selected Nigerian Green Leafy Vegetables from Two Different Zones". *Acta Scientific Nutritional Health* 3(11): 1-7.
15. Pellett, P. L. and Young, V. R. (2018). Nutritional evaluation of protein foods. *Food and Nutrition Bulletin.* 4(1):1-7.
16. Nnenne, S.K., Ubaaji, K.I., Ogbodo, U.C., Enemor, V.H.A. and Oladejo, A.A. (2020). Comparative Study on the Nutritional and Antioxidant Components of Fruit Parts of *Citrullus lanatus*. *European Journal of Nutrition and Food Safety*, 12(11): 39-51.
17. Enemor, V.H.A., Oguazu, C.E., Odiakosa, A.U. and Okafor, S.C. (2019). Evaluation of the medicinal properties and possible nutrient composition of *Citrullus lanatus* (watermelon) seed. *Res J Med Plants.* 13:129-135.