

Nanotechnology-Enhanced Phytonutraceuticals: Innovations in Characterization, Formulation, and Therapeutic Applications

ABSTRACT

Nanotechnology refers to the branch of science and engineering devoted to designing, producing, and using structures, devices, and systems by manipulating atoms and molecules at nano-scale, having one or more dimensions of the order of 100 nanometres (100 millionth of a millimetre) or less. Nanotechnology is the exploitation of the unique properties of materials at the nanoscale. Nanotechnology has gained popularity in several industries, as it offers better built and smarter products. The ability to manipulate structures at the atomic scale allows for the creation of nanomaterials. Nanomaterials have unique optical, electrical and/or magnetic properties at the nanoscale, and these can be used in the fields of electronics and medicine, amongst other scenarios. Nanomaterials are unique as they provide a large surface area to volume ratio. Unlike other large-scaled engineered objects and systems, nanomaterials are governed by the laws of quantum mechanics instead of the classical laws of physics and chemistry. Nanotechnology is the engineering of useful objects and functional systems at the molecular or atomic scale. Nano-formulations are widely used for phyto-nutraceuticals and drug delivery systems. These substances are generally having low solubility, leading to their poor absorption and bioavailability in the human body. In this regard, one of the most important applications of nanotechnology in food sector has been the formulation of novel nutraceutical compounds with improved properties viz., enhanced solubility, stability, bioavailability and efficacy. This is achieved by encapsulation of nutraceutical by nanoparticles, which modifies their pharmacokinetics (PK) and biodistribution (BD). Nano-formulations widely used for the purpose are nanoliposomes, nanoemulsions, nanoparticles, nanofibres. The particles are characterized by scanning probe microscope (SPM), Ultraviolet-visible spectroscopy (UV-Vis spectrophotometer, SEM, TEM, Dynamic light scattering (DLS).

Key words: Bioavailability, Characterization, Nanoformulations, Nutraceutical delivery, Phytonutraceuticals

Introduction

The prefix "nano" in the term *nanotechnology* denotes a factor of one billionth (1×10^{-9}). Nanotechnology involves the study and manipulation of materials at the nanoscale, with structural dimensions typically on the order of 1 nanometer ($1 \text{ nm} = 10^{-9}$ meters). Nanoparticles are commonly defined as aggregates of atoms or molecules, exhibiting sizes with a radius less than 100 nm. At this scale, unique physical, chemical, and biological properties emerge, primarily due to quantum effects and the increased surface area-to-volume ratio. The growing interest in nanoscience (the study of low-dimensional systems) is a realization of Richard Feynman's renowned statement, "There's Plenty of Room at the Bottom." Building upon Feynman's vision, K. E. Drexler introduced the concept of "molecular nanotechnology" in his 1986 book *Engines of Creation*, where he proposed the use of nanoscale molecular structures to function in a mechanistic capacity, directing and catalyzing the synthesis of larger molecules.

In contrast to conventional materials, which typically have grain sizes ranging from hundreds of micro-meters (μm) to millimeters (mm), nano-crystalline materials possess grains in the range of 1–100 nanometers. While bulk materials largely retain their properties when reduced to the micro-meter scale, materials at the nano-scale exhibit markedly distinct and novel physical, chemical, and mechanical properties. These changes are primarily attributed to the increased surface area-to-volume ratio and quantum effects that dominate at reduced dimensions.

“According to National science Foundation, Nanotechnology is the capability to understand, manipulate and control matter at the level of individual atoms and molecules .Science and engineering are the primary operators of global technological competition. Modern science based on the unifying features of nature at the nanoscale contributes a new foundation for innovation, knowledge, and integration of technology.Nanotechnology is sometimes preferred as a general purpose technology because in its advanced version it will have significant impact on almost all areas of society and all industries.There is a longitudinal process of convergence and divergence in extensive areas of engineering and science. For example the convergence of sciences at macro scale was intended during the Renaissance, and it was latterly followed by narrow disciplinary specialization (NDS) in science and engineering in the 18th-19th centuries. The convergence at the nanoscale reached its brawn in about year 2000, and an estimation of a divergence in the nano system architectures in the next decades” (Anonymous, 2009).

Distinctive Characteristics of Nanotechnology

“Nanoparticles, as implied by the term, are characterized by their diminutive size. Due to this reduced scale, nanoparticles possess an exceptionally high surface area-to-volume ratio when compared to bulk materials. This significant increase in surface area enhances their reactivity, interaction potential, and other surface-dependent properties, which are often absent or less pronounced in larger material forms. This feature enables them to possess different physical, chemical and optical properties as they have small space to confine their electrons and produce quantum effects. Nanoparticles are highly reactive because of their large surface area they have more surface energy. So more the energy with any particle more unstable it will be to share its energy. The size of nanoparticles ranges from 1-100nm. Nanoparticles surface is coated with PEG (Polyethylene Glycol), it provides the function of adhesion, stabilization etc. Nanoparticles are surface charged (positive as well as negative) and they have also functional groups on surface such as $-\text{SH}$, $-\text{COOH}$, $-\text{NH}_2$. Their surface is decorated with lipids and polymer chains of different molecular weights to provide them stealth properties and conjugated with different targeting ligands such as aptamers, peptides, antibodies to enhance the specific recognition of target cells. Nanoparticles can be of different shapes like rod shaped, cuboidal, spherical, star shaped etc” (Somasundaran et al., 2010).

Nanotechnology inAgriculture

“Nanotechnology has also major role in agriculture as is the backbone of most of the developing countries in which a major part of their income comes from agriculture sector and more than half of the population depends on it for their livelihood. The current global population is nearly 6 billion with 50% living in Asia. A large proportion of those livings in developing countries face daily food shortages as a result of environmental impacts or political instability, while in the developed world there is a food surplus. For developing countries, the drive is to develop drought and pest resistant crops which also maximize yield. In developed countries, the food industry is driven by consumer demand which is currently for fresher and healthier foodstuffs” (Anonymous, 2009). “Nanotechnology helps agricultural sciences and reduce environmental pollution by production of pesticides and chemical fertilizers by using the nano particles and nanocapsules with the ability to control or delayed delivery, absorption and more effective and environmentally friendly and production of nano-crystals to increase the efficiency of pesticides for application of pesticides with lower dose. Two principal factors cause the properties of nanomaterials to differ significantly from other materials increased relative surface area and quantum effects. Morphology-aspect ratio/size, hydrophobicity, solubility-release of toxic species, surface area/roughness, Surface species contaminations/adsorption, during synthesis/history, Reactive Oxygen Species (ROS) O₂ / H₂O, capacity to produce ROS, structure/composition, competitive binding sites with receptor and dispersion/ aggregation are the important properties of nanoparticles” (Somasundaran *et al.*, 2010)

Nanotechnology in control plant diseases

“About 20–40% of crops are lost due to plant pests and pathogens each year worldwide” (Flood, 2010). “In modern farming practices, pest management relies heavily on the application of pesticides, such as insecticides, fungicides, and herbicides. The development of cost-efficient, high-performing pesticides that are less harmful to the environment is crucial. The new concepts such as nanotechnology can offer advantages to pesticides, like reducing toxicity, improving the shelf-life, and increasing the solubility of poorly water-soluble pesticides, all of which could have positive environmental impacts. The significance of agricultural nanotechnology, mainly for controlling diseases and safety has been reported elsewhere. Nano-based conventional herbicides and pesticides assist in the slow and continued supply of nutrients and agricultural chemicals in a controlled amount to the plants” (Duhan *et al.*, 2017). “Nanoparticles may have also a key role in the control of insect pests and host pathogens. Type of polysaccharides such as chitosan, alginates, starch, and polyesters have been considered for the synthesis of nano-insecticide. In general, the use of nanoparticles to protect plants can occur via two different mechanisms: (a) nanoparticles themselves providing crop protection, or (b) nanoparticles as carriers for existing pesticides and can be applied by spray” (Worrall *et al.*, 2018). “However, the suse of nanomaterials in plant protection and production of food is under-explored” (Prasad *et al.*, 2017).

Nanotechnology to improve quality of soil and fertilizer distribution

“Nanotechnology for the management of crops is used as an essential technology for enhancing crop productivity. Nanomaterials and nanostructures, such as carbon nanotubes, nanofibers, and quantum dots are now exploited in agriculture research as biosensors for evaluating the quality of soil and fertilizer distribution. The purpose of nanoparticles is to minimize the spread of chemicals amount, reduce the nutrient loss during fertilization, and increase the quality and yield with proper

nutrient” (Sangeetha *et al.*, 2021). “The development and use of vermiculite, nanoclay, and zeolite could improve fertilizer efficacy and crop production for ecological agriculture in coarse-textured. Amending sandy loam soils with inorganic amendments reduce $\text{NH}_4\text{-N}$ passage and increasing the yield of N fertilizer in ecological agriculture systems” (Mazloomi and Jalali, 2019). “Nanoclay is systematized into a number of modules such as montmorillonite, bentonite, kaolinite, hectorite, and halloysite on the basis of chemical composition and nanoparticle morphology. Most of the productivity of agricultural practices is heavily dependent on fertilizer use. Studies show that crop production is linearly determined by exhaustive application of fertilizers to increase soil fertility” (Rehmanullah *et al.*, 2020). “The use of nano fertilizer is crucial to enhance crop production. Nano fertilizer is a material with nanometer-size which improves the delivery to plants and managed the slow release of nutrients into the soil gradually in a highly controlled way, hence stopping eutrophication and contamination of water” (Davari *et al.*, 2017). “Nanotechnology makes the exploitation of nanostructured or nanomaterials for fertilizer transport or limited release routes to construct smooth fertilizer as new opportunities to modify nutrient usage efficacy and reduce charges for environmental safety” (Hai *et al.*, 2011). “Nano-fertilizer could improve nutrient efficiency through encapsulation within nanoparticles which is conducted by three methods. (a) Nutrient encapsulation within nanoporous structures, (b) Coating of thin polymeric film, or (c) Delivery in the form of particle or suspensions with nanoscale sizes” (Davari *et al.*, 2017). “Nanoscale fertilizers could lead to the more effective delivery of nutrients as their small size may allow them access to plant surfaces and transport channels” (Mastronardi *et al.*, 2015). “Nano-fertilizer extracted and prepared from banana peels were used in the growth of tomatoes, peppers, or flowers. Nano fertilizers were used for the growth and improvement of different crops, for instance, nanoparticles of ZnO for chickpea, silicon dioxide and iron slag powder for maize, colloidal silica and NPK for tomato, TiO_2 for spinach, gold and sulfur fertilizers were used for the growth of grapes. Fertilizer usage with nanoscale transporters may be subjected in a way so that they anchor the roots of the plant with the surrounding soil contents and organic material hence decreasing chemical loss and lessening environmental issues” (Dasgupta *et al.*, 2015). “Nanoscale fertilizers can decrease the toxicity of soil and hence the potential undesirable impacts accompanied by high dosage are reduced” (Davari *et al.*, 2017). “Such nano fertilizers slow down the nutrients release and extend the duration of fertilizer impact. TiO_2 nanoparticles have shown a major effect on the growth of maize crop; moreover, SiO_2 plus TiO_2 nanoparticles elevated the action of nitrate and increased plant absorption potential, by controlled use of water and fertilizer with the efficient outcome” (Dasgupta *et al.*, 2015).

Nano-sensors in food and agriculture

“The use of biosensors combined with improved technologies in the field of molecular biology, nanomaterials, and microfluidics have enormous applications for the productivity of crops. These are also applied to monitor the activity of microorganisms in the soil and able to predict the likely incidence of soil infections. The basic principle related to soil examination with the biosensor is to find out the comparative action of positive and negative microorganisms in soil depends upon variation on oxygen usage during their breathing. They also offer many opportunities in sensing contaminants and their hindrance, via using new properties related to nanomaterials” (Baruah and Dutta, 2009). “Biosensors for detection of the nitrate concentration in plants as well as detection for

markers to identify infected plants are reported for methyl salicylate and azelaic acid” (Griesche and Baeumner, 2020). “The use of biosensors for monitoring of *Penicillium digitatum* infection in citrus fruit was reported” (Chalupowicz *et al.*, 2020). “Smart delivery systems and nanosensors are applied to help in efficient natural agricultural means like nutrients, water, and chemicals using precision farming for example satellite monitoring, geographic systems, and distant detecting tools that remotely can detect pests on crops or indication of strain like drought” (Sekhon, 2014). “The application of independent sensors connected to GPS monitoring in real-time is thought to play a key role in nanotechnology-assisted tools” (Davari *et al.*, 2017). “Arrangement of nanosensors can be carried out throughout the field for monitoring crop growth and soil parameters. Nanosensors are getting attention and playing a big role in the development of the food industry for its efficient responding tools to detect gases, microbes, or toxic substances in packed foods. Nanobiosensors have been testified for detecting pathogens in processing plants or alerting clients, protocols, and providers on the safety position of food” (Cheng *et al.*, 2006). “It has been also employed for the existence of impurities, mycotoxins, and microbes in food” (Bratovcic *et al.*, 2015). “Allergens have also been detected via biosensors tools with the assistance of nanoparticles and the report is about to commercialize” (Warriner *et al.*, 2014). These tools can also detect the history of time, expiration date, and temperature.



Purpose of nanotechnology in Phyto-nutraceuticals

“The term nutraceutical refers to a food substance, or a part of it, which possesses health benefits in terms of therapeutic or preventive effects”(Khorasani *et al.*, 2018). “Nutraceuticals include antioxidants, prebiotics, probiotics, herbal products, spices, polyunsaturated fatty acids, and many other compounds of natural and represent natural ways to achieve therapeutic goals”(Shinde *et al.*, 2014). “In the last few years, there was an increase in nutraceuticals consumption among the consumers due to an increase in natural-derived compounds interest; to date, an increasing interest in the development of novel functional food is pushing towards the incorporation of nutraceuticals within food products. The changes in lifestyle that have occurred in recent years have led to an increase in some diseases, such as type-2 diabetes and cardiovascular diseases”(Gruss *et al.*, 2019). “At the same time, the awareness among consumers of the close correlation between these dangerous diseases and eating habits has increased. For this reason, consumers are increasingly attentive to the quality of the consumed food and increasingly interested in food that can also have beneficial effects on their health and prevent the onset of dangerous diseases”(Goneares *et al.*,

2018). “All these aspects prompted researchers to study the potential beneficial effect on human health of nutraceuticals and their mechanism of action; at the same time, the industry was stimulated to develop innovative products capable of attracting consumers’ interest. Nanotechnology, based on structures with sizes in the order of nanometers is a revolutionary technology that has allowed researchers to overcome numerous limitations related to the use of nutraceuticals following their encapsulation into these structures, such as their stability, low solubility, and poor bioavailability”(Ting *et al.*, 2013). “It is noteworthy that the bioavailability, that is the fraction of a taken compound which is absorbed and available for physiological functions, is a key aspect for nutraceutical compounds, because their effectiveness is strictly related to their bioavailability. Unfortunately, different endogenous and exogenous factors can compromise the bioavailability of nutraceuticals, such as the biochemical transformations they may undergo into the epithelial cells, their physicochemical features, the food storage, and so on”(Leonard *et al.*, 2000). “For this reason, many innovative strategies have been thought to exert their beneficial effects when introduced into the organism. Among them, different nano-formulations have been designed to enhance the beneficial effects of nutraceuticals. Food nanotechnology could revolutionize the scenario of the food industry and agriculture by offering numerous advantages not only in increased bioavailability, but also promoting a controlled-release and targeted delivery of encapsulated bioactive natural compounds, which led to an increase in their biological efficacy representing exciting opportunities for the nutritional supplement industries. In particular, in the design of nutraceutical delivery systems, it must be taken into account that the formulation must have adequate chemical-physical properties, sustainable production costs, and food-grade materials must be used”(Huang *et al.*, 2010). Nutritional therapy and phyto-therapy have emerged as new concepts of health aid in recent years. Strong recommendations for consumption of nutraceuticals from plant origin have become progressively popular to improve health, and to prevent and treat diseases. Nutraceuticals are "naturally derived bioactive compounds that are found in foods, dietary supplements and herbal products, and have health promoting, disease preventing and medicinal properties." “Plant derived Nutraceuticals/functional foods have received considerable attention because of their presumed safety and potential nutritional and therapeutic effects. Some popular phyto-nutraceuticals include glucosamine from ginseng, Omega-3 fatty acids from linseed, Epigallocatechin gallate from green tea, lycopene from tomato etc. Some important phytonutraceuticals with their properties are Allicin from *Allium sativum*: It is a powerful antifungal antibacterial. It has been shown to be an antioxidant and has been used to treat arteriosclerosis and serum cholesterol. Betaine (Trimethyl Glycine) from green leafy vegetables and germinated grain it reduces toxic buildup of homocysteine” (Pulliainen *et al.*, 2010.) “Bromelain from *Ananas sp.* it is pineapple protease enzyme used to prevent heart disease, reduce the effects of aging, improve the immune system, and to reduce arthritis and inflammation. Camphor from *Cinnamomum camphora* used as an inhalant to treat cold and flu” (Chen *et al.*, 2002). Capsaicin or trans-8-methylN-vanillyl-5 nonenamide from *Capsicum annum* used for pain relief topically and as a digestive aid when taken internally. It is also seen as a possible antioxidant for the body. It can pose a risk of allergic reactions and the severe damage to the eyes or skin if used in higher doses. Ellagic Acid from strawberries and raspberries this phytochemical fights cancer in humans Vatter and Shetty, 2005. Ricinoleic acid from Castor oil or *Ricinus communis*. contains ricinoleic acid the active ingredient. Castor oil is used both externally (multiple skin problems) and internally for constipation, upper respiratory problems, and liver Okui, 1963. paday Curcumin from *Curcuma*

longa the colorant in turmeric a fraction of which has been shown by studies done at the University of California in Los Angeles to clear brain plaque caused by Alzheimer's disease mishra Ram *et al.*, 2000. "Isoquercitin from mangoes and from *Rheum nobile* (Enzyme Modified) increases blood flow for varicose veins, and possible use for arterial flow as well. Recent studies have shown possibilities in increased brain functions and it might be useful in the treatment of progressive Alzheimer's disease. Lutein and Lutein Esters from marigold extracted from marigold seeds, and also found in spinach, rosemary and kale, it is a carotenoid which shows healthful eye benefits. Resveratrol especially high in grape skin anti-inflammatory, inhibits COX-1 enzyme, blocks adhesion of blood cells to vessel walls shown to reduce skin and breast cancer"(Gehm *et al.*, 1997). "Zeaxanthin a carotenoid used as an antioxidant. It is the coloring agent in marigolds and is extracted from them. It is used for eye health and some claim will retard the effect of 'aging eyesight' or Age-Related Macular Degeneration (AMD). But, the main challenge still in fully realizing the potential of nutra-ceuticals has been their poor bioavailability after consumption. These substances are generally having low solubility, leading to their poor absorption and bioavailability in the human body. In this regard, one of the most important applications of nanotechnology in food sector has been the formulation of novel nutraceutical compounds with improved properties viz. enhanced solubility, stability, bioavailability and efficacy. This is achieved by encapsulation of nutraceuticals by nanoparticles, which modifies their pharmacokinetics (PK) and biodistribution (BD). The choice of the correct vehicle of delivery for the active nutraceutical or pharmaceutical ingredients into the human body is very crucial for facilitating the direct contact of the ingredients with the target site of action in the body. As most of these are either poorly soluble or lipophilic compounds their delivery is significantly enhanced by altering the physicochemical properties like water solubility, partition coefficient, lipophilicity, crystallinity, etc. The poor solubility of the active ingredients poses multiple challenges to their full utilization: route of administration, transport in the physiological system and reaching the site of action. All these in turn ultimately lead to the poor bioavailability of the same in the organisms. Bioavailability is one of the most crucial and indispensable property of any dietary or pharmaceutical ingredient, which decides what proportion of the therapeutically active component reaches the systemic circulation in the host and is available at the target site for action. Many types of nutraceuticals are available today in the market with various proclaimed health benefits. One such prominent example among nutraceuticals are the phytochemicals including plant polyphenols (curcumin, resveratrol) carotenoids, (lycopene, β carotene, lutein) etc. These are widely favoured by the researchers, food manufacturers and consumers alike, because of their multiple health benefits viz. blood pressure regulation, reducing the probability of having malignant diseases like cancer, promoting digestion, immunity and growth, regulating glucose and cholesterol levels and also reducing stress by acting as antioxidants. Nanotechnology is a latest and advanced field, wherein the unique physicochemical characteristics of nanoparticles are used to significantly alter the structure, texture and quality of phytoutaceuticals. Applications of nanotechnology in this sector has been the formulation of novel phyto nutraceutical compounds with improved properties viz. enhanced solubility, stability, bioavailability and efficacy. This is achieved by encapsulation of nutraceuticals by nanoparticles, which modifies their pharmacokinetics (PK) and bio-distribution (BD)" (Sadhana Raut., et al2022).

Characterization of nano-materials used for phyto-nutraceutical purposes

“Data about certain other physical parameters are needed for that e.g., the physical form, morphology, particle size distribution etc, which can be obtained through imaging techniques like scanning and transmission electron microscopy (SEM and TEM). Light scattering techniques e.g., Dynamic light scattering (DLS) can provide additional information about particle size distribution of NMs in liquid samples, like average size and even size distribution of NMs, both during the synthesis and in physiological processes like digestion. Stability, which is another very basic and important property of any of the active ingredients inside the body, like the various nutraceutical components, can be measured by laser Doppler micro-electrophoresis method over a period of time, as zeta potential” (Ting et al., 2013).

Importance of nanoparticles in Phytonutraceuticals

“There are mainly four types of nanoparticles in nutraceuticals viz; nanoliposomes, nanoemulsions, nanoparticles. The nano-emulsion formulations of active ingredients can be used for developing biodegradable coating and packaging films to enhance the quality, functional properties. Nanoencapsulation technique provides the possibility to protect the chemical structure of phyto nutraceuticals from environmental agents such as pH, light, temperature, radicals, or oxygen increases their bioavailability; allows specific delivery to target sites and allows a controlled release of the encapsulated compound”(Cristianno et al., 2021 ; Assadpur *et al.*, 2019). “Concerning the ability of nanosystems to control the release of the delivered active compounds, it consists of a specific concentration/time release profile at the desirable site of action and it is the main challenge for nutraceuticals encapsulation. Therefore, an ideal delivery system should be able to release its content following specific stimuli such as pH, moisture, enzymes, and temperature, and, at the same time, to protect the nutraceutical from the same stimuli”(Mclementis *et al.*, 2010). “Moreover, the encapsulation of nutraceutical compounds leads to an enhancement in their solubility, as, once the nutraceutical is loaded into the carrier, features are dependent on the physico-chemical characteristics of the vesicle rather than to the entrapped compound. Nanosystems also provide the possibility to co-deliver water- and lipid-soluble molecules, thus supporting their synergistic effect. They are also able to guarantee the physico-chemical stability and avoid undesirable changes in smell and taste that might result from the addition of nutraceuticals to food products. The materials used for the realization of the drug delivery system can be of various nature (lipid, polymeric, protein) as long as it has the Generally Recognized as Safe (GRAS) status”(Fox *et al.*, 2014).

Nanoparticles: Nanoparticles (NPs; 1–100 nm in size) have a special place in nanoscience and nanotechnology, not only because of their particular properties resulting from their reduced dimensions, but also because they are promising building blocks for more complex nano structures tritional value, and shelf life of foods. Nanoparticles are widely used drug delivery systems and can be made of different material, for example, polymers (poly-D,L-lactide-co-glycolide, polylactic acid, poly- ϵ -caprolactone), proteins, and lipids Desfrancois *et al.*, 2018. In particular, in order to be applied in food and nutraceutical fields, food-grade material for the fabrication of nanoparticles must be used. Among the food grade material zein, a maize protein, chitosan, and gelatin are widely used Feng *et al.*, 2020. Due to their biodegradability, bioavailability, and the possibility of encapsulating hydrophobic compounds, soy proteins have attracted the researchers' attention to be used in the design of nanocarriers for the delivery of bioactives, nutraceuticals included Verma *et al.*, 2018. Soy β -conglycinin (a storage globulin) was employed for the development of

nanoparticles for the encapsulation of hydrophobic curcumin, a polyphenol with anti-oxidant and anti-inflammatory activities. A new method, based on disassembly and reassembly of β -conglycinin, which is the vicilin storage protein of soybeans, was performed using urea and without adding any organic solvent; the obtained nanoparticles, produced with this new technology, turn out to be more natural and are characterized by a good solubility and encapsulation efficiency (around 80%) greater than that obtained in previous works. The bioaccessibility of curcumin was found to be around 40% (while that of free curcumin was found to be around 20%). β -conglycinin nanostructures represent promising biocompatible delivery systems for hydrophobic compounds Liu *et al.*, 2019. Another field of application of nanoparticles for the delivery of nutraceuticals is the field of cancer, to improve the activity of drug therapies and to decrease their side effects. Recently, Cosco and collaborators proposed hyaluronan-coated PLGA (Poly Lactic-co-Glycolic Acid) nanoparticles in which sclareol, a diterpene obtained from Clary sage (*Salvia sclarea* Linn.) Dimas *et al.*, 1999, was encapsulated to favor its administration in physiological media, thus improving its anticancer efficacy. Characterization studies showed that the realized nanoparticles had mean sizes of 100–150 nm showing a reduction in their diameter due to the addition of sclareol. The coating of nanosystems was performed to improve the anticancer efficacy of the delivered phytochemical, due to the interaction and internalization of the realized structures with HA+ cancer cells. In fact, it was concluded that the anticancer efficacy was properly related to the coating of nanoparticles, using hyaluronic acid (1.5 MDa), which promoted the interaction with the hyaluronan receptors expressed on breast cancer cell lines, MCF-7 and MDA-MB468. The amount of hyaluronic acid adsorbed on the surface of the nanosystems was detected through the carbazole assay showing that the coating efficiency did not increase over 1 mg of hyaluronic acid added to the formulation. Another nanosystem, comprising beta carotene-loaded zein nanoparticles, was developed by Jain and collaborators to study its potential use in breast cancer. The obtained system showed a greater anticancer activity, with respect to free beta-carotene both in vitro (MCF-7 cells) and in vivo (induced breast cancer in rats): this is probably due to the increased cellular intake of zein nanoparticles. It also is worth mentioning that the association between beta-carotene-loaded nanoparticles with free methotrexate, the most widely used anticancer drug, showed a double positive effect: a synergistic effect, obtaining a strong in vitro anticancer activity on MCF-7 (breast cancer cell line) cells and a reduction in methotrexate side effects on the liver and kidneys [59]. Resveratrol is another nutraceutical, which showed greater anticancer efficacy against MCF-7 when encapsulated into nanoparticles. This polyphenolic compound shows different anti-oxidant, anti-inflammatory, and anticancer activities. Unfortunately, it is practically insoluble in water (~ 0.03 mg/mL at 25 °C), and to overcome this difficulty, the encapsulation into cyclodextrins was recently proposed in particular, it was evidenced that complexation produced a consistent improvement in the solubility of resveratrol in water and consequently a significant improvement in the anticancer activity of resveratrol on several cell lines Chen *et al.*, 2019.

Liposomes: “Nanoliposome technology presents exciting opportunities for food technologists in areas such as encapsulation and controlled release of food materials, as well as the enhanced bioavailability, stability, and shelf-life of sensitive ingredients. Liposomes are lipid-based vesicles and represent a versatile and biocompatible drug delivery system used for the encapsulation of both hydrophilic and hydrophobic drugs. Liposomes are employed for the delivery of actives with different pharmaceutical activities nutraceuticals included”(Subramani *et al.*, 2020).The all trans-

retinoic acid (ATRA) is a metabolite of vitamin A. It is a nutraceutical compound widely studied for its anticancer property. It was encapsulated in liposomes, obtaining an entrapment efficiency of around 82% in order to protect it from degradation. In particular, Cristiano et al. demonstrated that the encapsulation of ATRA within liposomes allows it to protect the drug from photo-degradation phenomena that would compromise its pharmacological activity.

Nanoemulsions: Nanoemulsions have small droplet size and are kinetically stable colloidal systems. They have enhanced functional properties in comparison to conventional emulsions. The composition and structure of the nanoemulsions can be controlled for the encapsulation and effective delivery of bioactive lipophilic compounds. Nanoemulsions have potential application in the food industry for the delivery of phytonutraceuticals, coloring and flavoring agents, and antimicrobials. Nanoemulsions are formulations made of a water phase, an oily phase, and an emulsifier, and are characterized by a droplet size of around 100 nm. Gupta *et al.*, 2016. Oil-in-water and water-in-oil nanoemulsions are used for the encapsulation and protection of active ingredients and represent a suitable delivery system for the encapsulation of nutraceuticals, improving the efficacy of hydrophobic and hydrophilic active molecules and food components. Zhang *et al.*, 2020. Nanoemulsions were shown to be promising drug delivery systems for nutraceuticals, tomato extract rich in lycopene and curcumin, two antioxidant agents, was encapsulated into this system. Chang and collaborators nanoemulsion formulation was prepared by using the ultra-high-pressure homogenization method for the encapsulation of the oil extracted from the pulp of sea buck-thorn. Liu and collaborators were able to improve the oral bioavailability of the nutraceutical astaxanthin, a carotenoid which possesses numerous health benefits. Ambati *et al.*, 2014. Three types of long chain triglycerides, varying according to the fatty acid composition, were used for the nanoemulsions preparation: corn oil, olive oil, and flaxseed oil.

Development of phytonutraceuticals:

“During last few years, there was an increase in nutraceuticals consumption among the consumers due to an increase in natural-derived compounds interest; to date, an increasing interest in the development of novel functional food is pushing towards the incorporation of nutraceuticals within food products. Unfortunately, different endogenous and exogenous factors can compromise the bioavailability of nutraceuticals, such as the biochemical transformations they may undergo into the epithelial cells, their physicochemical features, the food storage, and so on. For this reason, many innovative strategies have been thought to exert their beneficial effects when introduced into the organism” (Punia *et al.*, 2019)..

Different nanoformulations designed to enhance the beneficial effects of nutraceuticals are:

Nanocurcumin: “Turmeric or Curcuma is a natural creation, whose therapeutic properties have been widely studied and an extensive variety of therapeutic effects on several diseases. Nano curcumin particles were prepared by Vibra-Cell Ultrasonic Liquid Processors device. The particles are characterized by scanning probe microscope (SPM) and Ultraviolet-visible spectroscopy (UV-Vis spectrophotometer). The results confirm that the prepared Nano curcumin has mean diameter 82 nm. The prepared Nano curcumin is exposed to plasma to enhance the properties of their Nano particle the result is improve the enhanced characterization” (Sabah *et al.*, 2020)

Allicin nanocapsules:“They were prepared via ionotropic pre-gelation. The wall materials were alginate-chitosan biopolymers. Nanocapsules were characterized using Fourier transform infrared spectroscopy (FT-IR) and field emission scanning electron microscopy (FESEM)”(Ghadir *et al.*, 2016)

Nano-Resveratrol:Resveratrol is another nutraceutical, which showed greater anticancer efficacy against MCF-7 when encapsulated into nanoparticles. This polyphenolic compound shows different anti-oxidant, anti-inflammatory, and anticancer activities. Unfortunately, it is practically insoluble in water (~0.03 mg/mL at 25 °C), and to overcome this difficulty, the encapsulation into cyclodextrins was recently proposed in particular, it was evidenced that complexation produced a consistent improvement in the solubility of resveratrol in water and consequently a significant improvement in the anticancer activity of resveratrol on several cell lines.

Nano- green tea extract: the wet milling technique is useful for making green tea nanoparticles for industrial scale.

Nano-Retinol:Retinol has been widely used as an anti-wrinkle active ingredient in cosmetic fields. Retinol-loaded lipid nano-carriers prepared via the vacuum emulsification method helps to increase the stability of retinol vulnerable to air and optimized encapsulation conditions and to increase the penetration efficiency into skin.

Nano-quercetin:Quercetin is a plant flavonol from the flavonoid group of polyphenols. It is found in many fruits, vegetables, leaves, seeds, and grains; capers, red onions, and kale are common foods containing appreciable amounts of it.

Conclusion

Nano-formulations widely used for the purpose are nanoliposomes, nanoemulsions, nanoparticles, nanofibres. The particles are characterized by scanning probe microscope (SPM), Ultraviolet-visible spectroscopy (UV-Vis spectrophotometer, SEM, TEM, Dynamic light scattering (DLS).

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that no generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

References

- Alorabi, A.Q., Hassan, M.S., Algethami, J.S. and Baghdadi, N.E., 2021. Synthesis and characterization of Ag-AgVO₃/Cu₂ heterostructure with improved visible-light photocatalytic performance. *Science Progress*, 104(4), p.00368504211050300.
- Anonymous.2009. Nanotechnology and Nanoscience applications: Revolution in India and beyond. *Strategic Applications Integrating Nano Science*.
- Baruah, S. and Dutta, J., 2009. Nanotechnology applications in pollution sensing and degradation in agriculture: a review. *Environmental Chemistry Letters*, 7, pp.191-204.
- Boddolla, S. and Thodeti, S. 2018. A review on characterization techniques of nanomaterials. *International Journal of Engineering, Science and Mathematics*,;7(1): 169-175.
- Bratovic, A., Odobasic, A., Ćatić, S. and Šestan, I., 2015. Application of polymer nanocomposite materials in food packaging. *Croatian journal of food science and technology*, 7(2), pp.86-94.
- Bruggemann, D., Wolfrum, B., Maybeck, V., Mourzina, Y., Jansen, M. and Offenhäusser, A., 2011. Nanostructured gold microelectrodes for extracellular recording from electrogenic cells. *Nanotechnology*, 22(26), p.265104.
- Cardoza, C., Nagtode, V., Pratap, A. and Mali, S.N., 2022. Emerging applications of nanotechnology in cosmeceutical health science: Latest updates. *Health Sciences Review*, p.100051.
- Chen, C.W., Lee, S.T., Wu, W.T., Fu, W.M., Ho, F.M. and Lin, W.W., 2003. Signal transduction for inhibition of inducible nitric oxide synthase and cyclooxygenase-2 induction by capsaicin and related analogs in macrophages. *British journal of pharmacology*, 140(6), pp.1077-1087.
- Chinnamuthu, C.R. and Boopathi, P.M., 2009. Nanotechnology and agroecosystem. *Madras Agricultural Journal*, 96(jan-jun), p.1.
- Das L., Bhaumik E. Raychaudhuri U. Chakraborty R. 2012. Role of nutraceuticals in human health.

- Dasgupta, N., Ranjan, S., Mundekkad, D., Ramalingam, C., Shanker, R. and Kumar, A., 2015. Nanotechnology in agro-food: from field to plate. *Food Research International*, **69**, pp.381-400.
- Davari, M.R., Bayat Kazazi, S. and Akbarzadeh Pivezhzani, O., 2017. Nanomaterials: implications on agroecosystem. *Nanotechnology: An Agricultural Paradigm*, pp.59-71.
- Duhan, J.S., Kumar, R., Kumar, N., Kaur, P., Nehra, K. and Duhan, S., 2017. Nanotechnology: The new perspective in precision agriculture. *Biotechnology Reports*, **15** :11-23.
- Fakoor Yazdan Abad, M., Rajabzadeh, G., Taghvaei Ganjali, S. and Tavakoli, R., 2016. Preparing allicin nanocapsules and determining the factors controlling their particle size through artificial intelligence. *International Journal of Food Engineering*, 12(3) :257-264.
- Flood, J., 2010. The importance of plant health to food security. *International Journal of Food Microbiology* **124**(2): 142-146.
- Gehm, B.D. McAndrews, J.M. Chien, P.Y. and Jameson, J.L., 1997. Resveratrol, a polyphenolic compound found in grapes and wine, is an agonist for the estrogen receptor. *Proceedings of the National Academy of Sciences*, **94**(25): 14138-14143.
- Gruss S.M. Nhim K., Gregg E., Bell M. Luman E., Albright A.2019. Public health approaches to type 2 diabetes prevention: *The US National Diabetes Prevention Program and beyond*. **19**:78.
- Huang Q. Yu H. Ru Q.2017. Bioavailability and delivery of nutraceuticals using nanotechnology. *Journal of Food Science* **75**:50–57.
- Jagannathan R., Patel S.A., Ali M.K., Narayan K.M.V.2019. Global Updates on Cardiovascular Disease Mortality Trends and Attribution of Traditional Risk Factors. 19:44.
- Jeevanandam, J., Barhoum, A., Chan, Y.S., Dufresne, A. and Danquah, M.K., Review on nanoparticles and nanostructured materials: history, sources, toxicity and regulations. *Beilstein journal of nanotechnology*, 2018;9(1):1050-1074.
- Joseph T, Morrison M.2006. Nanotechnology in Agriculture and Food Institute of Nanotechnology. *A Nano forum report*.

- Joshi, H., Choudhary, P. and Mundra, S.L., 2019. Future prospects of nanotechnology in agriculture. *International Journal of Chemistry Studies* **7**(2) : 957-963.
- Khorasani S. Danaei M. Mozafari M.R. Nanoliposome technology for the food and nutraceutical industries. *Trends Food Sci. Technology* 2018;**79**:106–115.
- Leonard N.B.2000. Handbook of Nutraceuticals and Functional Foods. CRC Press; Boca Raton, FL, USA: *Stability testing of nutraceuticals and functional foods*.
- Mishra, S. and Palanivelu, K. 2008. The effect of curcumin (turmeric) on Alzheimer's disease: An overview. *Annals of Indian Academy of Neurology*, **11**(1):13.
- Moulick, R.G., Das, S., Debnath, N. and Bandyopadhyay, K., 2020. Potential use of nanotechnology in sustainable and 'smart' agriculture: advancements made in the last decade. *Plant Biotechnology Reports*, **14** :505-513.
- Mourdikoudis, S., Pallares, R.M. and Thanh, N.T. 2018. Characterization techniques for nanoparticles: comparison and complementarity upon studying nanoparticle properties. *Nanoscale*, **10**(27) :12871-12934.
- Neme, K., Nafady, A. Uddin, S. and Tola, Y.B. 2021. Application of nanotechnology in agriculture, postharvest loss reduction and food processing: *Food security implication and challenges Heliyon*, **7**(12).
- Panyam, J. and Labhasetwar, V. 2003. Biodegradable nanoparticles for drug and gene delivery to cells and tissue. *Advanced Drug Delivery Review* **55**(3): 329-347.
- Pourpirali, R., Mahmoudnezhad, A., Oroojalian, F., Zarghami, N. and Pilehvar, Y. 2021. Prolonged proliferation and delayed senescence of the adipose-derived stem cells grown on the electrospun composite nanofiber co-encapsulated with TiO₂ nanoparticles and metformin-loaded mesoporous silica nanoparticles. *International journal of pharmaceutics*, **604**:120733.
- Prasad, R. Bhattacharyya, A. and Nguyen, Q.D., 2017. Nanotechnology in sustainable agriculture: recent developments, challenges, and perspectives. *Frontiers in microbiology*, **8**: 1014.
- Punia S., Sandhu K.S., Kaur M., Siroha A.K. Nanobiotechnology Bioformulation. Springer; Cham, Switzerland: 2019. Nanotechnology: A Successful Approach to Improve Nutraceutical Bioavailability; pp. 119–133.

- Rehmanullah, Muhammad, Z., Inayat, N. and Majeed, A., 2020. Application of nanoparticles in agriculture as fertilizers and pesticides: challenges and opportunities. *New frontiers in stress management for durable agriculture*, pp.281-293.
- Sangeetha, J. Hospet, R. Thangadurai, D., Adetunji, C.O. Islam, S. Pujari, N. and Al-Tawaha, A.R.M.S. 2021. Nanopesticides, nanoherbicides, and nanofertilizers: the greener aspects of agrochemical synthesis using nanotools and nanoprocesses toward sustainable agriculture. *Handbook of Nanomaterials and Nanocomposites for Energy and Environmental Applications*, pp.1663-1677.
- Sastry, R.K. Rashmi, H.B. and Rao, N.H. 2011. Nanotechnology for enhancing food security in India Food Policy, **36**(3) :391-400.
- Shinde N. Bangar B., Deshmukh S. Kumbhar P.2014. Nutraceuticals: A review on current status. *RJPT*. **7**:110–113.
- Shinde N., Bangar B., Deshmukh S., Kumbhar P. Nutraceuticals: A review on current status. *RJPT*. 2014;**7**:110–113.
- Singh, A., Sekhon, A.S. Unger, P. Babb, M., Yang, Y. and Michael, M. 2021. Impact of gas micro-nano-bubbles on the efficacy of commonly used antimicrobials in the food industry. *Journal of Applied Microbiology*, **130**(4) :1092-1105.
- Singh, A.R., Desu, P.K., Nakkala, R.K., Kondi, V., Devi, S., Alam, M.S., Hamid, H., Athawale, R.B. and Kesharwani, P., 2021. Nanotechnology-based approaches applied to nutraceuticals. *Drug Delivery and Translational Research*, pp.1-15.
- Somasundaran P, Fang X, Ponnurangam S, Li B.2010. Nanoparticles: Characteristics, mechanisms and modulation of biotoxicity. *Kona Powder and Particle Journal*. **28**:38-49.
- Spira, M.E. and Hai, A., 2020. Multi-electrode array technologies for neuroscience and cardiology. *Nano-Enabled Medical Applications*, pp.567-602.
- Thodeti, S., Reddy, R.M. and Kumar, J.S., Synthesis and characterization of pure and indium doped SnO₂ nanoparticles by sol-gel methods. *Int. J. Sci. Eng. Res*, 2016;**7**:310-317.
- Ting Y. Jiang Y., Ho C.-T. Huang Q.2014. Common delivery systems for enhancing in vivo bioavailability and biological efficacy of nutraceuticals. *Journal of Functional Foods*. **7**:112–128.

- Vattem, D.A. and Shetty, K., 2005. Biological functionality of ellagic acid: a review. *Journal of food biochemistry*, **29**(3) :234-266.
- Wani, K.A. and Kothari, R., 2018. Agricultural nanotechnology: applications and challenges. *Ann Plant Science*, **7**: 2146-2148.
- Warriner, K. Reddy, S.M. Namvar, A. and Neethirajan, S. 2014. Developments in nanoparticles for use in biosensors to assess food safety and quality. *Trends in food science & technology*, **40**(2):183-199.
- Worrall, E.A., Hamid, A., Mody, K.T., Mitter, N. and Pappu, H.R., 2018. Nanotechnology for plant disease management. *Agronomy*, **8**(12): 285.
- Wu, D. Wang, S. Yu, G. and Chen, X. 2021. Cell death mediated by the pyroptosis pathway with the aid of nanotechnology: prospects for cancer therapy. *Angewandte Chemie*, **133**(15): 8096-8112.
- Sadhana Raut., et al. "Pharmacognostic and Pharmacological Aspects on *Tabernaemontana divaricata* Plant". *Acta Scientific Pharmacology* 3.10 (2022): 22-34.