

VITAMIN D STATUS IN AFRICAN PATIENTS ON SUPPLEMENTED HEMODIALYSIS, FOLLOWED IN THE PUBLIC SECTOR IN CÔTE D'IVOIRE

Abstract

Introduction

Chronic kidney disease (CKD) causes decreased activity of the enzyme 1-alpha-hydroxylase and calcitriol deficiency with mineral and bone metabolism disorders or secondary hyperparathyroidism. Anticipation of the occurrence of these complications implies the systematic supplementation of all hemodialysis patients, with monitoring of biological parameters. It is in this context that this study was carried out, the general objective of which was to establish the profile of vitamin D concentration in hemodialysis and Ivorian public sector supplemented patients in Abidjan.

Material and methods

In a cross-sectional study conducted from September to December 2022, we included 89 patients with end-stage CKD treated with hemodialysis at the CNPTIR in Cocody. The collection of epidemiological data was done on the basis of a questionnaire on the one hand and the patients' medical records on the other. The vitamin D concentration was determined by the ELFA immunoassay method on mini Vidas (BIOMERIEUX).

Results

The mean age of the patients was 45 ± 13 and 65.17% of them were men. 55.06% of the patients had been treated with dialysis for at least 5 years. 48.31% of the patients had regular supplementation treatment. Of the two types of vitamin D supplementation, 94.38% of the study population was supplemented with cholecalciferol compared to 18% for alfacalcidol with 12% of patients included, using both types. From a biological point of view, it appears that the average concentration of vitamin D in the patients was 26.9 ng/mL, and only 16% have a concentration of less than 20 ng/mL, corresponding to a deficiency vitamin D status.

Conclusion

At the end of this cross-sectional study, it emerged that the average in adult hemodialysis subjects followed at the CNPTIR in Cocody had an insufficient vitamin D status. And 37% of the study population with normal vitamin D status. However, these results could be improved by better accessibility to the products, and regular compliance with supplementation.

Keywords: Vitamin D- African - hemodialysis - chronic kidney disease

1 – INTRODUCTION

In humans, cholecalciferol or vitamin D₃ (VitD₃) has a dual origin: an endogenous origin occurring in the epidermis and an exogenous origin (food and drug intake).

Its main biological role (the regulation of phosphocalcic metabolism) requires activation by two successive hydroxylations: the first, at the level of the liver and the second at the level of the kidney (by the enzyme 1-alpha-hydroxylase) to obtain calcitriol or active vitamin D3: 1,25(OH)₂D₃ (1, 17)

Also, chronic kidney disease (CKD), by causing the decrease or even suppression of the activity of the enzyme 1-alpha-hydroxylase, leads to calcitriol deficiency at the origin and to mineral and bone metabolism disorders such as osteoporosis and secondary hyperparathyroidism constituting a serious complication in dialysis patients.

Also, calcitriol deficiency leads to hypocalcemia due to a decrease in intestinal absorption and stimulates the synthesis of parathyroid hormone (PTH), which allows the hydroxylation of 25 hydroxy vitamin D (25(OH)D) to 1,25(OH)₂D₃, active. This parathyroid regulation allows the stability of 1,25(OH)₂D₃ levels at the expense of 25(OH)D, regardless of the amount of sunlight. This deficiency therefore induces secondary hyperparathyroidism (HPT) responsible for osteoporosis or osteomalacia in adults

A meta-analysis by Bischoff-Ferrari (18) shows a reduction in the risk of non-vertebral fracture in patients receiving vitamin D supplementation > 800 IU/day (effect more pronounced in patients deficient at the start of the study and those over 70 years of age). There is a lower response to anti-osteoporotic treatments when vitamin D stocks are insufficient, the necessary level being 30 to 40 ng/mL.

Thus, the measurement of vitamin D concentration in CKD patients can prevent or improve the management of complications related to vitamin D deficiency (18).

Thus, our study aims to establish the status of chronic kidney failure patients on dialysis in public centers in Abidjan (Côte d'Ivoire) receiving vitamin D supplementation

2- MATERIALS AND METHODS

2.1 - Study design and patients

This is a cross-sectional study, with a descriptive aim that took place over a period of 3 months. It was carried out at the biology laboratory of the SAMU (Emergency Medical Assistance Service) in Abidjan.

The study focused on a population of patients suffering from end-stage CKD, treated with hemodialysis and supplemented with vitamin D at the public hemodialysis center in Abidjan: CNPTIR (National Center for the Prevention and Treatment of Renal Failure).

Patient recruitment was done according to the inclusion criteria below:

Subjects of both sexes aged 18 years or older, with chronic kidney disease and vitamin D supplementation, who have given consent to participate in the study with an available medical record.

2.2 - Method

Sampling was carried out by successive recruitment of patients meeting the inclusion criteria over the study period. Questioning and consultation of patients' medical records made it possible to collect epidemiological-clinical data, in particular the characteristics of vitamin D supplementation treatment (prescribed specialty, dosage and dosage) and patient compliance.

The selected patients were subjected to a venous blood sample at the elbow crease on an empty stomach tube without anticoagulant (with separator gel). On the serum collected after centrifugation at 4000 rpm for 5 minutes, the vitamin D was measured.

The vitamin D assay was done by the ELFA (Enzyme Linked Fluorescent Assay) method on Mini VIDAS® (BIOMERIEUX).

The data was processed using the Office Excel 2016 software.
 The concentration of vitamin D was interpreted according to the following values:
 25(OH)D < 20 ng/ mL : Deficiency
 25(OH)D between 20 and 25 ng/mL: Impairment
 25(OH)D between 25 and 30 ng/mL: Insufficiency
 25(OH)D ≥ 30 ng/ mL : Normale

2.3 - Statistical analysis

Quantitative variables were described using the mean, standard deviation, extremes, median and interquartile range (25th percentile (P25)-75th percentile (P75)). Each of the qualitative variable modalities was described in terms of numbers and percentages.

3 - RESULTS

More than half of the study population is under the age of 49. The average age of patients is 45 years with a maximum of 80 years.

Male patients accounted for 65.17%, i.e. a sex ratio (M/F) of 1.87.

The majority of patients have been dialysis for more than 5 years. The average duration of hemodialysis treatment is 5.78 years with extremes of 1 to 20 years. This duration of treatment also corresponds to that of vitamin D supplementation (vit D).

Table I : Patient characteristics

Parameters	Values
Age (years)	
Average ± standard deviation	45 ± 7
Minimum - maximum	18 - 80
Gender : number (%)	
Men	58 (65,17%)
women	31 (34,83%)
Length of time on dialysis treatment	5,78 ± 1,5

Compared to the prescribed molecules, two types of supplementation are observed: on the one hand, supplementation with specialties containing the combination of calcium + cholecalciferol and on the other hand, supplementation with 1 alpha-hydroxycholecalciferol prescribed in combination with calcium only orally.

About half of the patients in our sample are unable to follow the supplementation treatment on a regular basis.

More than half of the patients have a vitamin D concentration (greater than or equal to 25 ng/mL). The average concentration of vitamin D in our series is 26.9 ± 8.8 ng/mL with a maximum of 52.4 ng/mL and a minimum of 8.1 ng/mL. Regarding supplementation treatment, 69.77% of patients who observed regular compliance had a vitamin D ≥ 25 concentration compared to 43.48% in non-compliant patients.

Table II : Distribution of the study population by vitamin D concentration

Plasma concentration of vitamin 25(OH)D (ng/mL)		Values
Mean \pm standard deviation		26,9 \pm 8,8
Médian		26,8
Minimum - maximum		8,1 - 52,4
Distribution Number (%)	< 20	14 (16%)
	[20 – 25[17 (19%)
	[25 – 30[25 (28%)
	≥ 30	33 (37%)
	total	89 (100%)

Table III: Distribution of the study population by vitamin D status and good compliance to supplementation

	Plasma concentration of vitamin 25(OH)D (ng/mL)				Total
	<20	[20-25[[25-30[≥ 30	
good compliance					
No	16	10	7	13	46
Yes	9	4	10	20	43
Total	25	14	17	33	89

DISCUSSION

The average age of our study population is 45 ± 7 years with extremes of 18 and 80 years. The distribution by age group showed that more than half of the study population is under 49 years of age; The relative youth of CKD patients highlighted in our study is also observed in other studies in Africa, particularly in Burkina Faso (2) with an average age of 45 years. On the other hand, in Caucasian populations with CKD, the average reported ages remain above 60 years (3,4). These observed differences are widely documented, involving in particular the youth of the general population in Africa as well as self-medication (5), the frequent and endemic nature of certain diseases such as schistosomiasis, HIV-AIDS (6).

The male predominance observed in our population has also been reported in numerous studies both in Africa and elsewhere in the world (7–9). Generally speaking, many authors agree that men are more affected by CKD than women due to the higher frequency of kidney disease in men (10), and the notion of smoking that could have a detrimental effect on the rate of progression of chronic kidney disease.

Regarding the length of time spent on hemodialysis treatment, the average of 6 years we found is dissimilar to those reported by El harraqui et al (11) in Morocco (7 years) and Dammack et al (12) in Tunisia (9 years). This duration can be superimposed on that of vitamin D supplementation.

The median concentration of vitamin D found in our study is 26.8 ng/mL. It is significantly lower than that reported by Cavalier et al. (13) in 2018 in CKD patients in the same city of Abidjan (40.2 ng/mL). Similarly, the proportions of patients with 25(OH)D levels greater than 30 ng/mL differ significantly (37% in our study vs. 80% in the Cavalier study). These differences can be explained by the platforms used for the determination of 25(OH)D in the 2 studies: VIDAS bioMérieux vs. Fujirebio Lumipulse. Indeed, several authors have noted the tendency of certain equipment (notably VIDAS) to under-recover 25(OH)D compared to other equipments such as Fujirebio Lumipulse (13,14).

Compared to other populations of chronic kidney disease, particularly in Morocco, our results remain relatively low, especially compared to those reported by Benabdellah et al (15) (mean vitamin D concentration equal to 34.8 ng/mL) without specifying the assay platform used.

Moreover, in the same city of Abidjan and for the same assay technique (Vidas bioMérieux), the proportion of 25(OH)D deficiency reported by Boyvin (16) in a healthy population is 33% compared to 16% in our dialysis population. This result suggests that systematic supplementation of these at-risk patients helps maintain an acceptable vitamin D stock. Of the different types of vitamin D supplementation, 94.38% of the study population was supplemented with cholecalciferol, 18% with alfacalcidol, and 12% used both types.

The low prescription rate of alfacalcidol could be explained by its high cost and low availability on the Ivorian market. Also, its prescription is generally used as a curative treatment after the onset of hyperparathyroidism or bone complications.

For cholecalciferol, the daily doses used varied from 400 IU to 6000 IU depending on the patient's blood calcium level and the pharmaceutical specialty prescribed.

Although relatively low, the proportion of dialysis patients with vitamin D deficiency could have been lower if compliance with supplementation was correct in all patients.

Indeed, the analysis of the average vitamin D concentration value as a function of the regularity of supplementation shows that 65% of deficient patients are non-compliant. This poor compliance is mainly linked to financial difficulties in honoring prescribed prescriptions, as vitamin D is not covered by the Ivorian state subsidy or reimbursed by insurance companies.

CONCLUSION

At the end of our cross-sectional study to establish the profile of vitamin D concentration in patients on hemodialysis and supplementation in public centers in Abidjan (Côte d'Ivoire), it appears that the proportion of patients who are deficient is low, and could be improved by greater accessibility to products leading to better compliance. These results highlight the interest of regular systematic supplementation in Ivorian hemodialysis subjects.

ETHICAL CONSIDERATIONS

The study was approved by the local ethical committee of the Ministry of Health. A free and informed consent form was obtained from all participants.

REFERENCES

1. Tissandié E, Guéguen Y, A. Lobaccaro JM, Aigueperse J, Souidi M. Vitamin D: Metabolism, regulation and associated diseases. *Med Sci (Paris)*. 1 Dec 2006; 22(12):1095-1100.
2. Gérard C, Augustin DS, Roger KA, Aida LMH, Gaoussou S, Hien KM, et al. Epidemiological Profile of Chronic Hemodialysis Patients in Ouagadougou. *Open Journal of Nephrology*. 25 avr 2016; 6(2):29-36.
3. Coresh J, Astor BC, Greene T, Eknoyan G, Levey AS. Prevalence of chronic kidney disease and decreased kidney function in the adult US population: Third National Health and Nutrition Examination Survey. *Am J Kidney Dis*. janv 2003; 41(1):1-12.
4. Villarroel M, Jorge J, Meredith D, Sutherland S, Pugh C, Tarassenko L. Non-contact vital-sign monitoring of patients undergoing haemodialysis treatment. *Sci Rep*. 28 oct 2020; 10(1):18529.
5. Ngoie SM, Mulenga P, Mukuku O, Kakisingi CN, Sangwa CM, Nawej PT, et al. Chronic kidney disease: associated factors, etiologies, clinical and biological characteristics in Lubumbashi in the Democratic Republic of Congo. *The Pan African Medical Journal*. 2017 Sep 15;28:41.
6. SOHOU S. Factors favouring CKD: study of 855 CKD cases admitted to the nephrology department of the YOPOUGON University Hospital from April 3, 1991 to December 31, 1996 [Internet]. [cited 2024 Oct 17]. Available on: <https://biblio.uvci.edu.ci/records/369vq-dr562>
7. Collins AJ, Foley RN, Herzog C, Chavers B, Gilbertson D, Ishani A, et al. United States Renal Data System 2008 Annual Data Report. *Am J Kidney Dis*. janv 2009; 53(1 Suppl):S1-374.
8. Goodkin DA, Young EW, Kurokawa K, Prütz KG, Levin NW. Mortality among hemodialysis patients in Europe, Japan, and the United States: case-mix effects. *Am J Kidney Dis*. nov 2004; 44(5 Suppl 2):16-21.
9. Halle MP, Takongue C, Kengne AP, Kaze FF, Ngu KB. Epidemiological profile of patients with end stage renal disease in a referral hospital in Cameroon. *BMC Nephrol*. 21 avr 2015;16:59.
10. Epidemiology and etiology of chronic kidney disease [Internet]. 2001 [cited 2024 Oct 17]. Available on: <https://www.larevuedupraticien.fr/archive/epidemiologie-et-etologie-de-linsuffisance-renale-chronique>

11. Harraqui RE, Naima A, Yassamine B, Haddiya I. Strategy for the management of arterial hypertension in chronic hemodialysis: an applied model of therapeutic patient education (TPE). *The Pan African Medical Journal* [Internet]. 2014 Sep 25 [cited 2024 Oct 17]; 19(86). Available on: <https://www.panafrican-med-journal.com//content/article/19/86/full>
12. Dammak N, Chaabouni Y, Kammoun F, Charfi A, Bouhamed L, Azaiz A, et al. Vitamin D supplementation in hemodialysis patients: a crossover clinical trial comparing cholecalciferol D3 and ergocalciferol D. *Nephrology & Therapeutics*. 1 Sep 2018; 14(5):320.
13. Cavalier E, Sagou Yayo E, Attoungbre-Hauhouot ML, Konan JL, Yao-Yapo C, Monnet D, et al. Vitamin D, bone alkaline phosphatase and parathyroid hormone in healthy subjects and haemodialysed patients from West Africa: impact of reference ranges and parathyroid hormone generation assays on the KDIGO guidelines. *Clin Kidney J*. avr 2019; 12(2):288-93.
14. Moreau E, Bächer S, Mery S, Le Goff C, Piga N, Vogeser M, et al. Performance characteristics of the VIDAS® 25-OH Vitamin D Total assay - comparison with four immunoassays and two liquid chromatography-tandem mass spectrometry methods in a multicentric study. *Clin Chem Lab Med*. janv 2016; 54(1):45-53.
15. Brooks SPJ, Ratnayake WMN, Rondeau I, Swist E, Sarafin K, Weiler HA. Inadequate vitamin D status is associated with lower food plus supplemental intake of vitamin D in children of South Asian ethnicity living in the National Capital Region of Canada. *Appl Physiol Nutr Metab*. 13 sept 2021; 1-8.
16. Boyvin L, Seri K. 25 (OH) Vitamin D level and calcium/phosphorus metabolism disorders in patients living with HIV in Abidjan. *Int J Biochem Res Rev*. 2017; 17:1-7.
17. Lee SA, Stein HH. Effects of dietary levels of calcium, phosphorus, and 1-alpha-hydroxycholecalciferol on digestibility, retention of calcium and phosphorus, and concentration of metabolizable energy in diets fed to sows in late-gestation. *Canadian journal of animal science*. 1 janv 2021; 102(1):184-8.
18. Heike A Bischoff-Ferrari 1, Walter C Willett, John B Wong, Andreas E Stuck, Hannes B Staehelin, E John Orav, Anna Thoma, Douglas P Kiel, Jana Henschkowski Prevention of nonvertebral fractures with oral vitamin D and dose dependency: a meta-analysis of randomized controlled trials, *Arch Intern Med* 2009 Mar 23; 169(6):551-61. doi: 10.1001/archinternmed.2008.600.