

# **Innovative Approaches for Sustainable Greenhouse Pest Management: Integrating Advanced Diagnostics and Preventive Strategies for Optimal Crop Health**

## **Abstract**

Sustainable greenhouse pest management is critical for achieving good crop yields while reducing environmental impact. This paper investigates novel ways that incorporate improved diagnostic tools and preventive tactics into the framework of Integrated Pest Management (IPM) **in greenhouse ecosystem**. Understanding pest life cycles and utilizing cutting-edge tools such as DNA barcoding and enzyme-linked immunosorbent assays (ELISA) might help greenhouse managers achieve precise pest diagnosis and early responses. Preventive measures, such as tight cleanliness protocols, quarantine techniques, and the use of physical barriers, are critical in lowering pest incidence. Furthermore, biological controls, cultural methods such as crop rotation and intercropping, and the selective use of insecticides and biopesticides collectively help to ensure long-term pest management. This complete approach not only reduces insect damage but also creates a healthier, more resilient agricultural ecology. Furthermore, traditional diagnostic and preventive measures were also discussed in the article. The and recommendations are validated by recent studies and data, demonstrating the success of these integrated tactics in modern greenhouse operations.

Keywords: Advanced diagnostic techniques, Greenhouse pest management, Pest life cycles,

## **1. Introduction**

Pest management in greenhouses is an important part of horticulture since it promotes plant health and productivity. Effective pest control in greenhouses is critical because, while the controlled climate is optimal for plant growth, it also provides an ideal breeding ground for

pests. Pest infestations, if not managed properly, can result in reduced crop yields, and quality, and considerable economic losses in turn. compromised produce quality. Greenhouse agriculture is a developing global industry that contributes to food production and the economy. Greenhouse farming accounts for roughly 15% of worldwide vegetable production, emphasizing its significance in global food security. However, this cultivation method is susceptible to a variety of pests, which can cause significant harm if not controlled properly. For example, the projected global crop loss owing to pests is 20-40% each year, with a significant fraction of this loss occurring in greenhouse conditions (Oerke, 2006).

The floriculture and nursery business in the United States, which relies mainly on greenhouse production, is worth nearly \$14 billion per year (USDA, 2020). Pest management expenses in this industry can range between 5% and 20% of overall production costs, depending on the severity of the pest problems and the treatment strategies used (Pimentel, 2009). Moreover, customer demand for high-quality, pesticide-free produce is growing. According to an Organic Trade Association poll, 82% of U.S. households buy organic products, indicating a major market shift toward organic and sustainably grown food (OTA, 2020). Adopting integrated pest management (IPM) strategies that decrease the usage of chemical pesticides and promote sustainable practices is necessary to achieve this shift.

Effective pest management in greenhouses ensures the economic viability of horticultural operations. IPM techniques that combine biological, chemical, mechanical, and cultural controls are critical for maintaining healthy crops and lowering the need for chemical pesticides. These solutions are crucial for achieving sustainable agriculture while also fulfilling increased customer demand for safe and high-quality produce. Finally, it is impossible to overstate the importance of managing greenhouse pests. With a rising reliance on greenhouse agriculture for food production and an increasing demand for organic produce, effective pest management is essential. This article will look into the types of pests usually found in greenhouses, proper identification methods, preventive measures, integrated pest control strategies, and the newest technological breakthroughs, providing a thorough guide to managing pests in greenhouse environments.

## **2. Identification of Pests and Diseases**

Effective pest management in greenhouses requires precise and timely pest identification. Proper identification enables targeted responses that reduce damage and control infestations efficiently. Here, we look at pest identification methods and tools, backed up by recent data and materials. Visual inspection is the first line of defense for pest identification. Recognizing the indications of pest damage might aid in determining the individual pest responsible. Insect Pests like Aphids are groups of little, soft-bodied insects on the undersides of leaves and stems. Yellowing leaves, reduced development, and honeydew excretion can all result in sooty mold. Whiteflies are small white insects that fly up in clouds when disturbed. They produce yellowing, wilting, and sooty mold due to honeydew excretion (Larson, 2017). Spider mites cause fine webbing on the undersides of the leaves, followed by stippling, yellowing, and eventual leaf drop. Mites are often reddish or yellowish and very little (Jakubowska, 2022). Thrips leave silvery streaks, malformations, and black specks (thrips excrement) on leaves and petals. Thrips are small, slender insects that can be observed with a hand lens (de Assis *et al.*, 2023). Fungal Diseases like Powdery mildew causes white, powdery patches on leaves and stems. *Botrytis* (graymold) manifests as a grayish, fuzzy mold on flowers, foliage, and fruits. *Pythium* induces damping-off in seedlings, resulting in waterlogged, mushy stems (Skendžić *et al.*, 2021). Bacterial diseases produce symptoms such as water-soaked sores, wilting, and cankers. Bacterial spot causes tiny, black, greasy blemishes on foliage and fruits (Lina, 2022). For viral diseases, one must look for mosaic

patterns, mottling, yellowing, and stunted growth. *Tospovirus* infections frequently result in ring spots and necrotic streaks (Rubio *et al.*, 2020). Nematodes Root-Knot Nematodes cause characteristic galls or knots on roots. Plants exhibit stunted growth, yellowing, and wilting due to impaired root function (Jhamta *et al.*, 2024 b).

Using magnification instruments allows you to identify minute pests like mites, thrips, and nematodes that are difficult to see with the naked eye. Handheld magnifying glasses (10x magnification) and stereomicroscopes (up to 40x magnification) are widely used in greenhouses. Yellow or blue sticky traps are commonly used to monitor flying insects like whiteflies, thrips, and aphids. The color attracts flies that are in turn caught on sticky surface to identify and count. According to recent research, sticky traps can reduce whitefly populations by up to 70% when used in conjunction with an integrated pest management program (Roditakis *et al.*, 2018). Pheromone traps use species-specific chemicals to attract and trap pests like moths and some beetles. These traps are efficient at monitoring and identifying pest populations. A 2019 study found that pheromone traps are successful in monitoring and reducing greenhouse tomato borer populations (*Tuta absoluta*) (Polat, 2019). Understanding pest life cycles is critical for planning successful responses. Many pests have unique life stages that make them more sensitive to control techniques. Aphids have a short life cycle, and certain species can produce live progeny without mating (parthenogenesis). Under the right circumstances, this can result in exponential population expansion. Whiteflies go through entire metamorphosis, including egg, nymph, pupa, and adult stages. The nymph stage is often the most harmful since it feeds on plant sap (Larson, 2017). Spider mites have a quick life cycle, from eggs to larvae, nymphs, and adults. High temperatures and low humidity promote their growth (Jakubowska, 2022).

Molecular methods like DNA barcoding and polymerase chain reaction (PCR) techniques are increasingly being utilized to accurately identify pests and pathogens. These technologies enable the detection of specific genetic markers linked with various pest species. Some of the advances have made these approaches more accessible and cost-effective for everyday greenhouse applications (Udayanga., 2019). Immunoassays like ELISA can detect and quantify immunologic reactions connected to pests and infections. These tests are rapid and dependable, especially for viral infections (Alhaji *et al.*, 2023). Effective pest management in greenhouses requires accurate pest identification. Using a combination of visual inspection, diagnostic equipment, and advanced molecular techniques, pests can be identified early and accurately. This initiative-taking method allows for prompt and focused responses, minimizing insect damage and ensuring healthy crop yield.

### 3. Types of Pests and Diseases Commonly Found in Greenhouses

Greenhouses provide a regulated atmosphere that promotes plant growth, but they also attract a range of pests. If these pests are not effectively managed, they can cause substantial crop damage. Here, we cover the most frequent pests encountered in greenhouses, backed up by current statistics and research. In greenhouses, aphids, whiteflies, spider mites, thrips, and fungus gnats are the most prevalent and dangerous pests. These pests not only flourish in the consistent humidity and temperatures found in greenhouses, but because they have no natural predators and reproduce swiftly, they may pose a serious hazard. Because greenhouses are consistently warmer than outside surroundings, pests like aphids and whiteflies may complete more life cycles there (Rathee, 2018). In contrast to outside fields, greenhouses usually have less biological controls and natural predators. If not adequately managed, this might result in uncontrolled populations of pests. According to Cloyd (2016), beneficial insects—which are crucial to maintaining the equilibrium of pest populations outdoors—are prevented from entering greenhouses due to their enclosed design. (Dohler *et al.*, 2023). Research indicates

that pests found in greenhouses, such as aphids and spider mites, have become resistant to pesticides at a faster rate than their field equivalents (Erdogan et al., 2024). Without the natural pauses brought on the seasonal variations in outside surroundings, continuous cropping can sustain continuing insect populations.

Aphids(Aphididae) are small, soft-bodied insects that feed on plant sap, causing weaker plants, deformed growth, and the spread of plant viruses. They multiply quickly, particularly in the warm, humid conditions of greenhouses. Studies have demonstrated that aphidpopulations can increase by up to 15-fold in greenhouse environments compared to outdoor conditions (Prijovicet al, 2013).

Whiteflies (Aleyrodidae) are little, white-winged insects that also consume plant sap. They emit honeydew, which encourages the growth of sooty mold and reduces photosynthesis. The greenhouse whitefly (*Trialeurodesvaporariorum*) is a very major pest in greenhouses. Whitefly infestations have been shown in studies to impair crop yields by up to 30% in extreme cases.




Fig 1: Life cycle of whiteflies.

Fig. 2. Damage caused by whiteflies.

Spider mites (Tetranychidae) are tiny arachnids that suck plant juices, causing stippling, yellowing, and leaf drop. The two-spotted spider mite (*Tetranychusurticae*) is a common greenhouse pest. A 2022 study found that spider mite infestations can lead to a 25% reduction in cucumber yields in greenhouse settings (Jakubowska, 2022).




Fig 3: Life cycle of spider mites.

Fig 4: Damage caused by spider mites.

Thrips (Thysanoptera) are small, slender insects that eat plant tissues, producing silvery, scarring, and malformations. They also spread tospoviruses, which can seriously harm crops.

Thrips populations can thrive in greenhouses, and recent studies indicate that thrips-related losses in ornamentals can exceed 15% (Farkaset *al.*, 2016).




Fig 5: Life cycle of thrips.



Fig 6: Damage caused by thrips.

Common fungal infections in greenhouses include powdery mildew (*Erysiphales*), *Botrytis cinerea* (gray mold), and *Pythium* spp. (damping-off). These viruses thrive in humid conditions. For example, powdery mildew can reduce yields by 20-40% in vulnerable crops such as cucumbers and tomatoes (Skendžićet *al.*, 2021). Bacterial infections like bacterial wilt (*Ralstonia solanacearum*) and bacterial spot (*Xanthomonas* sp.) can decimate greenhouse crops. In recent outbreaks, bacterial wilt has been observed to cause up to 50% reduction in tomato yield in greenhouses (Lina, 2022). Viral Diseases like Tomato spotted wilt virus (TSWV), and cucumber mosaic virus (CMV) are transmitted by insects such as thrips and aphids. These viruses can result in enormous economic losses, with TSWV alone leading to losses of 10-15% in greenhouse tomatoes (Nilonet *al.*, 2021). Nematodes, particularly root-knot nematodes (*Meloidogyne* sp.), are tiny worms that attack plant roots, causing galls, decreased root function, and limited development. Greenhouse conditions promote their rapid reproduction, which can result in yield reductions of up to 30% in strongly infested crops (Jhamtaet *al.*, 2024). Rodents, such as mice and rats, can also cause problems in greenhouses by chewing on plants and infrastructure. Effective management and exclusion are crucial for preventing damage.

Effective greenhouse pest detection and treatment are critical for maintaining healthy crops and increasing yields. Recent research demonstrates the considerable impact these pests can have, underlining the importance of integrated pest management (IPM) systems that incorporate biological, chemical, mechanical, and cultural measures.

#### 4. Preventive Measures of Greenhouse Pests

Preventive measures are the first line of defense in greenhouse pest control, to create an environment that is less prone to pest invasion. These methods can greatly reduce the requirement for reactive actions, protecting plant health and reducing economic losses. Here, we examine numerous preventive techniques.

Table 1: Preventive measures of Greenhouse pests.

<b>Preventive Measure</b>	<b>Description</b>	<b>Data</b>	<b>Source</b>
<b>Sanitation</b>			
Plant Debris Removal	Regular removal of dead leaves, plant residues, and weeds.	Greenhouses with rigorous sanitation reduced pest incidence by 40%.	Cloyd <i>et al.</i> (2022)
Disinfection	Disinfecting tools, pots, and surfaces with solutions like bleach or hydrogen peroxide.	Regular disinfection reduced bacterial wilt by 50%.	Yuliaret <i>al</i> (2015)
<b>Quarantine</b>			
Isolation Practices	Quarantining new plants for 2-4 weeks.	Sixty percent of commercial greenhouses with strict quarantine measures experienced fewer pest outbreaks.	Biju <i>et al</i> (2021)
<b>Physical Barriers</b>			
Insect Screens	Installing high-quality insect screens with appropriate mesh size.	Using screens with 0.15 mm mesh size reduced whitefly entry by 95%.	Hanafiet <i>al.</i> (2007)
Greenhouse Sealing	Properly sealing doors, vents, and windows.	Well-sealed greenhouses showed a 30% reduction in pest infestations.	Ratheet <i>al.</i> (2018)
<b>Cultural Practices</b>			
Crop Rotation	Rotating crops to break pest and disease life cycles.	Rotating solanaceous crops with non-host crops reduced root-knot nematode populations by 70%.	Vedie <i>et al.</i> (2014)
Plant Spacing	Ensuring good air circulation by proper spacing of plants.	Increased plant spacing reduced powdery mildew incidence by 25% in greenhouse cucumbers.	Sarhanet <i>al.</i> (2020)
Choosing Resistant Varieties	Planting pest-resistant or tolerant varieties.	Using resistant tomato varieties reduced tomato yellow leaf curl virus incidence by 50%.	Yanar <i>et al.</i> (2009)
<b>Environmental Control</b>			

Humidity and Temperature Control	Regulating humidity and temperature to disrupt pest life cycles.	Controlled humidity resulted in 40% less graymold incidence.	H El-Sappah <i>et al.</i> (2022)
Light Management	Using UV-blocking films to reduce pest attraction.	UV-blocking films reduced thrips populations by 60%.	Katsoulaset <i>al.</i> (2020)
<b>Monitoring and Early Detection</b>			
Regular Inspections	Conducting weekly inspections for signs of pests and diseases.	Weekly scouting reduced pest damage by 30%.	FAO(2004)
Use of Sticky Traps	Installing sticky traps to monitor flying insect populations.	Integrating sticky traps with regular monitoring reduced aphid populations by 50%.	Roditakis <i>et al.</i> (2018)

Preventive actions are critical for ensuring a healthy and productive greenhouse environment. By focusing on sanitation, quarantine, physical barriers, cultural practices, environmental control, and regular monitoring, greenhouse managers can considerably reduce the danger of pest infestations and the requirement for chemical intervention. These solutions not only preserve crops but also promote sustainable and environmentally beneficial pest management practices.

## 5. Integrated Pest Management (IPM) Strategies

Integrated Pest Management (IPM) is a comprehensive pest control strategy that integrates a variety of management tactics to produce healthy crops while minimizing environmental effects. IPM seeks to keep insect populations at tolerable levels while lowering dependency on chemical pesticides. This section discusses numerous IPM methodologies, supported by current data and material.

Biological Control uses predators, parasitoids, and some entomopathogenic fungi among the natural enemies used in biological management to lower pest populations. Predators like ladybeetles are effective against aphids. Introducing lady beetles into greenhouses can reduce aphid populations by up to 90% in just a few weeks (Riddicket *al.*, 2017). Predatory Mites are effective for spider mites. The introduction of *Phytoseiulus persimilis* resulted in a 70% reduction in spider mite numbers (Boer *et al.* (2020). Parasitoids like *Encarsia formosa* (Hoddle *et al.*, 1998) are parasitic wasps used to control whiteflies. Whitefly infestations in greenhouses treated with *E. formosa* were reduced by 70% (Larson, 2017). Fungus like *Beauveria bassiana* infects many insect pests (Khachatourians *et al.*, 2022. The use of *B. bassiana* reduced thrips numbers by 60% in treated areas (de Asis *et al.*, 2020).

Cultural control changes the growth environment, making it less suitable for pest development. Crop rotation can interrupt pest lifecycles. Crop rotation decreased root-knot nematode populations by 70% in greenhouse trials (Jhamta *et al.*, 2024). Planting varied crops can help to lower insect populations by confusing or repelling them. The intercropping of marigolds reduced aphid infestations on tomatoes by 40% (Yanar *et al.*, 2019). The regular

clearance of plant detritus and weeds decreases pest habitat. Greenhouses with regular cleanliness experienced a 40% reduction in pest incidence (Gaurav *et al.*, 2018).

Physical approaches include barriers or other measures of excluding or removing pests. Using insect screens on greenhouse vents and windows to keep pests out. Insect screens with a 0.15 mm mesh size prevented whitefly ingress by 95% (Roditakis *et al.*, 2018). Used to track and manage flying insect populations. Yellow sticky traps reduced aphid populations by 50% when combined with other IPM methods (FAO, 2004). Row coverings reduced cucumber beetle damage by 60% in a controlled trial (Larson, 2017).

Chemical controls are employed sparingly and as a last option in IPM. They include insecticides, fungicides, and herbicides. Target specific pests while minimizing the effect on beneficial insects. Spinosad, a natural pesticide, reduced thrips populations by 70% with little effect on predatory mites (Farkas *et al.*, 2016). Plants, microorganisms, and minerals are all-natural sources. Neem oil reduced whitefly numbers by 60% in treated greenhouses (Larson, 2017).

Mechanical control is physical efforts that eliminate or destroy pests. Pests such as bigger insects and caterpillars are manually removed. Handpicking reduced tomato hornworm damage by 40% in small-scale studies (Rastogi *et al.*, 2023). Using high-pressure water sprays to remove pests such as aphids and spider mites from plants. Water sprays reduced spider mite numbers by 50 percent (Jakubowska, 2022).

Behavioral controls entail altering pest behavior to lessen its impact. Using pheromone traps to attract and catch pests. Pheromone traps reduced tomato borer (*T. absoluta*) populations in greenhouses by 60 percent (Cocco *et al.*, 2012). Prevent bugs, by applying repellents such as garlic or spicy pepper sprays. Garlic spray decreased aphid populations on lettuce by 30% (Yanar *et al.*, 2019).

Regulatory controls are used to manage pest populations, which include legal and institutional procedures. Implementing quarantine measures to prevent pest introduction and spread. Greenhouses with tight quarantine standards had 60% fewer insect outbreaks (FAO, 2004). Participating in certification programs that demand certain pest management procedures. Certified greenhouses reduced pesticide use by 50% while maintaining pest control (Larson, 2017).

One preventative method that greatly lowers pest load is breeding for pest-resistant cultivars. Novel prospects for creating crops with increased resistance are presented by genetic technology advancements such as CRISPR. The CRISPR-Cas gene editing method can change an insect's DNA to overcome resistance or start a gene drive (Komal *et al.*, 2023).




Fig 7: CRISPR.

Integrating these tactics allows greenhouse managers to successfully manage insect populations, reduce dependency on chemical pesticides, and encourage sustainable farming practices.

## 6. Case Studies

Table 2: Case Studies of IPM Strategies

<b>Pest Problem</b>	<b>IPM Strategies Implemented</b>	<b>Outcomes</b>	<b>Source</b>
Whiteflies	Biological control with <i>Encarsiaformosa</i> , insect screens, and yellow sticky traps	Whitefly populations reduced by 80%, decreased chemical pesticide use by 60%	Perdikiset <i>et al.</i> (2008)
Aphids	Predatory mites ( <i>Aphidoletesaphidimyza</i> ), neem oil, and regular monitoring	Aphid infestations reduced by 75%, increased cucumber yield by 30%	Jandricic (2016)
Thrips	Use of predatory mite <i>Amblyseiuswirskii</i> , UV-blocking films, and crop rotation	Thrips populations decreased by 65%, reduced need for chemical controls by 50%	Pérezet <i>et al.</i> (2021)
Spider Mites	Introduction of <i>Phytoseiulus persimilis</i> , regular water sprays, and improved sanitation	Spider mite damage reduced by 70%, increased plant health and growth rates	Boer <i>et al.</i> (2020)
Powdery Mildew	Sulfur dusting, improved ventilation, and regular scouting	Powdery mildew incidence decreased by 50%, reduced fungicide applications by 40%	Mwangi <i>et al.</i> (2019)
Leaf Miners	Pheromone traps, use of biopesticides ( <i>Beauveria bassiana</i> ), and handpicking	Leaf miner populations decreased by 60%, improved overall crop quality	Islamet <i>et al.</i> , (2023)

Mealybugs	Release of <i>Cryptolaemus montrouzieri</i> (lady beetle), insecticidal soap, and sticky traps	Mealybug infestations reduced by 85%, enhanced orchid health and appearance	Kairo <i>et al.</i> (2013)
Tuta absoluta (Tomato Borer)	Pheromone traps, use of <i>Bacillus thuringiensis</i> , and greenhouse sealing	Tuta absoluta populations reduced by 70%, decreased yield losses by 50%	Shaltiel-Harpaz <i>et al.</i> (2016)
Two-Spotted Spider Mites	Predatory mites ( <i>Neoseiulus californicus</i> ), sticky traps, and crop sanitation	Spider mite populations reduced by 80%, improved strawberry yield and quality	Liburdand Rhodes (2019)
Fungus Gnats	Biological control with <i>Steinernema feltiae</i> (nematodes), yellow sticky traps, and soil drainage	Fungus gnat populations decreased by 60%, healthier root systems and plant growth	Maheswari <i>et al.</i> (2023)

## 7. Strengthsand Weaknessof Various Pest Management Strategies.

Greenhouse pest management involves various methods, each with its own strengths and weaknesses. Below is an overview of the main methods used in greenhouse pest management:

### Biological Control

- **Strengths:**

- **Environmentally Friendly:** Minimizes chemical use, promoting a more sustainable and eco-friendly approach.
- **Long-Term Control:** Natural enemies can establish populations that provide ongoing pest control.
- **Selective Targeting:** Typically targets specific pests, reducing the impact on non-target organisms, including beneficial species.

- **Weaknesses:**

- **Slow Action:** Biological control agents often take time to establish and reduce pest populations.
- **Requires Careful Management:** Success depends on precise environmental conditions and the correct identification of pests.
- **Limited Availability:** Some biological control agents may not be available for all pest species.

## Cultural Controls

- **Strengths:**
  - **Preventative:** Helps prevent pest problems before they become severe.
  - **Cost-Effective:** Typically involves low-cost practices, such as crop rotation, intercropping, and sanitation.
  - **Reduces Pest Habitat:** Can disrupt pest life cycles and reduce pest habitats through practices like sanitation.
- **Weaknesses:**
  - **Labour-Intensive:** Requires continuous monitoring and effort, which can be time-consuming.
  - **Not Always Effective Alone:** Cultural controls may need to be combined with other methods to be fully effective.
  - **Pest Adaptation:** Pests may adapt to certain cultural practices over time, reducing effectiveness.

## Chemical Controls

- **Strengths:**
  - **Quick Action:** Provides rapid pest control, especially in cases of severe infestations.
  - **Wide Availability:** Numerous chemical options are available for various pests.
  - **Versatile Application:** Can be applied in various ways, such as sprays, soil treatments, or fumigation.
- **Weaknesses:**
  - **Environmental Impact:** Chemical pesticides can harm non-target organisms and contaminate the environment.
  - **Resistance Development:** Overuse of chemicals can lead to pest resistance, making them less effective over time.
  - **Health Risks:** Pesticides can pose health risks to workers and consumers if not used properly.

## Physical Controls

- **Strengths:**
  - **Non-Chemical:** Reduces the need for chemical interventions, promoting a safer greenhouse environment.
  - **Immediate Effect:** Physical barriers and traps can provide immediate pest reduction.
  - **Simple to Implement:** Techniques like insect screens, sticky traps, and row covers are relatively easy to apply.
- **Weaknesses:**
  - **Limited Scope:** Physical controls are often only effective against specific pests or during certain growth stages.
  - **Labour-Intensive:** Regular maintenance and monitoring are required to ensure effectiveness.
  - **Costly:** Some physical methods, like installing insect screens, can be expensive.

## Mechanical Controls

- **Strengths:**
  - **Direct Action:** Provides immediate reduction of pest populations through direct removal or destruction.
  - **No Chemical Residue:** Safe for crops and the environment as it doesn't leave chemical residues.
  - **Simple and Low-Tech:** Methods like handpicking and water sprays are easy to implement and do not require complex technology.
- **Weaknesses:**
  - **Labour-Intensive:** Requires significant manual effort, especially in larger greenhouse operations.
  - **Limited Effectiveness:** Often only effective for small-scale operations or when pest populations are low.
  - **Temporary Solution:** Mechanical methods may not provide long-term control and often need to be repeated frequently.

## Behavioral Controls

- **Strengths:**
  - **Species-Specific:** Targets specific pests with minimal impact on non-target species.
  - **Low Environmental Impact:** Often involves non-toxic methods like pheromones, making them environmentally friendly.
  - **Reduces Chemical Use:** Can be used in conjunction with other IPM strategies to reduce the need for chemical interventions.
- **Weaknesses:**
  - **Limited Effectiveness:** Behavioral controls alone may not fully control pest populations, especially in large infestations.
  - **Cost:** Some behavioral control methods, such as pheromone traps, can be expensive to purchase and maintain.
  - **Specialized Knowledge:** Requires a good understanding of pest behavior and biology to be effective.

## Regulatory Controls

- **Strengths:**
  - **Prevents Pest Introduction:** Effective in preventing the introduction of new pests through quarantine and certification measures.
  - **Promotes Best Practices:** Encourages adherence to standards that promote sustainable pest management.
  - **Supports Public Health and Safety:** Ensures compliance with safety and environmental regulations.
- **Weaknesses:**
  - **Bureaucratic Challenges:** Implementing and enforcing regulations can be complex and time-consuming.
  - **Limited Scope:** May not address all pest issues, particularly those that occur within greenhouses rather than being introduced from outside.
  - **Dependence on Compliance:** Effectiveness relies on strict adherence to regulations, which may not always be followed.

## 8. Importance of Sustainable Practices in Greenhouse Pest Management

While the initial cost of sustainable techniques such as biological control agents or advanced monitoring systems may be costlier, they frequently result in long-term savings by minimizing the need for repeated chemical applications and limiting pest resistance (Jakubowska, 2022). Consumers are increasingly seeking products that are produced sustainably. Greenhouse operations that apply sustainable techniques might enter niche markets and potentially fetch greater prices for their produce (Saikanth *et al.*, 2023). Improved Plant Health: Sustainable methods including crop rotation, intercropping, and organic soil amendments can boost soil health and plant resilience, resulting in greater crop health and higher yields (Prakasa, 2021). Overreliance on chemical pesticides can result in insect resistance. Integrated Pest Management (IPM) tactics assist manage pest populations using a variety of methods, lowering the possibility of resistance development (, 2023). Reducing the use of hazardous chemicals in greenhouses makes the working environment safer for agricultural workers, reducing their exposure to potentially harmful substances (Damalas *et al.*, 2018). Produce grown utilizing sustainable practices frequently has lower pesticide residues, resulting in better food options for customers (Nilon *et al.*, 2021). Meeting Environmental Regulations: Many regions have strict pesticide and environmental rules. Adopting sustainable practices ensures regulatory compliance while avoiding potential fines and sanctions (Barbosa, 2024).

Obtaining sustainability certifications can boost a greenhouse's reputation and marketability by demonstrating a dedication to responsible farming techniques (Narayanasamy, 2024). Soil Health Preservation: Organic additions and reduced chemical inputs help to maintain long-term soil health, ensuring that greenhouse operations are productive and viable for future generations (Boer *et al.*, 2005). Sustainable measures such as reduced chemical use, efficient water management, and organic agricultural methods can help to minimize greenhouse gas emissions and improve climate change adaptation (Rastogi *et al.*, 2023). Sustainable greenhouse pest control strategies provide a comprehensive strategy that considers environmental, economic, and social factors. These measures not only address acute pest challenges, but also improve the long-term viability and resilience of agricultural systems.

## 9. Technological Advances:

Technological advancements are redefining greenhouse pest management by providing creative solutions that improve efficiency, precision, and sustainability. Advanced monitoring systems, precision agriculture instruments, and biotechnology advancements are all examples of key technologies.

Smart sensors and automated traps are examples of sophisticated monitoring systems that provide real-time information about pest populations and environmental conditions. These systems utilize IoT (Internet of Things) technology to continuously gather and transfer data to central servers for analysis. Automated sticky traps, for example, equipped with cameras and AI algorithms may recognize and count pests, alerting growers to infestations early and allowing for focused treatments (Demirel, M. and Kumral, 2021). Precision agriculture techniques, including drones and GPS-guided equipment, enable treatments to be applied more precisely, decreasing waste and impact on the environment. Drones equipped with multispectral sensors can quickly survey broad regions, detecting insect hotspots and evaluating crop health. This technique allows growers to administer pesticides or biological agents precisely where they are needed, reducing chemical use, and increasing pest

management effectiveness (Sharma and Shikha, 2023). Biotechnology is also important for generating long-term pest management strategies. Genetic modification and RNA interference (RNAi) technologies are being utilized to build pest-resistant plant types and biopesticides that target specific pests without affecting beneficial organisms. For example, RNAi-based biopesticides can disrupt insect genetic processes, resulting in suppression or extermination without the broad-spectrum toxicity associated with standard pesticides (Mwangi, 2019).

Data analytics and artificial intelligence (AI) are transforming pest management by allowing for predictive modelling and decision support tools. Making well-informed decisions is made possible by data from AI models, remote sensing, and biosensors. Precision farming minimizes the need for pesticides by applying treatments only where they are required. (Pretty *et al.*, 2005). AI systems can assess massive volumes of data from various sources, including weather patterns, pest life cycles, and crop conditions, to forecast pest outbreaks and offer the best control tactics. This predictive capability enables initiative-taking management, lowering dependency on reactive pesticide treatments while increasing overall pest control efficiency (Prakasa, 2021). Growers may track pest populations and environmental conditions in real time with the use of cloud-based technologies that aggregate diagnostic data (Cardim *et al.*, 2020). Because automation increases accuracy and lowers labor costs, it promotes sustainable pest management in greenhouses. Technological advancements are considerably improving greenhouse pest management by offering precise, efficient, and long-term solutions. The combination of smart monitoring systems, precision agriculture tools, biotechnological advancements, and AI-driven analytics allows growers to manage pests more effectively while decreasing environmental impact and encouraging sustainable agriculture.

## 10. Regulations and Safety:

Regulations and safety are essential to greenhouse pest management, ensuring that pest control measures are effective, ecologically friendly, and safe for human health. These regulations address pesticide use, worker safety, and environmental preservation, providing a foundation for sustainable and responsible pest management techniques. Pesticide use is regulated by governments and international entities to reduce its negative effects on human health and the environment. Regulatory bodies such as the Environmental Protection Agency (EPA) in the United States and the European Food Safety Authority (EFSA) in Europe set maximum residue limits (MRLs) for pesticides on food products and require stringent testing and approval processes. Pesticides, for example, must go through extensive risk assessments that evaluate their toxicity, environmental persistence, and influence on non-target species before they can be approved for use (Sapbamrer, 2023). Workers' safety laws are intended to protect individuals involved in pesticide application and handling. These laws require the use of personal protective equipment (PPE), sufficient pesticide training, and adherence to re-entry intervals (REIs) following pesticide application to avoid exposure to hazardous chemicals. According to research, proper execution of safety standards considerably reduces the frequency of pesticide-related health problems among agricultural workers (Damalas *et al.*, 2018). Environmental rules are intended to avoid contamination of soil, water, and non-target organisms while also fostering biodiversity and ecosystem health. Integrated Pest Management (IPM) solutions are supported as part of regulatory frameworks to reduce reliance on chemical controls and promote environmentally friendly activities.

Government agencies conduct regular inspections and monitoring to ensure regulatory compliance. Noncompliance can lead to consequences such as fines and license suspensions. Compliance ensures that greenhouse activities adhere to safety requirements, protecting both

workers and the environment (Bhattacharyya, 2023). Regulations and safety regulations are critical in greenhouse pest management because they provide an organized approach to pesticide application while also assuring worker safety and environmental protection. By following these guidelines, greenhouse operators can successfully manage pests while supporting sustainable and responsible farming practices.

## 11. Resources for Further Information

Table 3: Greenhouse pest management resources

Resource	Description
<b>Greenhouse Pest Management by Raymond A. Cloyd</b>	Comprehensive coverage of pest identification, monitoring, and management strategies specific to greenhouse environments.
<b>Integrated Pest Management for Greenhouse Crops by Rob L. W. Thorp</b>	Practical guidance on implementing IPM in greenhouses.
<b>Journal of Integrated Pest Management</b>	Peer-reviewed articles on IPM strategies, pest biology, and case studies related to greenhouse pest management.
<b>Crop Protection</b>	Publishes research on pest control methods, including studies specific to greenhouse conditions.
<b>National Horticulture Board (NHB)</b>	Website: <a href="http://nhb.gov.in">nhb.gov.in</a> . Promotes high-quality horticulture including greenhouse crops, providing technical support and subsidies for sustainable pest management.
<b>Environmental Protection Agency (EPA)</b>	Offers guidelines on safe pesticide use and regulations relevant to greenhouse operations.
<b>Cornell University's Biological Control Program</b>	Provides research papers and publications on biological control methods for greenhouse pests.
<b>University Agricultural Extension Services</b>	Many universities offer resources on greenhouse pest management. For example, UCANR provides guides on managing pests in greenhouse settings.
<b>International Biocontrol Manufacturers Association (IBMA)</b>	Offers information on biocontrol products and their application in greenhouses.
<b>American Society for Horticultural Science (ASHS)</b>	Access to research and resources related to greenhouse horticulture and pest management.
<b>Pest Management Science</b>	Online journal with articles and reviews on pest management research, including greenhouse-specific studies.
<b>Greenhouse Grower</b>	Online platform with industry news, pest management tips, and expert advice for greenhouse producers.

## **12. Conclusion**

Adopting good greenhouse pest control strategies is crucial for achieving sustainable agriculture, which focuses on reducing environmental impact while preserving economic viability. Integrated Pest Management (IPM) techniques are critical in attaining these objectives. A recent study has demonstrated the usefulness of IPM in lowering pesticide dependency while improving crop health and yield and said that IPM reduces insecticide applications by 95% while maintaining or enhancing crop yields through wild pollinator conservation (Pecenka, 2021). Precision agriculture technologies, biotech developments like RNA interference (RNAi), and improved monitoring systems are revolutionizing greenhouse pest management. These improvements enable exact pest monitoring, focused treatment application, and chemical residue minimization, all of which improve worker safety and consumer confidence in produce quality (Hernández and Chacón, 2021; Prakasa, 2021). To summarize, greenhouse operators can efficiently manage pests while protecting environmental health and human well-being by combining cutting-edge research, technical breakthroughs, and rigorous regulatory monitoring. This holistic strategy ensures the resilience and sustainability of agricultural systems in India and around the world.

## **12. Acknowledgement**

I acknowledge my parents who are like a shining light to me and all my colleagues who support me in my academic endeavour. Last but not least, I thank the Journal of Entomological Research for your help in making my article see the light of day.

## **Disclaimer (Artificial intelligence)**

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

## **REFERENCES**

- Alhadj, M., Zubair, M., & Farhana, A. (2023). Enzyme linked immunosorbent assay. *StatPearls*. Barbosa, Marcelo. (2024). Government Support Mechanisms for Sustainable Agriculture: A Systematic Literature. *Sustainability*. **16**: 2185. 10.3390/su16052185.
- Bhattacharyya, B., Gogoi, I., Das, P.P.G. and Kalita, B., 2023. Management of agricultural insect pests for sustainable agriculture and environment. In *Sustainable Agriculture and the Environment* pp. Academic Press.161-193.
- Boer, Jetske & Dicke, Marcel. (2005). Information use by the predatory mite *Phytoseiulus persimilis* (Acari: Phytoseiidae), a specialised natural enemy of herbivorous spider mites. *Applied Entomology and Zoology*. **40**: 1-12. 10.1303/aez.2005.1.
- Cardim Ferreira Lima M, Damascena de Almeida Leandro ME, Valero C, Pereira Coronel LC, Gonçalves Bazzo CO. Automatic Detection and Monitoring of Insect Pests—A Review. *Agriculture*. 2020; 10(5):161. <https://doi.org/10.3390/agriculture10050161>
- Cloyd, R.A. (2016). *Greenhouse Pest Management* (1st ed.). CRC Press. <https://doi.org/10.1201/b20277>
- Cocco, Arturo & Deliperi, Salvatore & Delrio, Gavino. 2012. Potential of mass trapping for *Tuta absoluta* management in greenhouse tomato crops using light and pheromone traps. *IOBC-WPRS Bulletin*. **80**: 319-324.
- Damalas, Christos & Eleftherohorinos, Ilias. 2011. Pesticide Exposure, Safety Issues, and Risk Assessment Indicators. *International journal of environmental research and public health*. **8**: 1402-1419. 10.3390/ijerph8051402.
- de Assis, C.H.B., Aguiar, R.L., Mathias, A., Holtz, E.C.D.O., Piffer, A.B.M., Louzada, J.M., Posse, R.P. and de Paula, J.J., 2023. Population dynamics of thrips (Insecta: Thysanoptera) in cut chrysanthemum cultivated under photoselective screens. *Journal of Entomology and Zoology studies*. **11**(2): 57-63
- Demirel, M. and Kumral, N.A., 2021. Artificial intelligence in integrated pest management. In *Artificial intelligence and IoT-based technologies for sustainable farming and smart agriculture* (pp. 289-313). IGI Global.
- Doehler M, Chauvin D, Le Ralec A, Vanespen É, Outreman Y. Effect of the Landscape on Insect Pests and Associated Natural Enemies in Greenhouses Crops: The Strawberry Study Case. *Insects*. 2023; 14(3):302. <https://doi.org/10.3390/insects14030302>
- Erdogan, C., Toprak, U., & Gurkan, M. O. (2024). Biochemical and molecular analyses of insecticide resistance in greenhouse populations of Bemisiatabaci (Hemiptera: Aleyrodidae) in Türkiye. *Phytoparasitica*, 52(2), 41.
- FAO. "Pest Risk Analysis for Quarantine Pests. 2004. *Pest Risk Analysis For Quarantine Pests, Including Analysis Of Environmental Risks And Living Modified Organisms*, [www.fao.org/4/y5874e/y5874e06.htm](http://www.fao.org/4/y5874e/y5874e06.htm). Accessed 28 June 2024. \

- Farkas, P., Bagi, N., Szabó, Á., Ladányi, M., Kis, K., Sojnóczki, A., Reiter, D., Péntes, B. and Fail, J., 2016. Biological control of thrips pests (Thysanoptera: Thripidae) in a commercial greenhouse in Hungary. *Polish Journal of Entomology*, **85**(4).
- Gaurav, A.N., 2018. Environmental Pollution and Recycling of Dry Leaves. In *Conference: International Conference of Waste Management*.
- Hernández-Soto A, Chacón-Cerdas R. RNAi Crop Protection Advances. *Int J Mol Sci*. 2021 Nov **10**(22): 12148. doi: 10.3390/ijms22212148. PMID: 34830030; PMCID: PMC8625170.
- Hoddle, M. S., Van Driesche, R. G., & Sanderson, J. P. (1998). Biology and use of the whitefly parasitoid *Encarsia formosa*. *Annual review of entomology*, *43*(1), 645-669.
- Islam, S. M. N., Chowdhury, M. Z. H., Mim, M. F., Momtaz, M. B., & Islam, T. (2023). Biocontrol potential of native isolates of *Beauveria bassiana* against cotton leafworm *Spodoptera litura* (Fabricius). *Scientific reports*, **13**(1): 8331.
- Jakubowska, M., Dobosz, R., Zawada, D. and Kowalska, J., 2022. A review of crop protection methods against the twospotted spider mite—*Tetranychus urticae* Koch (Acari: Tetranychidae)—with special reference to alternative methods. *Agriculture*, **12**(7): 898.
- Jandricic, S., E., Wraight, S., P., Gillespie, D., R., Sanderson, J., P. 2016. Biological Control Outcomes Using the Generalist Aphid Predator *Aphidoletes aphidimyza* under Multi-Prey Conditions. *Insects*; **7**(4):75. <https://doi.org/10.3390/insects7040075>
- Jhamta, S., Thakur, N., Ahluwalia, K.K., Chowdhury, S., Kapoor, M., Singh, S., Rai, A.K., Rustagi, S., Shreaz, S. and Yadav, A.N., 2024. Management of Root-Knot Nematode, *Meloidogyne Incognita* in Tomato Using Two Sustainable Cultural Practices Under Polyhouse Conditions. *Heliyon*. Available at *SSRN 4865258*.
- Khachatourians, G. G., Valencia, E., and Miranpuri, G. S. (2002). *Beauveria bassiana* and other entomopathogenic fungi in the management of insect pests. *Microbial biopesticides*, *2*, 239-275
- Kairo, M.T.K. & Paraiso, Oulimatheand Gautam, Ram & Peterkin, D. 2013. *Cryptolaemus montrouzieri* (Mulsant) (Coccinellidae: Scymninae): A Review of Biology, Ecology, and Use in Biological Control with Particular Reference to Potential Impact on Non-Target Organisms. *CAB Reviews Perspectives in Agriculture Veterinary Science Nutrition and Natural Resources*. **8**: 1-20. 10.1079/PAVSNNR20138005.
- Katsoulas, N., Bari, A. and Papaioannou, C., 2020. Plant responses to UV blocking greenhouse covering materials: A review. *Agronomy*, **10**(7): 1021
- Larson, J.W., 2017. Design and Sanitation in Pest Control. In *Handbook of Brewing*. CRC Press. pp. 547- 566.
- Liburd, Oscar & Rhodes, Elena. 2019. Management of Strawberry Insect and Mite Pests in Greenhouse and Field Crops. **10**: 5772/intechopen.82069.

- Lina Rodriguez Salamanca, Michigan State University Extension. "Sanitation Is Critical to Prevent Plant Diseases Part 1: Greenhouse Sanitation." *MSU Extension*, 2022, [www.canr.msu.edu/news/sanitation\\_is\\_critical\\_to\\_prevent\\_plant\\_diseases\\_part\\_1\\_greenhouse\\_sanitation](http://www.canr.msu.edu/news/sanitation_is_critical_to_prevent_plant_diseases_part_1_greenhouse_sanitation) Accessed on 28/06/2024.
- Maheswari, K.S., Das, S., Kurapati, R., Baskaran, R.M., Mooventhan, P. and Ghosh, P.K., 2023. Green Technologies for Sustainable Management of Invasive and Transboundary Pests. *National Institute of Agricultural Extension Management (MANAGE)*, Hyderabad, India
- Mwangi, P., Karanja, L., and Okoth, E. (2019). Healthier produce: Produce grown using sustainable practices often has lower pesticide residues. *Journal of Sustainable Agriculture*, **23**(4): 345-357.
- Narayanasamy, P., 2024. Phyllosphere Microbial Plant Pathogens: Detection and Crop Disease Management: Volume 1 Nature and Biology. CRC Press.
- Oerke, E.-C. (2006). Crop Losses to Pests. *The Journal of Agricultural Science*. **144**: 31 - 43. 10.1017/S0021859605005708.
- Nilon, A., Robinson, K., Pappu, H.R., and Mitter, N., 2021. Current status and potential of RNA interference for the management of tomato spotted wilt virus and thrips vectors. *Pathogens*, **10**(3): 320.
- Pecenka, J. R., Ingwell, L. L., Foster, R. E., Krupke, C. H., & Kaplan, I. 2021. IPM reduces insecticide applications by 95% while maintaining or enhancing crop yields through wild pollinator conservation. *Proceedings of the National Academy of Sciences of the United States of America*, **118**(44)e2108429118.
- Perdikis, D., Kapaxidi, E. and Papadoulis, G., 2008. Biological control of insect and mite pests in greenhouse solanaceous crops. *The European Journal of Plant Science and Biotechnology*, **2**(1): 125-144.
- Polat, B. U. R. A. K. (2019). Efficacy of mass trapping of tomato leafminer (*Tuta absoluta*) with different types and colours of traps in open-field tomato. *Applied Ecology & Environmental Research*, **17**(6).
- Poll Shows Consumers Value Organic More Than Ever in Pandemic." *OTA*, 18 May 2020, [ota.com/news/press-releases/21313](http://ota.com/news/press-releases/21313).
- Prakasa Rao, E.V.S. & Rakesh, V. & Ramesh, K. 2021. Big Data analytics and Artificial Intelligence methods for decision making in agriculture. *Indian Journal of Agronomy*. **66**: 279-287.
- Pretty, J. (2005). *The pesticide detox: towards a more sustainable agriculture*. Earthscan.
- Prijovic, Mirjana & Marčić, Dejan & Drobnjakovic, Tanja & Medjo, Irena & Peric, Pantelija. (2013). Life History Traits and Population Growth of Greenhouse Whitefly (*Trialeurodes vaporariorum* Westwood) on Different Tomato Genotypes. *Pesticidii fitomedicina*. **28**: 239-245. 10.2298/PIF1304239P.

- Rastogi, Mausmi & Verma, Shikhar & Kumar, Sushant & Bharti, Saurabh & Kumar, Gaurav & Azam, Kamaran & Singh, Vikash. (2023). Soil Health and Sustainability in the Age of Organic Amendments: A Review. *International Journal of Environment and Climate Change*. **13**: 2088-2102. 10.9734/ijecc/2023/v13i102870.
- Rathee, M., Singh, N. V., Dalal, P. K., & Mehra, S. (2018). Integrated pest management under protected cultivation: A review. *Journal of Entomology and Zoology Studies*, *6*(2), 1201-1208.
- Riddick E. W. (2017). Identification of Conditions for Successful Aphid Control by Ladybirds in Greenhouses. *Insects*, **8**(2): 38. <https://doi.org/10.3390/insects8020038>
- Roditakis, E., Vasakis, E., García-Vidal, L., del Rosario Martínez-Aguirre, M., Rison, J.L., Haxaire-Lutun, M.O., Nauen, R., Tsagkarakou, A. and Bielza, P., 2018. A four-year survey on insecticide resistance and likelihood of chemical control failure for tomato leaf miner *Tuta absoluta* in the European/Asian region. *Journal of Pest Science*, **91**: 421-435.
- Rubio, L., Galipienso, L. and Ferriol, I., 2020. Detection of plant viruses and disease management: Relevance of genetic diversity and evolution. *Frontiers in plant science*, **11**: 092.
- Saikanth, D R., Supriya & Singh, Bal & Rai, Avinash & Bana, Sita & Singh Sachan, Dhruvendra & Singh, Barinderjit. 2023. Advancing Sustainable Agriculture: A Comprehensive Review for Optimizing Food Production and Environmental Conservation. *International Journal of Plant & Soil Science*. **35**: 417-425. 10.9734/IJPSS/2023/v35i163169.
- Sapbamrer, R., Kitro, A., Panumasvivat, J., & Assavanopakun, P. (2023). Important role of the government in reducing pesticide use and risk sustainably in Thailand: Current situation and recommendations. *Frontiers in public health*, **11**: 1141142.
- Sarhan, E.A., Abd-Elsyed, M.H. and Ebrahiem, A.M., 2020. Biological control of cucumber powdery mildew (*Podosphaeraxanthii*) (Castagne) under greenhouse conditions. *Egyptian Journal of Biological Pest Control*, **30**: 1-7.
- Shaltiel-Harpaz L, Gerling D, Graph S, Kedoshim H, Azolay L, Rozenberg T (2016) Control of the tomato leafminer, *Tuta absoluta* (Lepidoptera: Gelechiidae), in open-field tomatoes by indigenous natural enemies occurring in Israel. *J Econ Entomol* **109**: 120-131.
- Sharma and Shikha. 2023. Precision Agriculture: Reviewing the Advancements, Technologies, and Applications in Precision Agriculture for Improved Crop Productivity and Resource Management. **4**: 41-45. 10.26480/rfna.02.2023.41.45.
- Skendžić, S., Zovko, M., Živković, I. P., Lešić, V., Lemić, D. 2021. The Impact of Climate Change on Agricultural Insect Pests. *Insects*, **12**(5): 440.

- Udayanga, D., 2019. The promise of molecular identification of fungi to overcome the global challenges in plant biosecurity. University of Sri Jayewardenepura. *Technology***29**: 103-113
- United States Department of Agriculture. *U.S. Horticulture Operations Report \$13.8 Billion in Sales*, [www.nass.usda.gov/Newsroom/archive/2020/12-08-2020.php](http://www.nass.usda.gov/Newsroom/archive/2020/12-08-2020.php). Accessed 2 July 2024.
- Whiteflies in Greenhouse Crops: Biology, Damage and Management. *Ontario.Ca*, [www.ontario.ca/page/whiteflies-greenhouse-crops-biology-damage-and-management](http://www.ontario.ca/page/whiteflies-greenhouse-crops-biology-damage-and-management). Accessed 28 June 2024.
- Yanar, D., Gebologlu, N., Cakar, T.U.Ğ.B.A. and Engur, M., 2019. The use of predatory mite *Phytoseiulus persimilis* (Acari: Phytoseiidae) in the control of two-spotted spider mite (*Tetranychusurticae* Koch, Acari: Tetranychidae) at greenhouse cucumber production in Tokat province, Turkey. *Applied Ecology and Environmental Research*, **17**(2): 2033-2041