

## **Effects of KNO<sub>3</sub> Seed Priming and Fertilizer Management on Growth and Yield of Green Gram (*Vigna radiata* L.) in a Temperate Agro-climate.**

### **Abstract**

Green gram (*Vigna radiata* L) is one of the most important nutritive pulse crop . It is an excellent source of protein with high quality of lysine. Seed priming techniques are used to improve germination, to reduce the time between sowing and emergence and for uniform emergence in the main field, especially under adverse environment. Keeping in view the nutritive importance of green gram, an investigation was carried out to assess the effects of seed priming and nutrient management for higher yield in green gram at research farm of Division of Agronomy, Faculty of Agriculture Wadura SKUAST-K Sopore during kharief-2021&2022 and the experiment was laid out in randomized complete block design (RCBD) with twelve treatments viz T<sub>0</sub>-Control, T<sub>1</sub>- Priming with 1% KNO<sub>3</sub>, T<sub>2</sub>-RFD, T<sub>3</sub>-RFD+ Priming with 1% KNO<sub>3</sub>, T<sub>4</sub>-1/2RFD, T<sub>5</sub>-1/2RFD+ Priming with 1% KNO<sub>3</sub>, T<sub>6</sub>-1/4RFD, T<sub>7</sub>-1/4RFD+ Priming with 1% KNO<sub>3</sub>, T<sub>8</sub>-1/2RFD+VAM, T<sub>9</sub>-1/2RFD+VAM+ Priming with 1% KNO<sub>3</sub>, T<sub>10</sub>-1/4RFD+VAM and T<sub>11</sub>-1/4RFD+VAM+ Priming with 1% KNO<sub>3</sub>. Among the treatments studied in both years, highest number of pod cluster per plant (9.66 and 9.33), number of pods per cluster (18.33 and 19.00), number of seeds per pod (13.00 and 13.33), 1000 -seed weight (43.66& 44.50 g) and yield per hectare (12.83 &13.30 q) respectively were noticed with full recommended dose of fertilizer along with seed priming by 1% KNO<sub>3</sub>.

**Key Words: Green gram, priming, potassium nitrate, growth, yield,**

### **Introduction**

Pulses are wonder gift of nature to the living universe and are the real gateway of sustainable agriculture(R.K.Bhairawa *et al.*,2014).Green gram (*Vigna radiata* L) is one of the most important pulse crop of the global economic importance; also called moong or mung bean and belongs to the family Leguminosae. It is an excellent source of protein (24.5%) with high quality of lysine 460mg/g, tryptophan 60mg/g and fair amount of fat0.6%, fiber 0.9% and ash 3.7% (Dharwe *et al.*, 2018).One possible way to mitigate the nutrients requirement of plant by the efficient utilization of biofertilizers as an eco-friendly technologies and based on renewable energy sources has gained momentum in recent years to supplement the parts of chemical fertilizers (Meena *et al.*, 2017). In leguminous crops the *Rhizobium* plays a significant role in maintaining and improving the soil

fertility and sustainability through its ability to fix atmospheric nitrogen in the soil through root nodules (Patel *et al.*, 2016). An efficient use of microbes increased yield reduces the soil borne pathogens and improves soil health (Haque *et al.*, 2012; Kumar *et al.*, 2017). Seed priming improves germination by reducing the time between sowing and emergence, especially under adverse environmental conditions (Gupta *et al.*, 2008). Seed priming is much beneficial in legumes as it substantially increases the yield of mung bean (Rashid *et al.*, 2004). Potassium nitrate ( $\text{KNO}_3$ ) is a chemical which can impact on growth, amount of nitrate, its reduction and assimilation in plants. Potassium deficiency reduces the photosynthetic  $\text{CO}_2$  fixation and transport and utilization of assimilates. Membrane and chlorophyll degradation are common in potassium deficient plants.  $\text{KNO}_3$  regulates water uptake and maintain seed moisture level which is crucial for initiation of several metabolic processes to break the seed dormancy even in drought or elevated temperatures (Javed *et al.*, 2020, Ali *et al.*, 2021). Foliar application of potassium can also be considered as one of the alternative in order to reduce such problems. Foliar application of nutrients may result in economic use of fertilizer by virtue of reducing loss through different processes and supplying nutrients instantly to the crop. Under the moisture stress conditions application of potassium may be considered as one promising option as it influences the water economy and crop growth, through its effect on water uptake, root growth, maintenance of turgor, transpiration and stomata behavior (Hsiao and Lauchi, 1986). Keeping in view the above facts, an investigation has been carried out to assess the effects of seed priming and nutrient management for higher yield in green gram under temperate agro climatic conditions.

## **Material and Methods**

The experiment was laid out at Experimental Farm of Division of Agronomy, Faculty of Agriculture Wadura SKUAST-K Sopore during kharief-2021&2022 and the experiment was laid out in randomized complete block design (RCBD) with twelve treatments viz T<sub>0</sub>-Control, T<sub>1</sub>- Priming with 1%  $\text{KNO}_3$ , T<sub>2</sub>-RFD, T<sub>3</sub>-RFD+ Priming with 1%  $\text{KNO}_3$ , T<sub>4</sub>-1/2RFD, T<sub>5</sub>-1/2RFD+ Priming with 1%  $\text{KNO}_3$ , T<sub>6</sub>-1/4RFD, T<sub>7</sub>-1/4RFD+ Priming with 1%  $\text{KNO}_3$ , T<sub>8</sub>-1/2RFD+VAM, T<sub>9</sub>-1/2RFD+VAM+ Priming with 1%  $\text{KNO}_3$ , T<sub>10</sub>-1/4RFD+VAM and T<sub>11</sub>-1/4RFD+VAM+ Priming with 1%  $\text{KNO}_3$ . The soil has pH 7.0, E.C 0.85 dsm<sup>-1</sup>, OC 0.51%, avg. N 340 kg/ha, avg. P 35 kg/ha, avg. K 230 kg/ha. The field was cleared manually, ploughed and weeded before seedbeds were marked and demarked. The seeds of green gram var. Shalimar mung-1 were sown in plots of 3.0 × 1.5 m size in the 1<sup>st</sup> week of June. The row to row spacing of 25 cm and plant to plant spacing

of 10 cm has been adapted. 15 days after germination, the thinning activities have been done. The sources of different organic manures tried in experiment were applied at the time of land preparation. Agronomic cultural practices recommended by SKUAST-K were adopted to raise the crop. The seeds were soaked in 1%  $\text{KNO}_3$  solution for 8 hours and dried under shade before sowing. Observations on various growth and yield related attributes were recorded, using standard procedures. The data thus collected was subjected to analysis of variance, using the method proposed by Panse and Sukhatme (1985).

## **RESULTS AND DISCUSSION**

### **Growth and Yield Parameters**

It is evident from the data of two years presented in Table 1, that there was significant difference among treatments tried in experiment related to growth and yield related attributes of green gram. Among different treatment combinations,  $T_3$ -Full recommended dose of fertilizer along with seed priming by 1% $\text{KNO}_3$  registered maximum number of pod cluster per plant (9.66 and 9.33), number of pods per cluster (18.33 and 19.00), number of seeds per pod (13.00 and 13.33), 1000 seed weight (43.66 & 44.50 g) and yield per hectare (12.83 & 13.30 q) respectively in both years followed by  $T_2$  and  $T_9$  and  $T_1$  was found statistically significant as compared to rest of the treatments. Seed priming by potassium nitrate in mung bean might be due to the better tolerance towards moisture stress and in turn might have boosted plant growth effectively. The nutrient status in the plant plays a vital role in improving the resistance of plant to stress conditions (Yadov, 2006). Similar findings have been reported by Bardhan *et al.*, (2007). Increase in number of branches by potassium nitrate application might be attributed to the fact that potassium enhances plant vigor and strengthens the stalk which might have resulted in better plant growth and more number of branches. Similar findings have been reported by Deotale *et al.*, (2015). The increase in number of seeds per pod might be due to the application of  $\text{KNO}_3$  which have active biochemical functions in plants, enzyme activation, photosynthesis and cell division that increased the number of seeds per pod. Similar findings have been reported by Golezani *et al.*, (2011).  $\text{KNO}_3$  is effective in improving the yield parameters under moisture stress conditions by delaying maturity due to enhancement of biochemical and physiological processes (Rahman and Attia, 2016). Potassium nitrate regulates the hormonal production such as auxins, cytokinins, and gibberellins positively while lower the production of abscisic acid in plants under high temperature stress (Bolat *et al.*, 2024). Use of potassium nitrate in various agricultural products including

fertilizers is very common as it improves the soil fertility and crop growth and yield by boosting the accumulation of nutrients like nitrogen, potassium, free amino acids, and proteins( Chalmardi *et al.*, 2014).

## **Conclusion**

Application of recommended dose of fertilizers along with seed priming by 1%KNO<sub>3</sub> had a significant response on growth and yield of green gram. The results obtained revealed that green gram responded well to the full recommended dose of fertilizers along with seed priming by 1%KNO<sub>3</sub> to other treatments in the study. Based on the findings of this study, it may be recommended that full recommended dose of fertilizers along with seed priming by 1%KNO<sub>3</sub> is adequate for maximum growth and yield of green gram in the study location.

## **References**

- Abd-El-Rhman, I.E. and Attia, M.F. (2016) Foliar spray with potassium nitrate and salicylic acid for improving growth, yield and nutrients uptake by olive trees under salinity stress conditions. *International Journal of Chem. Tech. Research.* 9(12): 230-245
- Ali LG, Nulit R, Ibrahim MH. Efficacy of KNO<sub>3</sub>, SiO<sub>2</sub> and SA priming for improving emergence, seedling growth and antioxidant enzymes of rice (*Oryza sativa*), under drought. *Sci Rep.* 2021;11:3864. <https://doi.org/10.1038/s41598-021-83434-3>
- Bairwa, R. K., Nepalia, V., Balai, C. M., Jalwania, R., &Meena, H. P. (2014). Yield and nutrient uptake of summer greengram (*Vigna radiata* L.) under different levels of phosphorus and sulphur fertilizations. *SAARC Journal of Agriculture*, 12(1): 162-172.
- Bardhan, K., Kumar, V. and Dhimmar, S.K (2007). An evaluation of the potentiality of exogenous osmoprotectants mitigating water stress on chickpea. *The Journal of Agricultural Sciences.* 3(2), 67–74. DOI: <http://doi.org/10.4038/jas.v3i2.8101>
- Bolat I, Korkmaz K, Dogan M, Turan M, Kaya C, Seyed Hajizadeh H, Kaya O. Enhancing drought, heat shock, and combined stress tolerance in Myrobalan 29 C rootstocks with foliar application of potassium nitrate. *BMC Plant Biol.* 2024;24(1):140.
- Chalmardi ZK, Abdolzadeh A, Sadeghipour HR. Silicon nutrition potentiates the antioxidant metabolism of rice plants under iron toxicity. *Acta Physiol Plant.* 2014;36:493–502

- Deotale, R.D., Mahale, S.A., Patil, S.R., Sahane, A.N. and Sawant, P.P. (2015) Effect of foliar sprays of nitrate salts on morphophysiological traits and yield of green gram. *International Journal of Researches in Biosciences, Agriculture and Technology*. 3(II) 63-67
- Dharwe, D. S., Dixit, H. S., Dotaniya, C. K., Mohbe, S., &Doutaniya, R. K. (2018). Influences of sulphur on the yield of summer greengram (*Vigna radiata* L.).*Journal of Pharmacognosy and Phytochemistry*, 7(2):147-149.
- Golezani, K.G., Jafari, S.F. and Kolvanagh, J.S. (2011). Seed priming and field performance of soybean (*Glycine max* L.) in response to water limitation..*Notulae Botanicae Horti Agrobotanici*. 39(2):186-189.
- Hsiao, T.C. and Lauchli, A. (1986) Role of potassium in plant water relations. In-Tinker, B. and Lauchli, A. (eds.). *Advances in Plant Nutrition*.2, 281-312.
- Gupta, A., Davlene, M., Arunkumar, M.B., Roy, M., Naseem, M., Choudhary, V.K. and Maib, R.K., Seed priming after math. *Int. J. Agric. Environ. Biotechnol.*, **1**: 199-209 (2008).
- Haque, M., Ilias, G.N.M., Molla, A.H. (2012) Impact of Trichoderma-enriched bio-fertilizer on the growth and yield of mustard (*Brassica rapa* L.) and tomato (*Solanum lycopersicon* mill.). *Agriculturists* **10**(2):109-119.
- Javed T, Ali MM, Shabbir R, Gull S, Ali A, Khalid E, Abbas AN, Tariq M. Rice seedling establishment as influenced by cultivars and seed priming with potassium nitrate. *J Appl Res Plant Sci*. 2020;1:65–75.
- Kumar, J., Chaurasia, A. K., & Bara, B. M. (2017). Effect of organic priming on germination and vigor of cotton (*Gossypium hirsutum* L.) seed. *Journal of Pharmacognosy and Photochemistry*, 6(3), 815-819.
- Meena, V.S., Meena, S.K., Verma, J.P., Kumar, A., Aeron, A., Mishra, P.K., Bisht, J.K., Pattanayak, A., Naveed, M., Dotaniya, M.L.(2017). Plant beneficial rhizospheric microorganism (PBRM) strategies to improve nutrients use efficiency: A review. *Ecological Engineering*, DOI: 10.1016/j.ecoleng.2017.06.058.
- Panse VG, Sukhatme PV. *Statistical Methods for Agricultural Workers*, ICAR, New Delhi, 1985.
- Patel, A.R., Patel, D.D., Patel, T.U. and Patel, H.M. (2016). Nutrient management in summer green gram (*Vigna radiata*). *International Journal of Applied and Pure Science and Agriculture* **2**(2):133-142.

Rashid, A., D. Harris, P. Hollington and S. Ali. 2004. On-farm seed priming reduces yield losses of mung bean (*Vigna radiata*) associated with mung bean yellow mosaic virus in NWFP of Pakistan. *Crop Prot.* 23:1119-1124.

Yadov, D.V. (2006) Potassium nutrition of sugarcane. In: Benbi DK *et. al.* Balanced fertilization for sustaining crop productivity. International Potash Institute, Horgen Pp. 275– 288.

**Table 1: Effects of KNO<sub>3</sub> Seed Priming and Fertilizer Management on Growth and Yield of Green Gram (*Vigna radiata* L.) in a Temperate Agro-climate.**

Treatments	No. of Pod Cluster/Plant		No. of Pods/Cluster		NO.OF Seeds/Pod		1000-Seed Weight(g)		Yield/ha(q)	
	2021	2022	2021	2022	2021	2022	2021	2022	2021	2022
T <sub>0</sub> –Control	2.33	2.66	9.00	8.66	5.00	5.33	34.16	33.83	6.30	6.06
T <sub>1</sub> – Priming with 1%KNO <sub>3</sub>	3.00	3.33	10.33	10.66	6.00	6.00	35.83	36.50	7.23	7.46
T <sub>2</sub> –(RFD)	8.66	8.33	17.00	17.33	12.33	12.33	42.33	43.33	11.90	12.13
T <sub>3</sub> – RFD+ Priming with 1%KNO <sub>3</sub>	9.66	9.33	18.33	19.00	13.00	13.33	43.66	44.50	12.83	13.30
T <sub>4</sub> – 1/2RFD	6.33	6.00	14.00	14.33	9.00	9.33	38.50	39.00	9.80	10.03
T <sub>5</sub> –1/2RFD + Priming with 1%KNO <sub>3</sub>	7.00	7.66	14.66	15.33	10.33	10.00	40.00	40.50	10.26	10.73
T <sub>6</sub> –1/4RFD	3.66	3.66	11.66	12.33	6.66	7.00	36.50	37.00	8.16	8.63
T <sub>7</sub> –1/4RFD + Priming with 1%KNO <sub>3</sub>	4.33	4.00	12.33	12.66	7.00	7.00	37.16	38.16	8.63	8.86
T <sub>8</sub> –1/2RFD + VAM	7.66	7.00	15.66	15.33	10.33	10.66	41.66	42.16	10.96	10.73
T <sub>9</sub> – 1/2RFD +VAM+ Priming with 1%KNO <sub>3</sub>	8.33	8.00	16.33	16.66	11.00	11.66	41.83	42.16	11.43	11.66
T <sub>10</sub> – 1/4RFD + VAM	5.00	5.00	11.33	11.66	8.33	7.66	37.50	37.50	7.93	8.16
T <sub>11</sub> – 1/4RFD +VAM+ Priming with 1%KNO <sub>3</sub>	5.66	5.33	12.66	13.00	8.66	8.00	38.00	38.50	8.86	9.10
C.D(P≤0.05)	0.78	0.86	1.23	1.41	1.25	1.35	0.91	1.42	0.86	0.99