

## Original Research Article

# Efficacy of biopesticides against sucking pests infesting moth bean [*Vigna aconitifolia* (Jacq.) Marechal]

---

### ABSTRACT

A field experiment was conducted at College of Agriculture, NAU, Bharuch (Gujarat) to study the efficacy of biopesticides against sucking pests of moth bean during *kharif* -2023. Results revealed that among the different biopesticides tested against sucking pests of moth bean, *Beauveria bassiana* ( $1 \times 10^8$  cfu g<sup>-1</sup>), neem oil 0.5% and NSKE 5% was found most effective whereas, treatments of *Lecanicillium lecanii* ( $1 \times 10^8$  cfu g<sup>-1</sup>), *Metarhizium anisopliae* ( $1 \times 10^8$  cfu g<sup>-1</sup>) and azadirachtin 0.15 EC found moderately effective against jassid, *Empoasca motti* (Pruthi.) and whitefly, *Bemisia tabaci* (Genn.) populations. Treatments of neem oil 0.5% and azadirachtin 0.15 EC was found most effective whereas, *Beauveria bassiana* ( $1 \times 10^8$  cfu g<sup>-1</sup>) and NSKE 5% found moderately effective against thrips, *Caliothrips indicus* (Bagnall). Treatment of novel plus 0.1% was found less effective against jassid, white fly and thrips.

*Keywords:* Moth bean, sucking pests, jassid, white fly, thrips, *Beauveria bassiana*

### 1. INTRODUCTION

*Vigna aconitifolia* (Jacq.) Marechal, also referred to as "moth," is a vital pulse crop that thrives in the country's dry and semi-arid regions. It is the *kharif* pulse that can withstand drought the best. In addition to covering a wide surface area, plants also retain moisture and prevent soil erosion. A member of the Papilionaceae subfamily of the Leguminosae family is the moth bean. An annual plant is the moth bean. Because of its deeper tap roots, it may draw moisture from the soil's lower strata. The plant height is between 30 and 35 cm, and the stem is branching. The trifoliolate leaves have lobed leaflets that are separated into three to five sections. Flowers are papilionaceous and mostly self-pollinated (Kukvaya et al., 2018). In India, moth bean occupies an area of 9.68 lakh ha with production of 3.21 lakh tonnes and productivity of 332 kg/ha whereas in Gujarat, it occupies an area of 0.12 lakh ha with production of 0.05 lakh tonnes and productivity of 462 kg/ha during year 2019 – 2020 (Anon., 2021). Jassids and whiteflies also act as vector of yellow mosaic virus apart from causing direct damage by desapping (Satyavir et al., 1984). Termites, galerucid beetles, mites and surface grass hoppers are minor pests, while jassid, whitefly, thrips, black weevil, pulse beetle and white grubs are major pests of moth bean (Bhathesar et al., 2021). Sucking pests

cause considerable damage on moth bean and management through chemical pesticides kill the natural enemies and cause pest resurgence [8-10]. Hence in this study biopesticides were tested for their efficacy against sucking pests of moth bean.

## 2. MATERIAL AND METHODS

A field experiment was conducted at College of Agriculture, NAU, Bharuch (Gujarat) to study the efficacy of biopesticides against sucking pests of moth bean during *kharif* -2023. For this, moth bean var. GMO-2 was selected and sown at a distance of 45 cm x 10 cm in Randomized Block Design with eight treatments and three replications having gross plot of Gross: 2.7 m x 4.0 m and net plot of 1.8 m x 3.8 m. The moth bean variety GMO-2 was raised by adopting all recommended agronomical practices. Two spray of biopesticides were given, first spray at time of appearance of pest *i.e.*, at 01/09/2023 and second spray was given at 10 days after first spray with manually operated knapsack sprayer. An untreated check was also maintained for comparison. The observations were recorded one day prior to first spray as well as 3<sup>rd</sup>, 7<sup>th</sup>, and 9<sup>th</sup> days after each spray. Standard methodology was adopted for recording pest population. Five plants were randomly selected from each plot and observations were recorded before and after each spray. The population of adults and nymphs of jassid [*Empoasca motti* (Pruthi.)], whitefly [*Bemisia tabaci*(Genn.)] and thrips [*Caliothrips indicus*(Bagnall)] were recorded from three leaves (upper, middle and lower) of each randomly selected plants and mean pest population was worked out.

## 3. RESULTS AND DISCUSSION

### 1. Jassid, *Empoasca motti* (Pruthi.)

#### First spray

Pooled over periods result of first spray (Table-1) revealed that the lower jassid population was observed in the plot treated with *Beauveria bassiana*( $1 \times 10^8$  cfu g<sup>-1</sup>) (2.70 per 3 leaves) and it was at par with neem oil 0.5% (2.77 per 3 leaves) and NSKE 5% (2.83 per 3 leaves). The next effective treatment was *Lecanicillium lecanii*( $1 \times 10^8$ cfu g<sup>-1</sup>) (4.13 per 3 leaves), it was at par with *Metarhizium anisopliae*( $1 \times 10^8$ cfu g<sup>-1</sup>) (4.20 per 3 leaves), azadirachtin 0.15 EC (4.26 per 3 leaves) and novel plus 0.1% (4.53 per 3 leaves) were found less effective. The highest jassid population was recorded in control (6.44 per 3 leaves).

#### Second Spray:

Pooled over periods result of second spray (Table-2) revealed that the lowest jassid population was observed in the plot treated with *Beauveria bassiana*( $1 \times 10^8$  cfu g<sup>-1</sup>) (1.67 per 3 leaves) and it was at par with neem oil 0.5% (1.73 per 3 leaves) and NSKE 5% (1.80 per 3 leaves). The next effective treatment was *Lecanicillium lecanii*( $1 \times 10^8$ cfu g<sup>-1</sup>) (2.76 per 3 leaves) and it was at par with *Metarhizium anisopliae*( $1 \times 10^8$ cfu g<sup>-1</sup>) (2.80 per 3 leaves) and azadirachtin 0.15 EC (2.88 per 3 leaves). The treatment of novel plus 0.1% (3.46 per 3 leaves) was found less effective. The highest jassid population was recorded in control (5.77 per 3 leaves).

#### Pooled over two sprays

The pooled over two spray (Table-2)revealed that all the biopesticides were found significantly superior over control. Among different treatments, significantly lowest population of jassid was recorded in plots treated with *Beauveria bassiana*( $1 \times 10^8$  cfu g<sup>-1</sup>) (2.18 per 3 leaves) and it was at par with neem oil 0.5% (2.25 per 3 leaves) and NSKE 5% (2.32 per 3 leaves). The next effective treatment was *Lecanicillium lecanii*( $1 \times 10^8$ cfu g<sup>-1</sup>) (3.44 per 3 leaves) and it was at par with *Metarhizium anisopliae*( $1 \times 10^8$ cfu g<sup>-1</sup>) (3.50 per 3 leaves) and

azadirachtin 0.15 EC (3.57 per 3 leaves). The treatment of novel plus 0.1% (3.99 per 3 leaves) was found less effective. The highest jassid population was recorded in control (6.11 per 3 leaves).

Sujatha and Bharpoda (2017) observed that among different treatments lowest jassid population was observed in *Beauveria bassiana* in green gram. Thus, present findings are in accordance with the earlier findings.

Sr. no.	Treatments	Mean no. of Jassid/3 leaves				
		Before spray	Days after spray			Pooled over periods
			3	7	9	
<b>T1</b>	<i>Beauveria bassiana</i> 1.15 WP(1 x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.51 (5.80)	1.90 (3.12)	1.66 (2.25)	1.79 (2.72)	1.78 (2.70)
<b>T2</b>	<i>Metarhizium anisopliae</i> 1.15 WP(1 x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.73 (6.95)	2.27 (4.64)	2.04 (3.68)	2.19 (4.28)	2.16 (4.20)
<b>T3</b>	<i>Lecanicillium lecanii</i> 2% AS(1x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.63 (6.42)	2.25 (4.55)	2.02 (3.60)	2.18 (4.24)	2.15 (4.13)
<b>T4</b>	Azadirachtin 0.15 EC (1500 ppm)	2.80 (7.36)	2.26 (4.62)	2.06 (3.74)	2.22 (4.42)	2.18 (4.26)
<b>T5</b>	Neem oil @ 0.5%	2.67 (6.65)	1.92 (3.20)	1.67 (2.30)	1.82 (2.80)	1.80 (2.77)
<b>T6</b>	Novel plus @ 0.1%	2.65 (6.53)	2.28 (4.70)	2.19 (4.32)	2.25 (4.56)	2.24 (4.53)
<b>T7</b>	NSKE @ 5%	2.62 (6.35)	1.94 (3.25)	1.69 (2.36)	1.84 (2.88)	1.82 (2.83)
<b>T8</b>	Control (water spray)	2.54 (5.94)	2.58 (6.15)	2.58 (6.17)	2.74 (7.01)	2.63 (6.44)
	<b>S.Em.±</b>	0.12	0.09	0.09	0.10	0.05
	<b>S.Em.± (P×T)</b>	-	-	-	-	0.09
	<b>CD at 5 %</b>	NS	0.29	0.28	0.29	0.15
	<b>CD at 5 % (P×T)</b>	-	-	-	-	NS
	<b>CV %</b>	7.38	7.67	7.90	7.84	7.82

Figure in parentheses are retransformed value whereas, those outside are  $\sqrt{x} + 0.5$  transformed

Sr. no.	Treatments	Mean no. of Jassid/3 leaves				
		Days after spray (DAS)			Pooled over periods	Pooled Over two sprays
		3	7	9		
T1	<i>Beauveria bassiana</i> 1.15 WP(1 x 10 <sup>8</sup> cfu g <sup>-1</sup> )	1.78 (2.68)	1.32 (1.24)	1.26 (1.08)	1.45 (1.67)	1.62 (2.18)
T2	<i>Metarhizium anisopliae</i> 1.15 WP(1 x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.08 (3.84)	1.77 (2.64)	1.56 (1.92)	1.80 (2.80)	1.99 (3.50)
T3	<i>Lecanicillium lecanii</i> 2% AS(1x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.07 (3.80)	1.76 (2.60)	1.54 (1.88)	1.79 (2.76)	1.97 (3.44)
T4	Azadirachtin 0.15 EC (1500 ppm)	2.09 (3.88)	1.79 (2.72)	1.60 (2.05)	1.83 (2.88)	2.00 (3.57)
T5	Neem oil @ 0.5%	1.80 (2.74)	1.36 (1.34)	1.27 (1.12)	1.48 (1.73)	1.64 (2.25)
T6	Novel plus @ 0.1%	2.24 (4.50)	2.02 (3.58)	1.67 (2.30)	1.98 (3.46)	2.11 (3.99)
T7	NSKE @ 5%	1.81 (2.79)	1.39 (1.44)	1.30 (1.18)	1.50 (1.80)	1.66 (2.32)
T8	Control (water spray)	2.71 (6.87)	2.66 (6.58)	2.09 (3.86)	2.49 (5.77)	2.56 (6.11)
	<b>S.Em.±</b>	0.09	0.08	0.10	0.05	0.05
	<b>S.Em.± (P×T)</b>	-	-	-	0.09	0.08
	<b>CD at 5 %</b>	0.28	0.26	0.29	0.15	0.14
	<b>CD at 5 % (P×T)</b>	-	-	-	NS	NS
	<b>CV %</b>	7.72	8.29	10.93	8.83	7.78

Figure in parentheses are retransformed value whereas, those outside are  $\sqrt{x} + 0.5$  transformed

## 2. White fly, *Bemisia tabaci*(Genn.)

### First spray

Pooled over periods result of first spray (Table-3) revealed that the lowest whitefly population was observed in the plot treated with *Beauveria bassiana*(1 x 10<sup>8</sup> cfu g<sup>-1</sup>) (2.76 per 3 leaves) and it was at par with neem oil 0.5% (2.90 per 3 leaves) and NSKE 5% (2.99 per 3 leaves). The next effective treatment was *Lecanicillium lecanii*(1x 10<sup>8</sup>cfu g<sup>-1</sup>) (4.10 per 3 leaves), it was at par with *Metarhizium anisopliae*(1 x 10<sup>8</sup>cfu g<sup>-1</sup>) (4.16 per 3 leaves ) and azadirachtin 0.15 EC (4.28 per 3 leaves). The treatment of novel plus 0.1% (4.87 per 3 leaves) was found less effective. The highest whitefly population was recorded in control (6.56 per 3 leaves).

### Second spray

Pooled over periods result of second spray (Table-4) revealed that the lowest whitefly population was observed in the plot treated with *Beauveria bassiana*(1 x 10<sup>8</sup> cfu g<sup>-1</sup>) (1.72 per 3 leaves) and it was at par with neem oil 0.5% (1.81 per 3 leaves) and NSKE 5% (1.91 per 3 leaves). The next effective treatment was *Lecanicillium lecanii*(1x 10<sup>8</sup>cfu g<sup>-1</sup>) (2.73 per

3 leaves) and it was at par with *Metarhizium anisopliae*( $1 \times 10^8$ cfu g<sup>-1</sup>) (2.86 per 3 leaves )and azadirachtin 0.15 EC (2.97 per 3 leaves). The treatment of novel plus 0.1% (3.65 per 3 leaves) was found less effective. The highest whitefly population was recorded in control (5.07 per 3 leaves).

Sr. no.	Treatments	Mean no. of Whitefly/3 leaves				
		Before Spray	Days after spray (DAS)			Pooled over periods
			3	7	9	
T1	<i>Beauveria bassiana</i> 1.15 WP( $1 \times 10^8$ cfu g <sup>-1</sup> )	2.57 (6.10)	1.90 (3.12)	1.66 (2.25)	1.85 (2.92)	1.80 (2.76)
T2	<i>Metarhizium anisopliae</i> 1.15 WP( $1 \times 10^8$ cfu g <sup>-1</sup> )	2.59 (6.22)	2.30 (4.80)	2.03 (3.62)	2.14 (4.07)	2.16 (4.16)
T3	<i>Lecanicillium lecanii</i> 2% AS( $1 \times 10^8$ cfu g <sup>-1</sup> )	2.61 (6.32)	2.28 (4.69)	2.02 (3.58)	2.13 (4.02)	2.14 (4.10)
T4	Azadirachtin 0.15 EC (1500 ppm)	2.56 (6.06)	2.33 (4.92)	2.05 (3.72)	2.17 (4.20)	2.18 (4.28)
T5	Neem oil @ 0.5%	2.76 (7.12)	1.97 (3.40)	1.68 (2.31)	1.87 (2.98)	1.84 (2.90)
T6	Novel plus @ 0.1%	2.61 (6.32)	2.37 (5.14)	2.22 (4.42)	2.36 (5.05)	2.32 (4.87)
T7	NSKE @ 5%	2.63 (6.42)	1.99 (3.48)	1.73 (2.48)	1.87 (3.00)	1.86 (2.99)
T8	Control (water spray)	2.54 (5.98)	2.65 (6.51)	2.66 (6.57)	2.66 (6.59)	2.66 (6.56)
	<b>S.Em.±</b>	0.12	0.09	0.08	0.09	0.05
	<b>S.Em.± (P×T)</b>	-	-	-	-	0.08
	<b>CD at 5 %</b>	NS	0.27	0.26	0.26	0.14
	<b>CD at 5 % (P×T)</b>	-	-	-	-	NS
	<b>CV %</b>	8.06	6.84	7.36	7.02	7.06

Figure in parentheses are retransformed value whereas, those outside are  $\sqrt{x} + 0.5$  transformed values

#### **Pooled over two sprays**

The pooled over two spray (Table-4)revealed that all the biopesticides were found significantly superior over control. Among different treatments, significantly lowest population of whitefly was recorded in plots treated with *Beauveria bassiana*( $1 \times 10^8$ cfu g<sup>-1</sup>) (2.24 per 3 leaves) and it was at par with neem oil 0.5% (2.35 per 3 leaves) and NSKE 5% (2.45 per 3 leaves). The next effective treatment was *Lecanicillium lecanii*( $1 \times 10^8$ cfu g<sup>-1</sup>) (3.41 per 3 leaves) and it was at par with *Metarhizium anisopliae*( $1 \times 10^8$ cfu g<sup>-1</sup>) (3.51 per 3 leaves) and azadirachtin 0.15 EC (3.63 per 3 leaves). The treatment of novel plus 0.1% (4.26 per 3 leaves) was found less effective. The highest whitefly population was recorded in control (5.81 per 3 leaves).

Singh et al. (2018) found that among different treatments *Beauveria bassiana* was highly effective against whitefly in green gram. thus, present findings are in confirmation with earlier findings.

Sr. no.	Treatments	Mean no. of Whitefly/3 leaves				
		Days after spray (DAS)			Pooled over periods	Pooled Over two sprays
		3	7	9		
<b>T1</b>	<i>Beauveria bassiana</i> 1.15 WP(1 x 10 <sup>8</sup> cfu g <sup>-1</sup> )	1.81 (2.77)	1.35 (1.32)	1.26 (1.08)	1.47 (1.72)	1.64 (2.24)
<b>T2</b>	<i>Metarhizium anisopliae</i> 1.15 WP(1 x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.10 (3.92)	1.79 (2.72)	1.56 (1.94)	1.82 (2.86)	1.99 (3.51)
<b>T3</b>	<i>Lecanicillium lecanii</i> 2% AS(1x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.09 (3.85)	1.72 (2.45)	1.54 (1.88)	1.78 (2.73)	1.96 (3.41)
<b>T4</b>	Azadirachtin 0.15 EC (1500 ppm)	2.13 (4.02)	1.82 (2.82)	1.61 (2.08)	1.85 (2.97)	2.02 (3.63)
<b>T5</b>	Neem oil @ 0.5%	1.83 (2.85)	1.41 (1.48)	1.26 (1.10)	1.50 (1.81)	1.67 (2.35)
<b>T6</b>	Novel plus @ 0.1%	2.24 (4.50)	2.06 (3.73)	1.80 (2.73)	2.03 (3.65)	2.17 (4.26)
<b>T7</b>	NSKE @ 5%	1.85 (2.92)	1.46 (1.62)	1.30 (1.20)	1.54 (1.91)	1.70 (2.45)
<b>T8</b>	Control (water spray)	2.64 (6.48)	2.38 (5.14)	2.02 (3.59)	2.35 (5.07)	2.50 (5.81)
	<b>S.Em.±</b>	0.09	0.08	0.07	0.04	0.04
	<b>S.Em.± (P×T)</b>	-	-	-	0.08	0.08
	<b>CD at 5 %</b>	0.27	0.25	0.21	0.13	0.13
	<b>CD at 5 % (P×T)</b>	-	-	-	NS	NS
	<b>CV %</b>	7.40	8.03	7.76	7.75	7.21

Figure in parentheses are retransformed value whereas, those outside are  $\sqrt{x} + 0.5$  transformed values

### 3. Thrips, *Caliothrips indicus*(Bagnall)

#### First spray

Pooled over periods result of first spray (Table-5) revealed that the lowest thrips population was observed in the plot treated with neem oil 0.5% (3.05 per three leaves) and it was at par with azadirachtin 0.15 EC (3.25 per 3 leaves). The next effective treatment was *Beauveria bassiana*(1 x 10<sup>8</sup> cfu g<sup>-1</sup>) (3.88 per 3 leaves) and it was at par with NSKE 5% (3.94 per three leaves). The remaining treatments viz., *Metarhizium anisopliae*(1 x 10<sup>8</sup>cfu g<sup>-1</sup>) (4.71 per 3 leaves ), *Lecanicillium lecanii*(1x 10<sup>8</sup>cfu g<sup>-1</sup>) (4.87 per 3 leaves) and novel plus 0.1% (5.11 per 3 leaves) were found less effective and they were at par with each other.

#### Second Spray

Pooled over periods result of second spray (Table-6) revealed that lowest thrips population was observed in the plot treated with neem oil 0.5% (2.14 per three leaves) and it was at par with azadirachtin 0.15 EC (2.22 per 3 leaves). The next effective treatment was *Beauveria bassiana*(1 x 10<sup>8</sup> cfu g<sup>-1</sup>) (3.10 per 3 leaves) and it was at par with NSKE 5% (3.10 per three

leaves). The remaining treatments viz., *Metarhizium anisopliae*(1 x 10<sup>8</sup>cfu g<sup>-1</sup>) (3.83 per 3 leaves), *Lecanicillium lecanii*(1x 10<sup>8</sup>cfu g<sup>-1</sup>) (4.07 per 3 leaves) and novel plus 0.1% (4.17 per 3 leaves) were found less effective and they were at par with each other. The highest thrips population was recorded in control (5.26 per 3 leaves).

Sr. no.	Treatments	Mean no. of Thrips/3 leaves				
		Before Spray	Days after spray (DAS)			Pooled over periods
			3	7	9	
T1	<i>Beauveria bassiana</i> 1.15 WP(1 x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.90 (7.93)	2.14 (4.08)	1.97 (3.39)	2.16 (4.18)	2.09 (3.88)
T2	<i>Metarhizium anisopliae</i> 1.15 WP(1 x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.66 (6.60)	2.35 (5.03)	2.13 (4.02)	2.36 (5.07)	2.28 (4.71)
T3	<i>Lecanicillium lecanii</i> 2% AS(1x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.65 (6.52)	2.39 (5.20)	2.16 (4.18)	2.39 (5.22)	2.31 (4.87)
T4	Azadirachtin 0.15 EC (1500 ppm)	2.90 (7.92)	2.07 (3.80)	1.83 (2.84)	1.90 (3.12)	1.93 (3.25)
T5	Neem oil @ 0.5%	2.83 (7.49)	1.99 (3.48)	1.77 (2.62)	1.88 (3.05)	1.88 (3.05)
T6	Novel plus @ 0.1%	2.67 (6.61)	2.41 (5.29)	2.25 (4.56)	2.45 (5.48)	2.37 (5.11)
T7	NSKE @ 5%	2.90 (7.93)	2.16 (4.19)	1.98 (3.43)	2.17 (4.20)	2.11 (3.94)
T8	Control (water spray)	2.55 (6.01)	2.70 (6.78)	2.71 (6.83)	2.74 (7.00)	2.72 (6.87)
	<b>S.Em.±</b>	0.13	0.09	0.09	0.09	0.05
	<b>S.Em.± (P×T)</b>	-	-	-	-	0.09
	<b>CD at 5 %</b>	NS	0.28	0.29	0.28	0.15
	<b>CD at 5 % (P×T)</b>	-	-	-	-	NS
	<b>CV %</b>	8.15	7.02	7.83	7.05	7.29

Figure in parentheses are retransformed value whereas, those outside are  $\sqrt{x} + 0.5$  transformed values.

### Pooled over two sprays

The pooled over two spray (Table-6) revealed that all the biopesticides were found significantly superior over control. Among different treatments, significantly lowest population of thrips was recorded in plots treated with neem oil 0.5% (2.60 per three leaves) and it was at par with azadirachtin 0.15 EC (2.74 per 3 leaves). The next effective treatment was *Beauveria bassiana*(1 x 10<sup>8</sup> cfu g<sup>-1</sup>) (3.49 per 3 leaves) and it was at par with NSKE 5% (3.52 per three leaves). The remaining treatments viz., *Metarhizium anisopliae*(1 x 10<sup>8</sup>cfu g<sup>-1</sup>) (4.27 per 3 leaves ), *Lecanicillium lecanii*(1x 10<sup>8</sup>cfu g<sup>-1</sup>) (4.47 per 3 leaves) and novel plus 0.1% (4.64 per 3 leaves) were found less effective and they were at par with each other. The highest thrips population was recorded in control (6.02 per 3 leaves).

Chaudhary et al. (2018) found that among different treatments neem oil 0.15%was highly effective against thrips in soybean which is in agreement with present findings.

Sr. no.	Treatments	Mean no. of Thrips/3 leaves				
		Days after spray (DAS)			Pooled over periods	Pooled Over two sprays
		3	7	9		
T1	<i>Beauveria bassiana</i> 1.15 WP(1 x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.14 (4.10)	1.89 (3.08)	1.62 (2.12)	1.89 (3.10)	1.99 (3.49)
T2	<i>Metarhizium anisopliae</i> 1.15 WP(1 x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.35 (5.04)	2.12 (4.01)	1.72 (2.45)	2.07 (3.83)	2.17 (4.27)
T3	<i>Lecanicillium lecanii</i> 2% AS(1x 10 <sup>8</sup> cfu g <sup>-1</sup> )	2.37 (5.12)	2.14 (4.06)	1.88 (3.02)	2.13 (4.07)	2.22 (4.47)
T4	Azadirachtin 0.15 EC (1500 ppm)	1.89 (3.08)	1.70 (2.39)	1.30 (1.18)	1.63 (2.22)	1.78 (2.74)
T5	Neem oil @ 0.5%	1.87 (2.99)	1.68 (2.32)	1.27 (1.12)	1.61 (2.14)	1.74 (2.60)
T6	Novel plus @ 0.1%	2.40 (5.26)	2.15 (4.12)	1.90 (3.12)	2.15 (4.17)	2.26 (4.64)
T7	NSKE @ 5%	2.15 (4.12)	1.89 (3.09)	1.61 (2.09)	1.88 (3.10)	1.99 (3.52)
T8	Control (water spray)	2.70 (6.76)	2.48 (5.47)	2.20 (3.54)	2.46 (5.26)	2.59 (6.02)
	<b>S.Em.±</b>	0.10	0.10	0.09	0.05	0.05
	<b>S.Em.± (P×T)</b>	-	-	-	0.09	0.08
	<b>CD at 5 %</b>	0.29	0.31	0.28	0.16	0.14
	<b>CD at 5 % (P×T)</b>	-	-	-	NS	NS
	<b>CV %</b>	7.39	8.82	9.68	8.57	7.30

Figure in parentheses are retransformed value whereas, those outside are  $\sqrt{x} + 0.5$  transformed values

#### 4. CONCLUSION

Among the different biopesticides tested against sucking pests of moth bean, *Beauveria bassiana*(1 x 10<sup>8</sup> cfu g<sup>-1</sup>), neem oil 0.5% and NSKE 5% was found most effective against jassid,*Empoasca motti* (Pruthi.) and whitefly,*Bemisia tabaci*(Genn.) whereas neem oil 0.5% and azadirachtin 0.15 EC was found most effective against thrips, *Caliothrips indicus*(Bagnall).

#### Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.

## REFERENCES

1. Kukvaya, D, Jakhar, B. L., Chaudhari, S. J. & Patel, B. C. (2018). Bio-efficacy of insecticides against sucking insect pests of moth bean, *Vigna aconitifolia* (Jacq.) Marechal. Journal of Entomology and Zoology Studies, 6(5), 2227-2230.
2. Anonymous (2021). Crop wise area, production and productivity of pulses from 2010-11 to 2020-21. Report published by Directorate of Pulses Development, Gov. of India, Bhopal, P. 32.
3. Satyavir, H., Jindal, S. K. & Lodha, S. (1984). Screening of moth bean cultivars against jassid, white fly and yellow mosaic virus. Annals of Arid Zone, 23, 99-103.
4. Bhatthasar, S., Khinchi, S. K., Kumawat, K. C. & Choudhary, S. (2021). Seasonal abundance of major sucking insect pests on moth bean, *Vigna aconitifolia* (Jacq.) Marechal. Journal of Pharma Innovation, 10(10), 968-971.
5. Sujatha, B. & Bharpoda, T. M. (2017). Bio-efficacy of biopesticides against sucking pests in green gram grown during *kharif*. International Journal of Pure & Applied Biosciences, 5(4), 1827-1834.
6. Singh, S. K., Singh, A. K., Singh, J. P. & Pathak V. (2018). Effect of application schedule of microbial and chemical insecticides on insect-pest control and grain yield of mung bean (*Vigna radiata* (L.) Wilczek). International Journal of Current Microbiology and Applied Sciences, 7(9), 1717-1727.
7. Chaudhary, D. M., Chaudhary, M. M. & Chaudhary, F. K. (2018). Evaluation of newer insecticidal formulation against sucking pests and effect on yield of soybean (*Glycine max* L.). International Journal of Current Microbiology and Applied Sciences, 7(8), 3834-3840.
8. Saxena HO, Tripathi YC, Pawar G, Kakkar A, Mohammad N. Botanicals as biopesticides: Active chemical constituents and biocidal action. Familiarizing with Local Biodiversity. 2014:222-40.
9. Srinivasan R, Sevgan S, Ekesi S, Tamò M. Biopesticide based sustainable pest management for safer production of vegetable legumes and brassicas in Asia and Africa. Pest management science. 2019 Sep;75(9):2446-54.
10. Bairwa DK, Sharma JK, Kumawat KC. Efficacy of insecticides, biopesticide and plant products against whitefly, *Bemisia tabaci* on mothbean, *Vigna aconitifolia*. Indian Journal of Plant Protection. 2006;34(2):210-2.

UNDER PEER REVIEW