

A PROSPECTIVE OBSERVATIONAL STUDY ON THE CLINICAL PROFILE ASSESSMENT OF MYOCARDIAL INFARCTION PATIENTS IN A TERTIARY CARE TEACHING HOSPITAL

ABSTRACT

Background: Cardiovascular diseases (CVDs) are the single largest contributor to the global burden of disease and are recognized as the leading cause of death globally. CHD has various clinical manifestations ranging from asymptomatic to acute conditions, namely acute coronary syndrome. Cardiovascular risk factors for acute coronary syndrome (ACS) are on the rise in people of Indian origin. Myocardial Infarction (MI) also called as heart attack, infarction means the death of an area of tissue because of interrupted blood supply. Myocardial Infarction has become a growing concern among the cardiovascular diseases, hence the study is conducted in order to assess the clinical profile of patients suffering from MI.

Aim: To assess the clinical presentations, risk factors, short-term complications and management of patients with MI.

Methodology: A prospective observational study was conducted in a tertiary care teaching hospital over six months. A total of 121 patients were selected based on inclusion and exclusion criteria.

Results: Among the 121 patients included, the predominant were males (63.63%). Most of the patients fell under the age group of 51-60 (n=46). The common presenting symptoms were chest pain (75.20%) followed by breathlessness (46.28%) and sweating (27.27%). Most patients had NSTEMI (62.80%) and STEMI (37.19%). Alcohol (40.49%) was major risk factor followed by smoking (38.84%) and diabetes (24.79%). The elevated biomarkers noted were Troponin I (42.14%), Troponin T (37.19%) and CK-MB (6.61%) respectively. The majority of infarctions occurred on inferior wall (30.57%). The most prevalent complication observed was heart failure (14.8%) followed by sinus tachycardia (6.61%), and sinus bradycardia (5.78%). The majority of patients received key medications like anti-platelet agents, statins and anti-coagulants during hospital stay.

Conclusion: The risk of men developing myocardial infarction is more compared to women due to various reasons. It is more likely that the diagnosis of myocardial infarction is neglected due to certain presenting symptoms like epigastric pain, syncope, vomiting, sweating etc., hence this study was conducted with an overview of creating awareness regarding the severity of myocardial infarction along with the advantages of early diagnosis.

Keywords: MI, CHD, STEMI, NSTEMI.

INTRODUCTION

Cardiovascular diseases (CVDs) are the single largest contributor to the global burden of disease and are recognized as the leading cause of death globally.¹ Among which ischemic heart disease (IHD), characterized by insufficient blood flow to the muscle tissue of the heart, is the world's biggest killer, responsible for 16% of global deaths in 2019 according to the world health organization (WHO).² CHD has various clinical manifestations ranging from asymptomatic to acute conditions, namely acute coronary syndrome (ACS).³ Cardiovascular risk factors for acute coronary syndrome (ACS) are on the rise in people of Indian origin.⁴

Acute coronary syndromes (ACSs) is a term that includes all clinical syndromes compatible

with acute myocardial ischemia resulting from an imbalance between myocardial oxygen demand and supply. ACSs are classified according to electrocardiographic (ECG) changes into:

- ST-segment elevation ACS (STEMI)
- Non-ST-segment elevation ACS (NSTEMI)
- Unstable Angina (UA).⁵

When patients with prolonged ischemic discomfort at rest are first seen, the working clinical diagnosis is that they are suffering from an acute coronary syndrome (ACS).⁶

MYOCARDIAL INFARCTION Commonly called as heart attack, infarction means the death of an area of tissue because of interrupted blood supply. Because the heart tissue distal to the obstruction dies and is replaced by noncontractile scar tissue, the heart loses some of its strength.⁷

STEMI	NSTEMI
STEMI is a medical emergency caused by acute total occlusion of an epicardial coronary artery, most often due to atherosclerotic plaque rupture/erosion and subsequent thrombus formation. ⁸	NSTEMI is defined by an elevation of cardiac enzymes (creatinine kinase MB (CK-MB) or troponin) and the absence of ST-segment elevation. Of NSTEMI patients, 25% develop Q wave MIs. ⁸

Patients with myocardial infarction present with two types of symptoms:

- *Typical symptoms* – usually include chest, arm, or jaw pain described as dull, heavy, tight or crushing.⁹
- *Atypical symptoms* – include giddiness, dyspnoea, vomiting, sweating, and epigastric pain in the absence of chest pain.¹⁰
- The median number of PCI procedures per million people in the 2023 survey was lower in middle-income compared with high-income countries (1355 vs 2330) but the number of primary PCI procedures was greater in middle-income countries (6248.8 vs 477.8), probably reflecting a greater burden of ST-elevation myocardial infarction.
- Recently, however, data in cohorts aged <40 years have shown that modestly abnormal lipid values are associated with higher risk of myocardial infarction over a median follow-up of 5.2 years.¹¹
- In 2020, approximately 19.1 million deaths were attributed to CVD globally. The age-adjusted death rate per 100,000 population was 239.8. The age-adjusted prevalence rate was 7354.1 per 1000,000.¹²
- The death rate of heart disease is estimated at approximately 31% globally, according to the World Health Organization (WHO) and over 23.6 million people worldwide may die from CHD.¹³
- Statistics suggest that among overall cardiovascular deaths, 0.9 million (68.4%) is caused by IHDs and is increasing to a greater number in the years to come.¹⁴
- Meanwhile, fatalities from stroke and **heart attack (MI)** have reached 17.7 million in the world from heart disease.¹³
- Global burden of disease studies have shown that cardiovascular diseases, and in particular CHD, are the greatest cause of morbidity and mortality in most of the world,

including both wealthy as well as poorer countries. ¹⁵

MATERIALS AND METHODS

This prospective observational study was carried out in the Department of General Medicine, Vijayanagara Institute of Medical Sciences (VIMS), Ballari, Karnataka over a period of six months (March 2023 to September 2023). Approval from Institutional Ethical Committee was taken before the commencement of the study.

Study Sample Size: Sample size was calculated by using the formula

$$n = Z^2 pq / d^2 \quad n = \text{Required sample size} \quad Z = \text{Reliability co-efficient}$$

$$n = (1.96)^2 \times 0.92 \times 0.08 / (0.05)^2 \quad p = \text{Estimated proportion} \quad d = \text{Margin of error}$$

$$n = 113$$

The minimum required sample size was 113 MI patients. 121 was the achieved sample size.

Study population:

Patients aged above 18 years with a diagnosis of Myocardial Infarction during the period of the study.

Inclusion criteria:

Patients aged above 18 years admitted with a diagnosis of Myocardial Infarction. Patients who were diagnosed with both STEMI, NSTEMI and other co morbid conditions were included in the study.

Exclusion criteria:

- Patients below 18 years
- Patients who are not willing to sign informed consent form.

Materials used:

Data collection form, Informed consent form, Patient information leaflet.

RESULTS

A total number of 121 subjects were covered during the study. Out of which 77 (63.63%) were male and 44 (36.36%) were female. The maximum number of patients were among the age group of 51-60 (n=46).




Figure 1: Age group wise distribution of patients

Among 121 patients, the majority of males had a habit of smoking (n=47) and alcohol consumption (n=47), followed by tobacco chewing (n=10). Whereas, some females had a habit of tobacco chewing (n=3) and alcohol consumption (n=2).




Figure 2: Distribution of patients according to social habits

The common presenting symptoms were chest pain (42.32%), followed by breathlessness (26.04%) and sweating (15.34%).

Presenting Symptoms	No of cases	Percentage
Chest pain	91	42.32%
Breathlessness	56	26.04%
Sweating	33	15.34%
Vomiting	9	4.18%
Abdominal pain	8	3.72%
Giddiness	7	3.25%
Epigastric pain	4	1.86%
Syncope	3	1.39%
Vertigo	2	0.93%
Palpitations	2	0.93%
Total	215	100%

Table 1: Presenting Symptoms

Among 121 patients, 45 (37.19%) had STEMI, and 76 (62.80%) NSTEMI.

The elevated biomarkers among the study subjects were like, 51 (42.14%) patients had elevated Troponin I, 45 (37.19%) patients had elevated Troponin T and 8 (6.61%) with elevated CK-MB.






Figure 3: Elevated Biomarkers

The most common site of infarction was the anterior wall, affected in 24 (19.83%) patients, 37 (30.57%) patients with inferior, 8 (6.61%) inferio-lateral, 29 (23.96%) antero-lateral, 8 (6.61%) antero-septal, 12 (9.91%) lateral and 3 (2.47%) others.



The changes in ECG have been recorded as follows: ST elevations were in 38 (31.40%) patients, ST depressions in 35 (28.92%), T-wave inversions in 30 (24.79%), pathological Q wave in 9 (7.43%), LBBB in 7 (5.78%) and 2 others (1.65%).

Among 121 patients, 49 (23.44%) were alcoholics, 47 (22.48%) smokers, 13 (6.22%) were tobacco chewers, 17 (8.13%) hypertensive patients, 30 (14.35%) diabetic patients, 14 (6.69%) patients with dyslipidemia, 13 (6.22%) patients with obesity and 26 (12.44%) did not have any risk factors.



Comorbidities are divided based on the system. 5 (4.13%) patients had central nervous system comorbidities, 3 (2.47%) endocrine system comorbidities, 16 (13.22%) respiratory system

comorbidities, 6 (4.95%) renal system comorbidities, and 6 (4.95%) other comorbidities and 85 (70.24%) patients without comorbidities.

Complications of myocardial infarction observed among the study subjects were, 18 patients with heart failure (14.8%), 8 patients with sinus tachycardia (6.61%), 7 patients with sinus bradycardia (5.78%), 7 patients with cardiogenic shock (5.78%), 2 patients with dilated cardiomyopathy (1.65%), 2 patients with atrial fibrillation (1.65%), 1 patient with atrial flutter (0.82%), and 76 (62.80%) patients had no complications.

Comorbidities	No of cases	Percentage
Central nervous system	5	4.13%
Endocrine system	3	2.47%
Respiratory system	16	13.22%
Renal system	6	4.95%
Others	6	4.95%
None	85	70.24%

Table 2: Comorbidities




Figure 6: Complications

Commonly prescribed agents – 121 (21.88%) patients received Aspirin, 115 (20.79%) Clopidogrel, 120 (21.69%) Atorvastatin, 93 (16.81%) Heparin, 46 (8.31%) Metoprolol, 32 (5.78%) Nitrates, 23 (4.15%) Streptokinase, 2 (0.36%) with ticagrelor and 1 (0.18%) with fondaparinux

Name of drugs	No of patients	Percentage
Aspirin	121	21.88%
Clopidogrel	115	20.79%
Atorvastatin	120	21.69%
Heparin	93	16.81%
Streptokinase	23	4.15%
Metoprolol	46	8.31%
Nitrates	32	5.78%
Ticagrelor	2	0.36%
Fondaparinux	1	0.18%
Total	553	100%

Table 3: Commonly prescribed agents during hospital stay

DISCUSSION

A total of 121 patients were included in the study, among the study population **males** 77 (63.63%) **predominate** over **females** 44 (36.36%) which was similar to Niveditha Alok Swamy et al,¹⁶ & most of the patients fall under the age group of 51-60 (n=46) which was nearly identical to Sapkal Harish Barsu et al,¹⁷ study.

Chest pain (75.20%) followed by breathlessness (46.28%) and sweating (27.27%) were the common symptoms observed in patients. These findings were similar to Bhagwan Das Negi et al,¹⁸ study.

Among the study population, the most common ECG interpretation **was** found as ST-elevation (31.40%) ST-depression (28.92%) T wave Inversion (24.79%) and others which **were** homogeneous with the study conducted by Sadeq Tabatabai et al.¹⁹

In the current study, the most **infarctions** occurred in inferior wall (30.57%) followed by anterio-lateral (23.96%), **anterior (19.83%) and others.** These findings were incompatible with **the** Bhagwan Das Negi et. al.,¹⁸ study.

Among the study population, **our results on the type** of MI are as follows NSTEMI (62.80%), **and** STEMI (37.19%) which is in contrast with Sadeq Tabatabai et al.,¹⁹ study. Our conclusion on biomarkers, Troponin I (42.14%), followed by Troponin T (37.19%) and CK-MB (6.61%) were distinctive **from** the Anand Premanand et. al,²⁰ study.

Among the study population, the common complications were found as HF (14.8%) followed by sinus tachycardia (6.61%), sinus bradycardia (5.78%) and others. These findings **are** in

liaison with the studies conducted by Sadeq Tabatabai et al¹⁹ and Govind Adhikari et al.²¹ respectively.

Contrary to the Niveditha Alok Swamy et al¹⁶ study, the major risk factor observed in our study was alcohol, followed by smoking & tobacco chewing.

CONCLUSION

The risk of men developing myocardial infarction is more compared to women due to various reasons. It is more likely that the diagnosis of myocardial infarction is neglected due to certain presenting symptoms like epigastric pain, syncope, vomiting, sweating etc., hence this study was conducted with an overview of creating awareness regarding the severity of myocardial infarction along with the advantages of early diagnosis.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

CONSENT

All the authors have declared that written informed consent form was obtained from all the patients.

ETHICAL APPROVAL

All the authors hereby declare that the study was conducted by the approval of the institutional ethics committee. (Reg.No – TVMCP/IEC/V PD/2022-2023/03)

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES

1. Safiri S, Karamzad N, Singh K, Carson-Chahhoud K, Adams C, Nejadghaderi SA, Almasi-Hashiani A, Sullman MJ, Mansournia MA, Bragazzi NL, Kaufman JS. Burden of ischemic heart disease and its attributable risk factors in 204 countries and territories, 1990–2019. *European journal of preventive cardiology*. 2022 Jan 1;29(2):420-31
2. Huang KS, He DX, Huang DJ, Tao QL, Deng XJ, Zhang B, Mai G, Guha-Sapir D. Changes in ischemic heart disease mortality at the global level and their associations with natural disasters: A 28-year ecological trend study in 193 countries. *Plos one*. 2021 Jul 9;16(7):e0254459.
3. Ginanjar E, Sjaaf AC, Alwi I, Sulistyadi W, Suryadarmawan E, Wibowo A, Liastuti LD.

CODE STEMI program improves clinical outcome in ST elevation myocardial infarction patients: A retrospective cohort study. *Open Access Emergency Medicine*. 2020 Nov 2:315-21

4. Gupta S, Lakhani KK, Munshi H. A study of risk factors in young patients of acute coronary syndrome. *Int J Contemporary Med Res*. 2017;4(10):2144-7
5. Dipiro JT, Talbert RL, Yee GC, Matzke GR, Wells GC, Posey LM. *Pharmacotherapy: A pathophysiologic approach: 7th edition*: NY: Mc Graw-Hill Book Company: 2008, Section 2, Chapter 18, Page No 250,252
6. Longo L, Fauci AS, Kasper L, Hauser SL, Jameson JL, Lozcalzo J. *Harrison's principles of internal medicine: 18th edition*: NY: Mc Graw-Hill Book Company: 2012; Part 10, Section 5, Chapter 244, 245, 246 and e33; Page No: 2015-2024, 33-1, 2035-2037.
7. Gerard J. Tortora, Sandra Reynolds Grabowski. *Principles of Anatomy and physiology: 8th Edition*: Harper Collins College Publishers. 1996; Unit 4, Chapter 20; The Cardiovascular System: The Heart; Page No 589.
8. Godara H, Hirbe A, Nassif M, Otepka H, Rosenstock A. *The Washington Manual of Medical Therapeutics: 34 edition*: Lippincott Williams and Wilkins: 2013; Chapter 4, Page No: 126-128, 142, 144, 145, 147, 165.
9. DeVon HA, Mirzaei S, Zègre-Hemsey J. Typical and atypical symptoms of acute coronary syndrome: time to retire the terms?. *Journal of the American Heart Association*. 2020 Apr 9;9(7):e015539.
10. Savith A. Clinical profile of acute myocardial infarction in elderly patients: A cross-sectional study. *INTERNATIONAL JOURNAL OF SCIENTIFIC STUDY*. 2015;3(6):65-8.
11. Timmis A, Aboyans V, Vardas P, Townsend N, Torbica A, Kavousi M, Boriani G, Huculeci R, Kazakiewicz D, Scherr D, Karagiannis E. *European Society of Cardiology: the 2023 Atlas of cardiovascular disease statistics*. *European Heart Journal*. 2024 Oct 7;45(38):4019-62.
12. Tsao CW, Aday AW, Almarzooq ZI, Alonso A, Beaton AZ, Bittencourt MS, Boehme AK, Buxton AE, Carson AP, Commodore-Mensah Y, Elkind MSV, Evenson KR, Ezzamel C, Ferguson JF, Generoso G, Ho JE, Kalani R, Khan SS, Kissela BM, Knutson KL, Levine DA, Lewis TT, Liu J, Loop MS, Ma J, Mussolino ME, Navaneethan SD, Perak AM, Poudel R, Rezk-Hanna M, Roth GA, Schroeder EB, Shah SH, Thacker EL, VanWagner LB, Virani SS, Voecks JH, Wang N-Y, Yaffe K, Martin SS; on behalf of the American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics-2022 update: a report from the American Heart Association [published online ahead of print Wednesday, January 26, 2022]. *Circulation*. doi: 10.1161/CIR.0000000000001052
13. Prabu S, Thiyaneswaran B, Sujatha M, Nalini C, Rajkumar S. Grid Search for Predicting Coronary Heart Disease by Tuning Hyper-Parameters. *Computer Systems Science & Engineering*. 2022 Nov 1;43(2).
14. Chaithra N. Applying the Binary Logistic Regression Model for Prediction of the Ischemic Heart Disease.
15. William S Weintraub, William E Boden, Achieving successful prevention of coronary artery disease: a worldwide call to action, *European Heart Journal*, Volume 45, Issue

16. Niveditha Alok Swamy, B Kerodi,, M S Mulimani, A P Ambali, S R Raiker. Clinical profile of myocardial infarction in elderly of rural population. *MedPulse International Journal of Medicine*. December 2019; 12(3); 131-133
17. Sapkal Harish Barsu, Chandrakant Raibhoge, Deshpande Neelima S, Harish Sapkal. Clinical profile of acute myocardial infarction in the young. *MedPulse International Journal of Medicine*. August 2019; 11(2): 148-150.
18. Negi BD, Katyayani, Kumar D. Clinical profile of patients presenting with acute ST elevated myocardial infarction presenting at tertiary health care institute of Himachal Pradesh, India. *International Journal of Recent Advances in Multidisciplinary Research*. 2021 January 30;8(1): 6566-6568
19. Tabatabai S, Bazargani N, Al Hashmi JM. Clinical Profile of Patients Presenting with Acute Myocardial Infarction in United Arab Emirates: A Tertiary Care Center Government Hospital Experience. *Dubai Medical Journal*. 2021 Apr 22;4(2):93-9.
20. Ambali AP, Bomman JV. Clinical profile of 150 elderly patients with acute myocardial infarction. *Hypertension*. 2018;36:24-0
21. Adhikari G, Baral D. Clinical profile of patients presenting with acute myocardial infarction. *Int J Adv Med* 2018;5: 228-33.
22. Aydin S, Ugur K, Aydin S, Sahin İ, Yardim M. Biomarkers in acute myocardial infarction: current perspectives. *Vascular health and risk management*. 2019 Jan 17:1-0.
23. Rathore V, Singh N, Mahat RK, Kocak MZ, Fidan K, Ayazoglu TA, Aydin Karahan YG, Onk D, Akar E, Yolcu A. Risk factors for acute myocardial infarction: a review. *Ejmi*. 2018 Jan;2(1):1-7
24. Zhang L, Hailati J, Ma X, Liu J, Liu Z, Yang Y, He P, Wulasihan M. Analysis of risk factors for different subtypes of acute coronary syndrome. *Journal of International Medical Research*. 2021 May;49(5):03000605211008326
25. Dai H, Much AA, Maor E, Asher E, Younis A, Xu Y, Lu Y, Liu X, Shu J, Bragazzi NL. Global, regional, and national burden of ischaemic heart disease and its attributable risk factors, 1990–2017: results from the Global Burden of Disease Study 2017. *European Heart Journal-Quality of Care and Clinical Outcomes*. 2022 Jan;8(1):50-60
26. Wagh Y, Wagh S, Gawale Y, Gawale P. Clinical profile of patients presenting with acute coronary syndromes at tertiary healthcare hospital in Northern Maharashtra. *Asian Journal of Medical Sciences*. 2022 Aug 1;13(8).
27. Shanchay MS, Sultana F, Begum A, Miah MT. Atypical Presentation of Acute Myocardial Infarction. *Journal of Medicine*. 2021 Dec 1;22(2):88-94.
28. Sushritha K, Sharma A, Suman K, Goud PH, Akari S. Epidemiological Study of Risk Factors in Myocardial Infarction Patients. *Journal of Young Pharmacists*. 2020;2020(2s):s108.
29. Islam MN, Chowdhury SN, Rahman MS, Hossain SM. Short Term Complications of Acute Myocardial Infarction in a Tertiary Hospital. *Medicine Today*. 2021 Feb 25;33(1):42-4.
30. Ginanjar E, Sjaaf AC, Alwi I, Sulistyadi W, Suryadarmawan E, Wibowo A, Liastuti LD.

CODE STEMI program improves clinical outcome in ST elevation myocardial infarction patients: A retrospective cohort study. *Open Access Emergency Medicine*. 2020 Nov 2;3:15-21

31. Ferry AV, Anand A, Strachan FE, Mooney L, Stewart SD, Marshall L, Chapman AR, Lee KK, Jones S, Orme K, Shah AS. Presenting symptoms in men and women diagnosed with myocardial infarction using sex-specific criteria. *Journal of the American Heart Association*. 2019 Sep;8(17):e012307.
32. Bahall M, Seemungal T, Khan K, Legall G. Medical care of acute myocardial infarction patients in a resource limiting country, Trinidad: a cross-sectional retrospective study. *BMC Health Services Research*. 2019 Dec;19(1):1-1.
33. Lindow T, Pahlm O, Khoshnood A, Nyman I, Manna D, Engblom H, Lassen AT, Ekelund U. Electrocardiographic changes in the differentiation of ischemic and non-ischemic ST elevation. *Scandinavian Cardiovascular Journal*. 2020 Mar 3;54(2):100-7.
34. Ralapanawa U, Kumarasiri PV, Jayawickreme KP, Kumarihamy P, Wijeratne Y, Ekanayake M, Dissanayake C. Epidemiology and risk factors of patients with types of acute coronary syndrome presenting to a tertiary care hospital in Sri Lanka. *BMC cardiovascular disorders*. 2019 Dec;19(1):1-9.
35. Gebhard C, Gebhard CE, Stähli BE, Maafi F, Bertrand MJ, Wildi K, Fortier A, Galvan Onandia Z, Toma A, Zhang ZW, Smith DC. Weather and risk of ST-elevation myocardial infarction revisited: Impact on young women. *PLoS One*. 2018 Apr 9;13(4):e0195602.
36. Gupta S, Lakhani KK, Munshi H. A study of risk factors in young patients of acute coronary syndrome. *Int J Contemporary Med Res*. 2017;4(10):2144-7.
37. Sharma S, Rashid A, Ashraf M, Ismail M, Tanvir M, Sharma P, Ajaz S. Clinical Profile of Acute Coronary Syndromes (ACS) in North Indian Population: A Prospective Tertiary Care Based Hospital Study. *Annals of International Medical and Dental Research*.;3(5):50.
38. Seetharama N, Mahalingappa R, Ranjith KG, Veerappa V, Aravindh CL. Clinical profile of acute myocardial infarction patients: a study in tertiary care centre.
39. Jacobsen L, Grenne B, Olsen RB, Jortveit J. Feasibility of prehospital identification of non-ST-elevation myocardial infarction by ECG, troponin and echocardiography. *Emergency Medicine Journal*. 2022 Sep 1;39(9):679-84
40. Puelacher C, Gualandro DM, Glarner N, Lurati Buse G, Lampart A, Bolliger D, Steiner LA, Grossebacher M, Burri-Winkler K, Gerhard H, Kappos EA. Long-term outcomes of perioperative myocardial infarction/injury after non-cardiac surgery. *European Heart Journal*. 2023 May 14;44(19):1690-701.
41. Shehab A, AlHabib KF, Bhagavathula AS, Hersi A, Alfaleh H, Alshamiri MQ, Ullah A, Sulaiman K, Almahmeed W, Al Suwaidi J, Alsheikh-Ali AA. Clinical presentation, quality of care, risk factors and outcomes in women with acute ST-elevation myocardial infarction (STEMI): an observational report from six middle Eastern countries. *Current vascular pharmacology*. 2019 Jul 1;17(4):388-95
42. Baechli C, Koch D, Bernet S, Gut L, Wagner U, Mueller B, Schuetz P, Kutz A. Association of comorbidities with clinical outcomes in patients after acute myocardial infarction. *IJC Heart & Vasculature*. 2020 Aug 1;29:100558