

Original Research Article

Assessment and delineation of GPS-GIS based Soil Fertility Maps of Instructional Farm, Malegaon tehsil Baramati

ABSTRACT

The present investigation entitled "Assessment and delineation of GPS-GIS based fertility maps of Instructional Farm, Malegaon tehsil Baramati" was conducted in the year 2023-24 at division of Soil Science, Dr. Sharadchandra Pawar College of Agriculture, Baramati with an objectives to assess the soil available nutrient status and delineation of GIS based maps of soil available nutrients. Using Arc-GIS 10, Version 10.8, fertility maps of the available macro nutrients were created. The area under the Instructional Farm had soil pH ranged from 7.11 to 8.32, categorized into the slightly alkaline to moderately alkaline categories while the EC varied from 0.21 to 1.99 dSm^{-1} , indicating that most of the soils were non-saline. The soil's organic carbon and calcium carbonate contents ranged from 0.21 to 0.97 per cent and 1.50 to 19.75 per cent and they were classified into low to high and low to very high categories respectively. In the Instructional Farm, the ranges of available nitrogen, phosphorus and potassium content were 85.50 to 360.75, 23.29 to 80.08 and 202.5 to 968.68 kg ha^{-1} respectively. The Instructional Farm soils had very low to moderate levels of available nitrogen, moderately high to very high levels of available phosphorus and moderately high to very high levels of available potassium.

Key words: *Soil, Fertility maps, GPS-GIS, Instructional Farm*

1. INTRODUCTION

A necessary resource soil is referred to as the 'Soul of Infinite Life'. The most important and valuable natural resource that keeps life alive on earth is soil. An inch of top soil requires nearly ten thousand years to produce but in today's world, pollution and soil contamination are main issues. The development and application of soil nutrient management technologies that improves plant productivity and soil quality are biggest challenge for today. The adoption of green revolution technologies has led to dependence on different synthetic inputs predominantly fertilizers which are derived from fossil fuels. Excessive use of chemical fertilizers negatively impacts soil, leading to a significant decline in soil fertility.

Major nutrients also known as macronutrients are those that are needed in high amounts like C, H, O, N, P, K, Ca, Mg and S are some of them. C, H and O are known as structural elements which accounts for 95 percent of the plant's dry matter weight and provided through carbon dioxide (CO_2) and water (H_2O). Macronutrients includes both primary and secondary nutrients that are derived from soil, fertilizers and other sources. The key and vital primary nutrients nitrogen, phosphorus and potassium are introduced into the soil either during sowing or through subsequent intercultural operations. Imbalances in nutrient application either deficiency or excess can result in various disorders affecting both the plant and its fruit.

Retention and availability of nutrients, minerals content, presence of organic matter and soils environment are significantly influenced by the chemical characteristics of the soils. The

chemistry of soils involves the chemical reactions that influence the development of plants, animals and humans (Tale, 2015). Chemical characteristics of soils are developed from weathering processes, parent material transportation and soil forming processes. The chemical properties of soils includes different parameters such as pH, electrical conductivity (EC), cation exchange capacity (CEC), exchangeable Ca and Mg, organic carbon content and calcium carbonate content. The amount of organic matter, calcium carbonate, microbial activity, pH changes, amount of clay and soil moisture status all affect the soil's nutrient availability (Naphadeet *et al.* 2022).

The preparation of soil fertility maps is now possible through the applications of modern technologies such as GPS and GIS. The United States military administrated the Global Positioning System, a space-based navigation and tracking system that accurately determines the location of the object on earth's surface using geographical coordinates. A Geographic Information System (GIS) is a computer system used for capturing, querying and displaying geographic data. It is easy to change the data regarding to fertility status of an area even after the soil fertility maps are prepared (Surabhi *et al.* 2017).

2. MATERIAL AND METHODS

The **Instructional Farm**, Malegaon tehsil Baramati is located between 18° 08' N latitude and 74° 31' E longitude. Soils are resultant of the igneous rocks viz. Basalt (Deccan trap) which is basic in nature containing mainly feldspars (plagioclase), augite and small amount of titaniferrous magnetite mineral. The **climate** is classified as semi-arid tropical. Total 104 Geo-referenced surface (0-22.5 cm) soil samples representing **different soils** were collected from Instructional Farm, Malegaon tehsil Baramati. The latitude and longitude of sampling sites were recorded with the help of differential Global Positioning System (GPS). The delineation of the area for mapping different nutrient levels in soils was carried out using Arc-GIS 10, Version 10.8 software.

The pH was measured by in 1:2.5 soil water suspension using glass electrode pH meter and EC (dSm^{-1}) was measured in the supernatant solution of 1:2.5 soil water suspension using conductivity meter (Jackson, 1973). Organic carbon by wet oxidation method (Walkely and Black, 1934). Available Nitrogen was estimated by alkaline permanganate method of (Subbiah and Asija, 1956). Available phosphorus was extracted with 0.5 M NaHCO_3 solution buffered at pH-8.5. (Watanabe and Olsen 1965). Available Potassium was estimated by shaking the requisite amount of soil sample with 1 N neutral ammonium acetate solution at pH-7.0, (Jackson, 1973)

3. RESULT AND DISCUSSION

3.1 Soil reaction (pH)

The Soil pH of the Instructional Farm ranged from 7.11 to 8.32 with mean of 7.8. The alkaline nature of the soil could be attributed to the semi arid climatic condition of study area, annual evaporation rate is higher in comparison to rainfall hence there is less possibility of leaching down of soluble salts which results in accumulation of salts in soil's root zone. The

Comment [A1]: Mention area?

Comment [A2]: Land use?

Comment [A3]: Criteria ?

similar results were recorded by Shirgaveet *et al.* (2015) in soils around Arjunnagar of Kolhapur district, Jagtap *et al.* (2018) in Ajang village of Dhule tehsil of Dhule district (MH).

3.2 Electrical conductivity (EC)

The range and mean of electrical conductivity of soils from the Instructional Farm were 0.21 to 1.99 dSm⁻¹, 0.73 dSm⁻¹ respectively. The majority of the soils exhibited normal electrical conductivity, likely due to careful management of irrigation water. The accumulation of soluble salts on the surface of soil along with Ca⁺² and Mg⁺² carbonates may have led to increase EC greater than 1 dSm⁻¹. Naphadeet *et al.* (2022) reported comparable results in soils of Jalgaon district of Maharashtra and Ushasri *et al.* (2019) in soils of Bhudargad block, Kolhapur district. Similar observations were noted by Mandavgadeet *et al.* (2015) in soils of Jintur, Selu and Patthri tehsils of Parbhani district.

3.3 Organic carbon

The organic carbon of soils from the Instructional Farm was ranged from 0.21 to 0.97 per cent with mean of 0.62 per cent. Moderate to moderately high levels of organic carbon in soils were observed which may be caused by the addition of organic residues in soil. Wide variation in content of organic carbon might be due to high temperature dominant during summer increase the decomposition rate of organic matter. Aich *et al.* (2017) reported corresponding results in the soil of Organic Farm at College of Agriculture, Pune. Kumar and Palwe (2017) documented similar observations in the soils of Girawali village of Ambegaon tehsil, Pune (MH). Kashiwaret *et al.* (2023) also recorded similar findings in Pauni block of Bhandara district (MH).

Table 1 Chemical properties status in soils of Instructional farm, Malegaon tehsil Baramati.

Particular	pH (1:2.5)	EC (dSm ⁻¹) (1:2.5)	Organic Carbon (%)	CaCO ₃ (%)
Mean	7.85	0.73	0.62	8.98
Standard Error	0.02	0.05	0.02	0.51
Standard Deviation	0.17	0.48	0.19	5.22
Minimum	7.11	0.21	0.21	1.50
Maximum	8.32	1.99	0.97	19.75
CV%	2.18	65.39	30.05	58.12
Category	Slightly alkaline (84.61)	Normal (78.85)	Low (17.30)	Low (7.69)
	Moderately Alkaline (15.39)	Poor Seed emergence (21.15)	Moderate (27.20)	Moderate (21.16)
			Moderately high (38.25)	Moderately high (37.5)
			High (17.25)	Very high (33.65)

Comment [A4]: Use recent units

(Total no. soil samples analyzed - 104, Figures in Parenthesis are in Per cent)

3.4 Calcium carbonate

The content of calcium carbonate in the soils of the Instructional Farm ranged from 1.5 per cent to 19.75 per cent where mean was 8.98 per cent. Low precipitation and accelerated rate of evaporation in the semi-arid climatic condition within the study area led to more accumulation and precipitation of calcium carbonate resulting in high calcareousness in soil. Similar results revealed by Surabhi *et al.* (2017) in the soil of Shirol tehsil of Kolhapur district. Ingole *et al.* (2018) recorded similar findings in the soil of Nanded district (Maharashtra) and also reported by Hadoleet *et al.* (2019) in the soil of Jalgaon district (Maharashtra).

3.5 Available nitrogen

Range of available nitrogen was 85.5 to 360.5 kg ha⁻¹ with mean of 244.23 kg ha⁻¹ respectively. The low availability of nitrogen in most of the soils may be due to higher temperature in the semi-arid climate, which may have accelerated the rate of denitrification resulted in low levels of available nitrogen. The variation in soil nitrogen content may be related to management of soil, application of fertilizers and farm yard manure to previous crops etc. Similar results were recorded by Nagawade (2014) in Central Research Farm MPKV, Rahuri, Surabhi *et al.* (2017) in soils of Shirol tehsil of Kolhapur and Kondvilkar and Thakre (2018) in soils of Sakri tehsil of Dhule.

Table 2. Available nutrient status in soils of Instructional farm, Malagaon tehsil Baramati.

Particular	Available Nutrients (kg ha ⁻¹)		
	N	P	K
Mean	244.23	47.23	566.38
Standard Error	5.62	1.48	21.76
Standard Deviation	57.30	15.10	221.89
Minimum	85.50	23.29	202.5
Maximum	360.75	80.08	968.68
CV%.	23.46	31.98	39.18
Category	Very low (3.85)	Moderately high (7.69)	Moderately high (8.66)
	Low (65.38)	High (18.27)	High (4.80)
	Moderate (30.77)	Very high (74.04)	Very high (86.54)

(Total no. soil samples analyzed - 104, Figures in Parenthesis are in Per cent)

3.6 Available phosphorus

The status of available phosphorus in the soils of the Instructional Farm ranged from 23.29 to 80.08 kg ha⁻¹ with mean of 47.23 kg ha⁻¹. The high range of soils available phosphorus under study area may be mostly caused by several factors including past fertilization practices, organic matter content, soil texture, various soil management practices, semi-arid environment

with low rainfall. The continuous use of high-analysis fertilizers especially DAP, SSP has led to the phosphorus build up and resulted in high levels of available phosphorus in the soils. Similar findings were noted by Surabhi *et al.* (2017) in soils of the Shirol tehsil of Kolhapur district and Rajmaniet *al.* (2020) in soils of the KrishiVigyan Kendra, Palem, Telangana.

3.7 Available potassium

The available potassium in soils of the Instructional Farm ranged from 202.50 to 968.68 kg ha⁻¹ with mean of 566.38 kg ha⁻¹. The available potassium content of significant portion of the study areas soil was very high, this could be attributed to the predominance of mica and feldspar minerals in the parent materials. Similar outcomes of available potassium was documented by Pulkeshiet *al.* (2012) in soils of Mantagani village under northern zone of Karnataka, Palwe and Yelwe (2018) in soil of blocks A and B of central campus MPKV, Rahuri and also similar results recorded by Patil *et al.* (2016) in soil of Panhala tehsil of Kolhapur.

UNDER PEER REVIEW




Fig. 1 Location map of Instructional Farm, Malegaon tehsil Baramati




Fig. 2 Map of soil pH of Instructional Farm, Malegaon tehsil Baramati




Fig. 3 Map of soil EC of Instructional Farm, Malegaon tehsil Baramati




Fig. 4 Map of soil organic carbon of Instructional Farm, Malegaon tehsil Baramati




Fig. 5 Map of Calcium Carbonate of Instructional Farm, Malegaon tehsil Baramati




Fig. 6 Map of soil available nitrogen of Instructional Farm, Malegaon tehsil Baramati




Fig. 7 Map of soil available phosphorus of Instructional Farm, Malegaon tehsil Baramati




Fig. 8 Map of soil available potassium of Instructional Farm, Malegaon tehsil Baramati

CONCLUSIONS

The study reveals the several findings into the nutrient level of soils at the Instructional Farm, Malegaon tehsil Baramati. The soils were slightly alkaline with most samples showed suitable electrical conductivity and good for crop growth. Organic carbon content varied widely but a major proportion of soils were moderately high in organic carbon. calcium carbonate content was also moderate to very high.

Nitrogen levels were low, likely due to high temperatures accelerating denitrification processes. Availability of phosphorus levels were high to very high, possibly due to past fertilizer application practices. Potassium levels were also very high attributed to the presence of potassium rich minerals in the parent material.

DISCLAIMER

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts. COMPETING INTERESTS: Authors have declared that no competing interests exist.

References

1. Aich, V., Patil, G. D., More, N. B. and Udayana, S. K. (2017). GPS-GIS based soil maps of micronutrients status in organic farms at College of Agriculture, Pune (M.S.), India. *International Journal of Current Microbiology and applied Sciences*. **6**(3): 855-861.
2. Alison, L. E. and Moodier, C. D. (1965). Method of soil analysis, Part 2; Chemical and Microbiological Properties.
3. Hadole, S. S., Sarap, P. A., Lakhe, S. R., Dhule, D. T., Parmar, J. N. (2019). Status of micronutrients in soil of Jalgaon district, Maharashtra, India. *International Journal of Current Microbiology and Applied Science*. **8**(7): 1432-1439.
4. Ingole, A. J., Vaidya, P. H., Aundhakar, A. V. and Ghode, M. K. (2018). Effect of moisture and calcium carbonate content on chemical properties of soil and yield of cotton in Nanded district of Maharashtra. *Journal of Pharmacognosy and Phytochemistry*. **7**(5): 1971- 1976.
5. Jackson, M. L. (1973). Soil Chemical Analysis - Advanced Course, 2nd Edn. Publ. by the author, University of Wisconsin, Madison, USA.
6. Jagtap, M., Chaudhari, R. D., Thakare, R. and Patil, T. D. (2018). Mapping of soil micronutrient status based on GPS- GIS and biological properties of Ajang village of Dhule tehsil of Dhule district Maharashtra. *Journal of Pharmacognosy and Phytochemistry*. **27**(5): 3270-3275.
7. Kashiwar, S. R., Kundu, M. C. and Dongarwar, U. R. (2023). GIS based spatial mapping of soil nutrients status of Pauni block of Maharashtra, India. *Annals of Plant and Soil Research*. **25**(3): 446-454.
8. Kondvilkar, N. and Thakre, R. (2018). Mapping of soil macro and secondary nutrients by GIS in Sakri tehsil of Dhule district (M.S.). *Journal of Pharmacognosy and Phytochemistry*. **7**(4): 3158-316.
9. Kumar, S. and Palwe, C. R. (2017). Soil fertility maps of Girawali village of Ambegaon tehsil, Pune district, Maharashtra (India). *Trends In Biosciences*. **10**(31): 6446-6451.
10. Mandavgade, R. R., Waikar, S. L., Dhamak, A. L. and Patil, V. D. (2015). Evaluation of micronutrient status of soils and their relation with some chemical properties of soils of Northern tehsils (Jintur, Selu and Pathri) of Parbhani district. *IOSR Journal of Agriculture and Veterinary Science*. **8**(2): 38-41.

11. Nagawade, S. G. (2014). GPS- GIS based soil fertility map of central farm of MPKV, Rahuri (M.S.) 2014. Student of M.Sc. (Agri.) Department of Soil Science and Agricultural Chemistry, Rahuri. Thesis submitted to Mahatma PhuleKrishiVidyapeeth, Rahuri.
12. Naphade, M., Shinde, R. and Suryavanshi, R. T. (2022). Micronutrient status and it's correlation with chemical properties of soil of Jalgaon district, Maharashtra. *The Pharma Innovation Journal*. **11**(11): 1711-1720.
13. Palwe, C. R. and Yelwe, L. J. (2018). GPS GIS based soil fertility status of Mahatma PhuleKrishiVidyapeeth, Rahuri, (M.S.), India. *International Journal of Agriculture Innovations and Research* Volume. **7**(2): 2319-1473.
14. Panse, V. G. and Sukhatme, P. V. (1985). Statistical methods for agricultural workers. ICAR Publication, New Delhi. 3 rd Revised Edition.
15. Pulkeshi, H. B. P., Patil, P. L., Dasog, G. S., Radder, B. M., Bidari, B. I. and Mansur, C. P. (2012). Mapping of nutrients status by GIS in Mantagani village under northern transition zone of Karnataka. *Karnataka Journal of Agriculture Science*. **25**(3): 332-335.
16. Rajamani, K., Hari, N. and Rajashekar, M. (2020). Soil fertility evaluation and GPS-GIS based soil nutrient mapping of KrishiVigyan Kendra, Palem, Telangana. *International Journal of Pure and Applied Chemistry*. **21**(23): 139-145.
17. Shirgave, P. and Ramteke, A. (2015). Physiochemical status of fertile soil around Arjunnagar, district Kolhapur, Maharashtra. *International Journal of Chemical Studies*. **3**(2): 98-101.
18. Surabhi, H. K., Annapurna, M., Kondvilkar, N. B. and Pawar, R. B. (2017). Mapping of soil macro and secondary nutrients by GIS in Shirol tehsil of Kolhapur district. (M.S.). *Journal of chemical studies*. **5**(6): 892-896.
19. Ushashri, B. and Kumar, M. (2019). GPS-GIS based soil fertility maps of Bhudargadtehsil of Kolhapur district (M.S) India. *International Journal of Current Microbiology and Applied Sciences*. **8**(09): 523-532.
20. Walkley, A. J. and Black, A. I. (1934). Estimation of organic carbon by chromic acid titration method. *Soil Science*, 25:255-259.