

HEPATOPROTECTIVE INVESTIGATION OF *FICUS LYRATA* AND *FICUS ELASTICA* LEAVES EXTRACT

ABSTRACT

In hepatotoxic Albino rats caused by isoniazid and paracetamol, the hydroalcoholic extract of *Ficuslyrata* and *Ficus elastic* leaves was evaluated for hepatoprotective activity. The degree of protection was measured by estimating biochemical parameters such as serum glutamate Oxaloacetate transaminase (SGOT), serum glutamate pyruvate transaminase (SGPT), ALP and the level of total serum bilirubin. In rats that had been toxicated with isoniazid and paracetamol. Hydroalcoholic extract (100 mg/kg, 200 mg/kg) demonstrated a noteworthy hepatoprotective effect in a dose-dependent manner. The extract's hepatoprotective benefits were on par with those of the prescribed medication silymarin 2.5 mg/kg body weight.

Key Words: *Ficuslyrata*, *Ficus elastic*, Hepatoprotective activity

1. Introduction

Herbal remedies derived from nature are carefully considered to be safe and effective natural treatments for a range of illnesses. The purpose of this study was to demonstrate the hepatoprotective, antioxidant, and preliminary chemical screening properties of *Ficus elastic* and *lyrata*. After thoroughly washing the dried Leaves material under running water, an electronic grinder was used to grind it. The powder was extracted in stages using the maceration process and an ethanol-water solvent combination. *Ficuslyrata* and *Ficus elastic* were evaluated using various standards criteria, including phytochemical screening, percentage loss, yield, and organoleptic evaluation.

The India rubber plant, or *Ficuselastica*, is a huge tree in the Moraceae family that was formerly a significant supplier of subpar natural rubber. In the early 20th century, the unrelated rubber tree primarily took its position as a source of rubber. Although it is native to Southeast Asia, the India rubber plant is widely grown indoors in pots elsewhere. *Ficuslyrata* is a species of plant in the mulberry and fig family Moraceae. It is sometimes referred to as the fiddle-leaf fig, banjo fig, fiddle-leaved fig tree, lyre leaf fig tree, or lyre-leaved fig tree.

Ficuslyrata and *Ficus elastic* leaf are significant medicinal plants that are used to cure a variety of illnesses in traditional medicine. For two reasons, the liver may be regarded as the most significant organ in drug toxicity: first, it serves as a functional barrier between the site of absorption and the systemic circulation and is a key location for the metabolism and removal of foreign substances; second, these characteristics also make the liver a prime target for drug toxicity. Consequently, drug-induced liver damage presents a significant clinical challenge. Since the liver is essential for the detoxification and excretion of several endogenous and exogenous substances, damage to it or impairment of its function can have a number of negative effects on health.. Management of liver diseases is still a challenge to modern medicine (Handa and Sharma, 1990).

2. METHODOLOGY

2.1. Plant material

2.1.1 Selection&Collection of Plant Material

Ethnobotanical surveys were carried out in tribal localitie ofSamastipurDistrict north Bihar region. Fresh Leaves of *Ficus lyrata* and *Ficus elastic* were collected from area adjoining forests of SamastipurDistrict north Bihar region in the month of November.

2.1.2 Drying &StorageCollection of Plant Material

Leaves of *Ficus lyrata* and *Ficus elastic* was carried out in sun but under the shade.Leaves of *Ficus lyrata* and *Ficus elastic* were preserved in plastic bags and closed tightly and powdered as per the requirements.

2.2. Extraction procedure

100 grams of dried Ficuslyrata plant material and 100 grams of Ficus elastic were thoroughly extracted using the maceration process. Using a solvent mixture of ethanol-water(70:30),using maceration process for 48 hrs, filtered and dried using vaccum evaporator at 40⁰C. 70% ethanol was used to extract plant material constantly and for 48 hours at room temperature. It was proved that the extraction technique left no trace upon evaporation by adding a few drops of extract from the thimble. (Pradhan *et al.*, 2010, Ansari 2001, Mukherjee 2007).

The following formula was used to calculate the chosen plant materials' percentage yield. The percentage yield of each extract was calculated by using following formula:

$$\text{Percentage Yield} = \frac{\text{Weight of Extract}}{\text{Weight of Powder drug taken}} \times 100$$

2.3 Selection of animals

Swiss albino rats weighing between 150 and 200 g of were employed in the current study. Before the trial started, the animals were given two weeks to acclimate. They were then kept in typical laboratory settings, with a temperature of $25 \pm 2^\circ\text{C}$, a relative humidity of 45–65%, and a 12-hour light/dark cycle. Throughout the investigation, the animals were given unlimited water and regular laboratory animal nutrition. The Institution Animal Ethical Committee formally accepted the protocol for animal experimentation. The Organization for Economic Co-operation and Development (OECD) guideline No. 420 was used while testing for acute oral toxicity. (CPCSEA NO.1582/PO/Re/S/11/CPCSEA) In this investigation, water was available to the Swiss albino rats at all times during an overnight fast. For the first 24 hours and then for the following 14 days, the animals were monitored for death or any aberrant behavior after the extract was given orally at a concentration of 2000 mg/kg body weight. There were further behavioral, neurological, and autonomic reactions seen.

2.4 Isoniazid and Paracetamol induced hepatoprotective activity of Leaves extract of *Ficus lyrata* and *Ficus elastic*,

2.4.1 Experimental designs for *Ficus lyrata* leaves Extract Isoniazid induced hepatoprotective

Group –I: Normal control (Sterile distilled water ml/kg, p.o.)

Group –II: Isoniazid solutions were prepared in sterile distilled water (100 mg/kg, p.o.)

Group -III: *Ficus lyrata* Extract (100mg/kg, p.o.) + Isoniazid (100 mg/kg, p.o.)

Group –IV: *Ficus lyrata* Extract (200mg/kg, p.o.) + Isoniazid (100 mg/kg, p.o.)

Group –V: Silymarin (2.5 mg/kg, p.o.) + Isoniazid (100 mg/kg, p.o.)

Five groups, each with six creatures, were created from the animals. Group –I received 1ml/kg p.o. of sterile distilled water was given to the first group. Group II was received Isoniazid solutions at a dose of 100 mg/kg p.o. Group-III and Group -IV were received 100 mg/kg and 200 mg/kg of

hydroalcoholic leaves extract of *Ficuslyrata*, respectively. For 21 days, Group 5 was given silymarin (2.5 mg/kg p.o.) once daily. The animal was anesthetized with ether after 21 days in order to draw blood from the retroorbital plexus, and it was subsequently slaughtered while still under ether anesthesia in order to remove the liver. Numerous biochemical analyses were conducted. (Jiang *et al.*, 2004; Saleem *et al.*, 2008).

2.4.2 Experimental designs for *Ficus lyrata* leavesExtract Paracetamol induced hepatoprotective

Group –I: Normal control (Sterile distilled water ml/kg, p.o.)

Group –II: Paracetamol solutions were prepared in sterile distilled water (50 mg/kg, p.o.)

Group -III:*Ficus lyrata* Extract (100mg/kg, p.o.) + Paracetamol, (50 mg/kg, p.o.)

Group –IV: *Ficus lyrata* Extract (200mg/kg, p.o.) + Paracetamol, (50 mg/kg, p.o.)

Group –V: Silymarin (2.5 mg/kg, p.o.) + Paracetamol, (50 mg/kg, p.o.)

Five groups, each with six creatures, were created from the animals. Group –I received 1 ml/kg p.o. of sterile distilled water was given to the first group. Group II was received paracetamol solutions at a dose of 50 mg/kg p.o. Group-III and Group -IV were received 100 mg/kg and 200 mg/kg of hydroalcoholic leaves extract of *Ficuslyrata*, respectively. For 21 days, Group 5 was given silymarin (2.5 mg/kg p.o.) once daily. The animal was anesthetized with ether after 21 days in order to draw blood from the retroorbital plexus, and it was subsequently slaughtered while still under ether anesthesia in order to remove the liver. Numerous biochemical analyses were conducted.

2.4.3 Experimental designs for *Ficus elastic* leavesExtractIsoniazid induced hepatoprotective

Group –I: Normal control (Sterile distilled water ml/kg, p.o.)

Group –II: Isoniazid solutions were prepared in sterile distilled water (100 mg/kg, p.o.)

Group -III:*Ficus elastic* Extract (100mg/kg, p.o.) + Isoniazid (100 mg/kg, p.o.)

Group –IV:*Ficus elastic* Extract (200mg/kg, p.o.) + Isoniazid (100 mg/kg, p.o.)

Group –V: Silymarin (2.5 mg/kg, p.o.) + Isoniazid (100 mg/kg, p.o.)

Five groups, each with six creatures, were created from the animals. Group –I received 1 ml/kg p.o. of sterile distilled water was given to the first group. Group II was received Isoniazid solutions at a dose

of 100 mg/kg p.o. Group-III and Group -IV were received 100 mg/kg and 200 mg/kg of hydroalcoholic leaves extract of *Ficus elastic*, respectively. For 21 days, Group 5 was given silymarin (2.5 mg/kg p.o.) once daily. The animal was anesthetized with ether after 21 days in order to draw blood from the retroorbital plexus, and it was subsequently slaughtered while still under ether anesthesia in order to remove the liver. Numerous biochemical analyses were conducted.

2.4.4 Experimental designs for *Ficus elastic* leaves Extract Paracetamol induced hepatoprotective

Group –I: Normal control (Sterile distilled water ml/kg, p.o.)

Group –II: Paracetamol solutions were prepared in sterile distilled water (50 mg/kg, p.o.)

Group -III: *Ficus elastic* Extract (100mg/kg, p.o.) + Paracetamol, (50 mg/kg, p.o.)

Group –IV: *Ficus elastic* Extract (200mg/kg, p.o.) + Paracetamol, (50 mg/kg, p.o.)

Group –V: Silymarin (2.5 mg/kg, p.o.) + Paracetamol, (50 mg/kg, p.o.)

Five groups, each with six creatures, were created from the animals. Group –I received 1 ml/kg p.o. of sterile distilled water was given to the first group. Group II was received paracetamol solutions at a dose of 50 mg/kg p.o. Group-III and Group -IV were received 100 mg/kg and 200 mg/kg of hydroalcoholic leaves extract of *Ficus elastic*, respectively. For 21 days, Group 5 was given silymarin (2.5 mg/kg p.o.) once daily. The animal was anesthetized with ether after 21 days in order to draw blood from the retroorbital plexus, and it was subsequently slaughtered while still under ether anesthesia in order to remove the liver. Numerous biochemical analyses were conducted.

3. RESULT AND DISCUSSION

3.1 Result of Percentage yield

The percentage yield of leaf extracts from *Ficuslyrata* was displayed in Table No. The ethanol-water extract had a greater yield of 12.39%, respectively. The ethanol-water leaf extract yielded 8.15%, which was comparable to the *ficus elastic* of the leaves extract.

Table 1: Percentage yield of leaves of *Ficuslyrata* and *Ficus elastic*

S.N.	Plant Name	Percentage yield (%)
1	<i>Ficuslyrata</i>	12.39

2	<i>Ficus elastic</i>	8.15
---	----------------------	------

3.2In –Vivo hepatoprotective activity of *Ficuslyrata*&*Ficus elastic* leaves extract

Dose-dependent hepatoprotective effect was demonstrated by both test groups, i.e., high dose and low dosage treated groups. Liver activity improved in the test groups that received only the plant extract. It is evident that "*Ficuslyrata* and *Ficus elastic* leaves" are plants with hepatoprotective properties. According to this study, the leaves of *Ficuslyrata* and *Ficuselastica* significantly reduce the hepatotoxicity caused by the medications used to treat tuberculosis. *Ficuslyrata* and *Ficus elastic* leaves may have a hepatoprotective effect because of their antioxidant capability, which implies that plant extract may be helpful in preventing liver damage brought on by oxidative stress.

Elevated blood levels of SGPT, SGOT, and ALP activities indicate hepatocyte injury to the liver and, in turn, indirect impairment of liver function after APAP-induced hepatotoxicity. Following APAP treatment, there was a significant ($p < 0.05$) increase in SGPT, SGOT, and ALP activities, as shown in Table. *Ficuslyrata* and *Ficus elastic* leaf extract treatments at 100 and 200 mg/kg both markedly decreased the increase of these enzymes ($p < 0.05$). The observed decrease in hepatic enzymes was comparable to both the control group's and the group that received silymarin pretreatment. The apparent leakage of cellular enzymes into plasma is one of the classic indicators of hepatic injury or damage.

Furthermore, the presence or lack of particular enzymes in the bloodstream can be used to determine the kind and extent of liver damage. Alkaline phosphatase, aspartate aminotransferase, and alanine aminotransferase are frequently utilized as marker enzymes in general alanine aminotransferase measurements to assess APAP-induced hepatotoxicity. This study's improvement in SGPT, SGOT, ALP, and serum bilirubin levels demonstrates the hepatoprotective impact of *Ficuslyrata* and *Ficus elastic* leaves. Serum bilirubin increases caused by isoniazid, paracetamol, and *Ficuslyrata* and *Ficus elastic* leaf extract are suppressed. Transaminase levels have been shown to rise following Isoniazid therapy in earlier investigations (Reddy et al., 2013). The increase is time-dependent, with a significant elevation observed after 48 hours ($p < 0.05$), indicating severe

hepatocellular damage from these typically cytoplasmic enzymes leaking into the bloodstream. (Asha, *et al.*, 2004).

3.2 Hepatoprotective activity of *Ficuslyrataleaves* extract in Isoniazid induced hepatotoxicity

Hepatoprotective activity of hydroalcoholic extract of *Ficuslyrataleaves* and Silymarin on % SGOT levels, %SGPT levels, % serum bilirubin levels, % ALP levels in Isoniazid induced hepatotoxicity in rats table 1, 2, 3, 4. Results of Hepatoprotective activity of hydroalcoholic extract of *Ficus elasticleaves* and Silymarin on % SGOT levels, %SGPT levels, % serum bilirubin levels, % ALP levels in Isoniazid induced hepatotoxicity in rats table 5, 6, 7, 8.

Table 2: Effect of Hydroalcoholic extract of *Ficuslyrataleaves* and Silymarin on % SGOT levels in Isoniazid induced hepatotoxicity in rats. [19]

Treatment	Dose	SGOT (%)
Normal	1 ml/kg, p.o.	162 ± 2.5
Isoniazid	100 mg/kg, p.o.	341.45 ± 6.5
<i>Ficuslyrataleaves</i>	100 mg/kg p.o.	214.0 ± 3.0 ^{***}
<i>Ficuslyrataleaves</i>	200 mg/kg p.o.	189.0 ± 3.5 ^{***}
Silymarin	2.5 mg/kg p.o.	168.0 ± 2.9 ^{***}

Table 3: Effect of Hydroalcoholic extract of *Ficuslyrataleaves* and Silymarin on %SGPT levels in Isoniazid induced hepatotoxicity in rats [19]

Treatment	Dose	SGPT (%)
Normal	1 ml/kg, p.o.	160.0 ± 2.0
Isoniazid	100 mg/kg, p.o.	321.0 ± 3.50
<i>Ficuslyrataleaves</i>	100 mg/kg p.o.	194.0 ± 4.0 ^{***}
<i>Ficuslyrataleaves</i>	200 mg/kg p.o.	179.0 ± 3.50 ^{***}
Silymarin	2.5 mg/kg p.o.	155.0 ± 3.50 ^{***}

Table 4:

Effect of

***Ficuslyrataleaves* and Silymarin on % serum bilirubin levels in Isoniazid induced hepatotoxicity in rats**

Treatment	Dose	Serum Bilirubin (%)
Normal	1 ml/kg, p.o.	120.0 ± 5.50
Isoniazid	100 mg/kg, p.o.	285.0 ± 3.50
<i>Ficuslyrataleaves</i>	100 mg/kg p.o.	159.0 ± 4.51 ^{***}
<i>Ficuslyrataleaves</i>	200 mg/kg p.o.	129.0 ± 1.60 ^{***}
Silymarin	2.5 mg/kg p.o.	115.0 ± 4.50 ^{***}

Table 5: Effect of *Ficuslyrataleaves* and Silymarin on % ALP levels in Isoniazid induced hepatotoxicity in rats

Treatment	Dose	ALP (%)
Normal	1 ml/kg, p.o.	162.0 ± 3.20
Isoniazid	100 mg/kg, p.o.	325.0 ± 5.57
<i>Ficuslyrataleaves</i>	100 mg/kg p.o.	219.0 ± 4.30 ^{***}
<i>Ficuslyrataleaves</i>	200 mg/kg p.o.	186.0 ± 3.78 ^{***}
Silymarin	2.5 mg/kg p.o.	170.0 ± 5.40 ^{***}

3.3 Hepatoprotective activity of *Ficus elasticleaves* extract Isoniazid induced hepatotoxicity

Table6: Effect of Hydroalcoholic extract of *Ficus elasticleaves* and Silymarin on % SGOT levels in Isoniazid induced hepatotoxicity in rats

Treatment	Dose	SGOT (%)
Normal	1 ml/kg, p.o.	160 ± 2.5
Isoniazid	100 mg/kg, p.o.	328.45 ± 6.5
<i>Ficus elasticleaves</i>	100 mg/kg p.o.	205.0 ± 3.5 ^{***}

<i>Ficus elasticleaves</i>	200 mg/kg p.o.	177.0 ± 3.5 ^{***}
Silymarin	2.5 mg/kg p.o.	165.0 ± 2.5 ^{***}

Table 7: Effect of Hydroalcoholic extract of *Ficus elasticleaves* and Silymarin on %SGPT levels in Isoniazid induced hepatotoxicity in rats

Treatment	Dose	SGPT (%)
Normal	1 ml/kg, p.o.	155.0 ± 2.0
Isoniazid	100 mg/kg, p.o.	321.0 ± 3.50
<i>Ficus elasticleaves</i>	100 mg/kg p.o.	188.0 ± 4.0 ^{***}
<i>Ficus elasticleaves</i>	200 mg/kg p.o.	165.0 ± 3.50 ^{***}
Silymarin	2.5 mg/kg p.o.	162.0 ± 3.50 ^{***}

Table 8: Effect of *Ficus elasticleaves* and Silymarin on % serum bilirubin levels in Isoniazid induced hepatotoxicity in rats

Treatment	Dose	Serum Bilirubin (%)
Normal	1 ml/kg, p.o.	120.0 ± 5.50
Isoniazid	100 mg/kg, p.o.	274.0 ± 3.50
<i>Ficus elasticleaves</i>	100 mg/kg p.o.	147.0 ± 4.81 ^{***}
<i>Ficus elasticleaves</i>	200 mg/kg p.o.	123.0 ± 1.50 ^{***}
Silymarin	2.5 mg/kg p.o.	113.0 ± 4.50 ^{***}

Table 9: Effect of *Ficus elastic* leaves and Silymarin on % ALP levels in Isoniazid induced hepatotoxicity in rats

Treatment	Dose	ALP (%)
Normal	1 ml/kg, p.o.	162.0 ± 3.20
Isoniazid	100 mg/kg, p.o.	340.0 ± 5.57
<i>Ficus elastic</i> leaves	100 mg/kg p.o.	215.0 ± 4.30 ^{***}
<i>Ficus elastic</i> leaves	200 mg/kg p.o.	178.0 ± 3.78 ^{***}
Silymarin	2.5 mg/kg p.o.	167.0 ± 5.40 ^{***}

3.4Hepatoprotective activity of *Ficuslyrata*leavesextractParacetamol induced hepatotoxicity model

Hepatoprotective activity of extract of *Ficuslyrata*leaves and Paracetamol on % SGOT levels, %SGPT levels, % serum bilirubin levels, % ALP levels in Paracetamol induced hepatotoxicity in rats. Table 9, 10,11, 12. results ofHepatoprotective activity of hydroalcoholic extract of *Ficus elastic*leaves and Silymarin on % SGOT levels in Paracetamol induced hepatotoxicity in rats. table 13, 14, 15, 16.

Table 10: Effect of Hydroalcoholic extract of *Ficuslyrata*leaves and Silymarin on % SGOT levels in Paracetamol induced hepatotoxicity in rats

Treatment	Dose	SGOT (%)
Normal	1 ml/kg, p.o.	158 ± 2.5
Paracetamol	50 mg/kg, p.o.	352.45 ± 6.5
<i>Ficuslyrata</i> leaves	100 mg/kg p.o.	227.0 ± 3.5 ^{***}
<i>Ficuslyrata</i> leaves	200 mg/kg p.o.	203.0 ± 3.0 ^{***}
Silymarin	2.5 mg/kg p.o.	179.0 ± 3.5 ^{***}

Table 11: Effect of Hydroalcoholic extract of *Ficuslyrataleaves* and Silymarin on %SGPT levels in Paracetamol induced hepatotoxicity in rats

Treatment	Dose	SGPT (%)
Normal	1 ml/kg, p.o.	151.0 ± 2.0
Paracetamol	50 mg/kg, p.o.	316.0 ± 3.50
<i>Ficuslyrataleaves</i>	100 mg/kg p.o.	189.0 ± 4.0 ^{***}
<i>Ficuslyrataleaves</i>	200 mg/kg p.o.	174.0 ± 3.50 ^{***}
Silymarin	2.5 mg/kg p.o.	152.0 ± 3.50 ^{***}

Table 12: Effect of *Ficuslyrataleaves* and Silymarin on % serum bilirubin levels in Paracetamol induced hepatotoxicity in rats

Treatment	Dose	Serum Bilirubin (%)
Normal	1 ml/kg, p.o.	122.0 ± 5.50
Paracetamol	50 mg/kg, p.o.	304.0 ± 3.50
<i>Ficuslyrataleaves</i>	100 mg/kg p.o.	201.0 ± 4.51 ^{***}
<i>Ficuslyrataleaves</i>	200 mg/kg p.o.	165.0 ± 1.60 ^{***}
Silymarin	2.5 mg/kg p.o.	125.0 ± 4.50 ^{***}

Table 13: Effect of *Ficuslyrataleaves* and Silymarin on % ALP levels in Paracetamol induced hepatotoxicity in rats

Treatment	Dose	ALP (%)
Normal	1 ml/kg, p.o.	155.0 ± 3.20
Paracetamol	50 mg/kg, p.o.	337.0 ± 5.57
<i>Ficuslyrataleaves</i>	100 mg/kg p.o.	232.0 ± 4.30 ^{***}

<i>Ficuslyrata</i> leaves	200 mg/kg p.o.	191.0 ± 3.78 ^{***}
Silymarin	2.5 mg/kg p.o.	175.0 ± 5.40 ^{***}

3.5 Hepatoprotective activity of *Ficus elasticleaves* extract Paracetamol induced hepatotoxicity model

Table 14: Effect of Hydroalcoholic extract of *Ficus elasticleaves* and Silymarin on % SGOT levels in Paracetamol induced hepatotoxicity in rats

Treatment	Dose	SGOT (%)
Normal	1 ml/kg, p.o.	165 ± 2.5
Paracetamol	50 mg/kg, p.o.	339.45 ± 6.5
<i>Ficus elasticleaves</i>	100 mg/kg p.o.	219.0 ± 3.5 ^{***}
<i>Ficus elasticleaves</i>	200 mg/kg p.o.	181.0 ± 3.5 ^{***}
Silymarin	2.5 mg/kg p.o.	168.0 ± 2.5 ^{***}

Table 15: Effect of Hydroalcoholic extract of *Ficus elasticleaves* and Silymarin on %SGPT levels in Paracetamol induced hepatotoxicity in rats

Treatment	Dose	SGPT (%)
Normal	1 ml/kg, p.o.	152.0 ± 2.0
Paracetamool	50 mg/kg, p.o.	314.0 ± 3.50
<i>Ficus elasticleaves</i>	100 mg/kg p.o.	196.0 ± 4.0 ^{***}
<i>Ficus elasticleaves</i>	200 mg/kg p.o.	174.0 ± 3.50 ^{***}
Silymarin	2.5 mg/kg p.o.	156.0 ± 3.50 ^{***}

Table 16: Effect of *Ficus elasticleaves* and Silymarin on % serum bilirubin levels in Paracetamol induced hepatotoxicity in rats

Treatment	Dose	Serum Bilirubin (%)
-----------	------	---------------------

Normal	1 ml/kg, p.o.	120.0 ± 5.50
Paracetamol	50 mg/kg, p.o.	290.0 ± 3.50
<i>Ficus elasticleaves</i>	100 mg/kg p.o.	155.0 ± 4.81 ^{***}
<i>Ficus elasticleaves</i>	200 mg/kg p.o.	131.0 ± 1.50 ^{***}
Silymarin	2.5 mg/kg p.o.	125.0 ± 4.50 ^{***}

Table 17: Effect of *Ficus elastic* leaves and Silymarin on % ALP levels in Paracetamol induced hepatotoxicity in rats

Treatment	Dose	ALP (%)
Normal	1 ml/kg, p.o.	162.0 ± 3.20
Paracetamol	50 mg/kg, p.o.	331.0 ± 5.57
<i>Ficus elasticleaves</i>	100 mg/kg p.o.	227.0 ± 4.30 ^{***}
<i>Ficus elasticleaves</i>	200 mg/kg p.o.	172.0 ± 3.78 ^{***}
Silymarin	2.5 mg/kg p.o.	165.0 ± 5.40 ^{***}

CONCLUSION

Hepatic damage is the main obstacle to drug development and a major factor in the reason why pharmaceutical companies have stopped selling their products. Liver ailment caused by medication might be either exceptional (low frequency and probably dosage related) or unsurprising (high rate and measurement related). Responses that are eccentric, often referred to as unusual, can be classified as non-immune or resistive to extreme touchiness. Since ancient times, people have turned to plants as a source for healing illnesses, providing long-lasting excellent health, relieving severe suffering, and adding flavor and fragrance to food. India and China have been using a human services framework centered on plants for more than 5,000 years. Up until about 50 years ago,

botanicals were used as common medications in Europe. Ancient advances from China, India, and Greece were independently incorporated into the pharmacological systems of the Arabic countries. However, the Ayurvedic framework is the most unparalleled in terms of study breadth and profundity. This framework, which is said to have started approximately 6000 BC, is one of the oldest prescription frameworks. In many countries, medicinal plants have become an integral element of traditional medical frameworks. Documentation of the examination work done on conventional medications is required. These investigations aid in the identification of plant material. When synthetic analgesics and antimicrobial drugs were not yet widely available in the middle of the 20th century, natural drugs dominated the therapeutic landscape. People began to switch to this framework when allopathic pharmaceutical arrangements for manufactured drugs gained rapid traction and faster symptomatic relief.

Ethical Approval

Animal Ethic committee approval has been collected and preserved by the author(s)

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCE

1. AnggunSelvyaArsyad, AriefNurrochmad, UniversitasGadjahMada, NanangFakhrudinPhytochemistry, traditional uses, and pharmacological activities of *Ficuselastica*Roxb. ex Hornem: A review, , Journal of Herbmed Pharmacology 12(1):41-53 December 2022. DOI:10.34172/jhp.2023.04
2. HandaSS.Futuretrendsofplantsasdrugs.PharmaTimes.1991;4:13-23.
3. Hassan AbdallaAlmahyl, MawardiRahmani%, MohdAspollahSukarP& Abdul Manaf Ali4, Investigation on the Chemical Constituents of the Leaves of *Ficuselastica*Roxb. and Their Antimicrobial Activity, *Pertanika J. Sci. & Technol.* 11(1): 57 - 63 (2003)
4. Jean Emmanuel MbossoTeinkela Xavier SiweNoundou Edwige Laure Nguemfo Franck Meyer Rene Wintjens Michelle Isaacs Albert Emmanuel MpondoMpondo Heinrich C. Hoppe Rui Werner Maçedo Krause and Anatole Guy Blaise AzebazeBiological activities of plant extracts from *Ficuselastica* and *Selaginellavogelli*: An antimalarial, antitrypanosomal and cytotoxicity evaluation. *Saudi J Biol Sci.* 2018 Jan; 25(1): 117–122. Published online 2017 Jul 20. doi: 10.1016/j.sjbs.2017.07.002
5. KatzungG.(2005),*BasicandClinicalPharmacology*, 6thedi., Appleton and Lange, Connecticut, Pg. 297-9.
6. Keppler D, Lesch R, Reutter W, Decker K. Experimental hepatitisinducedbyd-galactosamine.*ExpMolPathol.*1968Oct;9(2):279-90.
7. Khandelwal KR. Ed. *Practical Pharmacognosy Technique and Experiments*, 23rdEdn: 2005; 15.
8. Kokate CK. Ed. *Practical Pharmacognosy*, 4thEdn., *Vallabh Prakashan*: 1994; 112:120.
9. KulikovaO.(2015),*HepatoprotectiveActivityofPlantPeptides*,*Ame.J. ofPlantSci.*,6,848-55.
10. Levy SB, Marshall B. Antibacterial resistance worldwide: causes, challenges and responses. *Nature Medicine Supplement.* 2007; 10(12):122-129.
11. M. P. Gonthier, C. Remesy, A. Scalbert, V. Cheynier, J. M. Souquet, K. Poutanen, and A. M. Aura, “Microbial metabolism of caffeic acid and its esters chlorogenic and caftaric acids by human faecal microbiota *in vitro*,” *Biomedicine & Pharmacotherapy*, vol. 60, pp. 536-540, 2006.

12. McIntyre N, Benhamou JP, Bircher J, Rizzetto M, Rodes J. Oxfordtext book of clinical hepatology. Vol 2. 2nd ed. New York: OxfordUniversityPress;1991.p.907-09.
13. Mohamed Z. M. Salem¹ , A. Z. M. Salem² *, L. M. Camacho³ and Hayssam M. Ali
Antimicrobial activities and phytochemical composition of extracts of Ficus species: An over view, African Journal of Microbiology Research page no 4211-4219.
14. Mohd Mujeeb, Shah Alam Khan, Vidhu Aeri, and Babar Ali , Hepatoprotective Activity of the Ethanolic Extract of *Ficus carica* Linn. Leaves in Carbon Tetrachloride-Induced Hepatotoxicity in Rats, Iran J Pharm Res. 2011 Spring; 10(2): 301–306.
15. Nadeem M.etal.,(1996),“Hepatoprotective activity of some herbal formulations in India”. Indian Drugs,33(8):pp390-395.
16. OECD, Guideline for Testing of Chemicals-Acute Oral Toxicity-Acute Toxic Class Method. Paris: OECD; 2001.
17. Patel R.B.,J.D.Raval,(1998),“Hepatoprotective effect of Indian Medicinal Plants–Part I”.Indian Drugs,25(6),pp224-27.
18. Reddy KN, Subbaraju GV, Reddy CS, Raju VS. Ethnoveterinary medicine for treating livestock in Eastern Ghats of Andhra Pradesh. Indian Journal of Tradition Knowledge 2013; 5(3):368-378.
19. Deepak kumar, Mr. Dhyanendra Singh, Mrs. Pranali Mishra, Dr. Vishal Gupta pharmacological evaluation of Hepatoprotective activity of fruits Extract of *Syzygium aeoranthum* 2456-4184