

Original Research Article

Effect of scheduling of irrigation on growth, yield and water use efficiency of linseed (*Linum usitatissimum* L.)

ABSTRACT

A field experiment was conducted to study the effect of scheduling of irrigation on growth, yield and water use efficiency of linseed (*Linum usitatissimum* L.) at Main Agricultural Research Station (MARS), Raichur during *rabi* season of 2023-24. The experiment was laid out in randomized complete block design with five treatments, replicated four times. There were five treatments viz., T₁: Pre sowing irrigation only, T₂: Pre sowing irrigation *fb* one irrigation at vegetative stage (30-35 DAS), T₃: Pre sowing irrigation *fb* two irrigations at vegetative stage (30-35 DAS) and flowering stage (40-45 DAS), T₄: Pre sowing irrigation *fb* three irrigations at vegetative stage (30-35 DAS), flowering stage (40-45 DAS) and capsule development stage (60-65 DAS) and T₅: Rainfed condition. The results revealed that, significantly higher plant height (60.94 cm), number of branches plant⁻¹ (6.18), leaf area (36.94 cm² plant⁻¹), total dry matter production (16.41 g plant⁻¹), number of capsules plant⁻¹ (44.27), number of seeds (8.18 capsule⁻¹), seed weight (2.98 g plant⁻¹), seed yield (1106 kg ha⁻¹) and straw yield (2344 kg ha⁻¹) was registered with pre sowing irrigation *fb* three irrigations at vegetative stage (30-35 DAS), flowering stage (40-45 DAS) and capsule development stage (60-65 DAS) which was found to be on par with pre sowing irrigation *fb* two irrigations at vegetative stage (30-35 DAS) and flowering stage (40-45 DAS). Further, (T₄) recorded significantly lower water use efficiency (5.26 kg ha-mm⁻¹). Whereas, the treatment under rainfed condition (T₅) recorded significantly higher water use efficiency (13.59 kg ha-mm⁻¹) compared to all other treatments.

Keywords: Linseed, Irrigation, growth, Yield, Water use efficiency.

INTRODUCTION

Linseed (*Linum usitatissimum* L.) is a self-pollinated crop widely adapted to temperate climate of the world. It is also known as Agase in Kannada, Javas or Alashi in Marathi, Alsi in Hindi and Ousahalu in Telugu. It is an annual plant belongs to the genus *linum* of the family *Linaceae*. The name *Linum* is originated from the Celtic word 'lin' or 'thread' and the name *usitatissimum* is latin word meaning "most useful". It is believed that linseed, also called as flax is originated in the Middle East or Indian regions, one of the oldest crop plants cultivated in around 47 countries for the dual purpose of seed oil and fibre.

Linseed occupies an area of 32.23 lakh ha yielding 30.68 lakh tonnes with an average productivity of 952 kg ha⁻¹ in the world. Whereas in India, it occupies an area of 2.39 lakh ha with a production and productivity of about 1.67 lakh tonnes and 698 kg ha⁻¹, respectively. India holds fifth position in area and ranks sixth in production [2]. Though there has been slight improvement in average productivity over the previous years, but it is still far below than the potential yield (2000-2200 kg ha⁻¹) of improved linseed varieties in the major linseed growing nations such as Canada (1432 kg ha⁻¹), China (1308 kg ha⁻¹), USA (1258 kg ha⁻¹) and Kazakhstan (809 kg ha⁻¹) underlying the need for upscaling the production and productivity of this crop. The present status of linseed production could be increased to 2-3 folds through the adoption of improved varieties coupled with recommended production and protection technologies. In Karnataka, it is grown over an area of 26 twenty six thousand ha with a production of 25.27 thousand tonnes and productivity of 972 kg ha⁻¹ [3]. In Karnataka, it is mainly grown in northern districts viz., Raichur, Vijayapura, Kalaburagi, Bidar, Koppal, Yadagiri and Bellary during October to November under conserved soil moisture and limited nutrient conditions with poor management practices.

Linseed yields ~~seed(delete)~~ which is a rich source of both non-edible and edible oil. The industrial oil is an important ingredient in the manufacture of paint, varnish and linoleum [12]. Edible linseed oil is used for human consumption and contains α -linolenic acid (ALA), a polyunsaturated fatty acid that has nutritional and health benefits [17]. Linseed oil is found to be containing 5 major fatty acids, viz., palmitic (about 7 %), stearic (3.4-4.6 %), oleic (18.5-22 %), linoleic (14.2-17 %) and linolenic acid (51.9-55.2 %) [15]. Linseed is one of the richest sources of lignin (800 times more than any other plant seed except sesame seeds 47 times more) which provides protection against certain form of cancer due to estrogenic and anti-estrogenic activity in the body. The oil cake is a good feed for milch cattle and poultries and hence priced 50% higher than rapeseed-mustard cake. It is good in taste and contains 36% protein of which 85% is digestible. It is also used as organic manure. It contains about 5% N, 1.4% P₂O₅ and 1.8% K₂O. Linseed is globally cultivated for its fibres and is called flax. The stem yields good quality fibre having high strength and durability.

Among all, water is one of the most important critical inputs for agriculture which consumes more than 70 per cent of the water resources of the country. Availability of adequate quantity and quality of water is key factor for achieving higher productivity levels. Investments in conservation of water, improved techniques to ensure its timely supply and improve its efficient use are some of the imperatives which the country needs to enhance. Poor irrigation efficiency of conventional irrigation system has not only reduced the anticipated outcome of investments made towards water resource development, but has also resulted in environmental problems like water logging and soil salinity thereby affecting crop yields. Thus, it calls for massive investments in adoption of improved methods of irrigation such as drip and sprinkler including fertigation [1].

There are several reports of marked response of linseed to irrigation. Significant yield increase from irrigation between early November and January when total rainfall and its distribution were poor. The significant positive effect of irrigation on seed yield was attributed to higher capsule

number and more seeds/capsules under irrigation [10]. The main effect of water stress on linseed yield attributes is on the number of capsules and seeds/plant or unit area [7]. Severe water stress can reduce mean seed weight and consequently lower yield [8]. Water deficit lowers seed yield mainly by reducing PAR interception by shortening growth duration and affecting the canopy development [16]. Water stress accelerated leaf, stem and pod senescence in linseed and thus reduced capsule and seed growth [8]. Usually, it is cultivated in rainfed areas. If winter rains fail, it creates soil moisture stress. To overcome this supplemental irrigation is required. Hence scheduling of irrigation at different growth stages help to increase the growth and yield by reducing moistures stress. Efficient water management is important in getting higher yield with good quality produce, higher water use efficiency, increased soil productivity, higher fertilizer use efficiency and lesser irrigation cost which can be achieved by following optimum irrigation schedule.

MATERIAL AND METHODS

A field experiment was conducted during *rabi* season of 2023-24 at Main Agricultural Research Station (MARS), Raichur, Karnataka. The field where the experiment was conducted is located at a Latitude 16.15° N and Longitude 77.20° E with an altitude of 407 meters above the Mean Sea Level (MSL). The soil of the experimental field was sandy loam in texture with alkaline pH (8.39), bulk density (1.61 g cm⁻³), organic carbon content (0.66 %), available nitrogen (206.58 kg ha⁻¹), phosphorus (23.25 kg ha⁻¹) and potassium (236.98 kg ha⁻¹) contents. The experiment was laid out in randomized complete block design (CRBD) with five treatments and replicated four times. There were five treatments viz., T₁: Pre sowing irrigation only, T₂: Pre sowing irrigation *fb* one irrigation at vegetative stage (30-35 DAS), T₃: Pre sowing irrigation *fb* two irrigations at vegetative stage (30-35 DAS) and flowering stage (40-45 DAS), T₄: Pre sowing irrigation *fb* three irrigations at vegetative stage (30-35 DAS), flowering stage (40-45 DAS) and capsule development stage (60-65 DAS) and T₅: Rainfed condition. The linseed variety NL-115 having duration of 110-115 days was sown with spacing of 30 x 5 cm with a seed rate of 25 kg ha⁻¹. The recommended dose of fertilizer being 40:20:20 kg NPK ha⁻¹, the entire quantity of P₂O₅, K₂O and 50 per cent of nitrogen were applied as basal dose at the time of sowing in the form of Urea, DAP and Muriate of Potash, respectively. The remaining 50 per cent of nitrogen was top dressed at 25 days after sowing (DAS) (How the urea was applied in dry soil as first irrigation was given at 30-35 days after sowing ?) using urea as band application. Suitable plant protection measures were taken during the cropping season. The irrigation treatments were imposed as per the schedule. Five plants were randomly selected for taking observations on growth and yield attributing parameters as per the schedule. The crop in each net plot was harvested separately as per treatment and the values were converted into hectare basis and expressed in kilograms per hectare. The Water Use Efficiency of each treatment was computed using the following formula [14]

$$WUE = \frac{Y}{WR}$$

Formatted: Font color: Red

Formatted: Font color: Red

Where, WUE = Water use efficiency (kg ha-mm⁻¹)

Y = Crop yield (kg ha⁻¹)

WR = Total water requirement /used in the field (mm)

Comments : If the total water requirement has been considered here for calculation of WUE , then what about the soil moisture contribution for growth of crop ? . If soil moisture contribution has not considered, then it is irrigation water use efficiency and not total water use efficiency.

Formatted: Font color: Red

In case of rainfed treatment, author cannot calculate WUE as it was calculated based on irrigation water applied. If soil moisture contribution for crop ET was considered alongwith crop ET from irrigation water , then the above equation is ok and the WUE is designated based on crop ET and not based on total water use.

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

The experimental data obtained were subjected to statistical analysis adopting Fisher's method of analysis of variance as outlined by [5]. The level of significance used in "F" test was given at 5 per cent. Critical difference (CD) values are given in the table at 5 per cent level of significance, wherever the "F" test was found significant.

RESULTS AND DISCUSSION

Effect on crop growth parameters

The results of the experiment revealed that, among the treatments, at harvest significantly higher plant height, number of branches per plant, leaf area and total dry matter production (60.94 cm, 6.18, 36.94 cm² plant⁻¹ and 16.41 g plant⁻¹, respectively) were recorded with pre sowing irrigation *fb* three irrigations at vegetative stage (30-35 DAS), flowering stage (40-45 DAS) and capsule development stage (60-65 DAS) which was at par with pre sowing irrigation *fb* two irrigations at vegetative stage (30-35 DAS) and flowering stage (40-45 DAS) which recorded on par growth parameters (plant height, number of branches per plant, leaf area and total dry matter production (58.39 cm, 5.93, 34.71 cm² plant⁻¹ and 15.11 g plant⁻¹, respectively). Whereas, rainfed condition recorded significantly lower plant height (41.52 cm), number of branches (4.28), leaf area (24.30 cm² plant⁻¹) and total dry matter production (10.90 g plant⁻¹) (Table 1). This is mainly due to the plants under ambient and sufficient moisture conditions will utilise other growth resources and accelerate the enzymatic activity, photosynthesis, carbohydrate metabolism, protein synthesis and cell division which in turn enhanced growth and development of plant owing to right amount of water at right time. Similar results were also obtained by many workers [4] and [11].

Table 1: Growth parameters of linseed at harvest as influenced by scheduling of irrigation

Treatment	Plant height (cm)	Number of branches plant ⁻¹	Leaf area (cm ² plant ⁻¹)	Total dry matter production (g plant ⁻¹)
T ₁	42.39	4.46	25.11	11.50

T₂	45.95	4.69	28.89	12.32
T₃	58.39	5.93	34.71	15.11
T₄	60.94	6.18	36.94	16.41
T₅	41.52	4.28	24.30	10.90
S. Em. ±	2.17	0.47	1.57	0.32
C.D. (P = 0.05)	6.69	1.44	4.83	0.99
CV %				

Formatted: Font color: Red


Effect on yield parameters and yield

Among the treatments, significantly higher yield attributes viz., number of capsules plant⁻¹ (44.27), number of seeds capsule⁻¹ (8.18), seed weight plant⁻¹ (2.98 g) and 1000 seed weight (8.16 g) were recorded with irrigation scheduled at pre sowing irrigation *fb* three irrigations at vegetative stage (30-35 DAS), flowering stage (40-45 DAS) and capsule development stage (60-65 DAS) that resulted in higher yield attributing parameters and yield levels as compared to other treatments. Whereas, number of capsules plant⁻¹ (41.53) and seeds capsule⁻¹ (7.75), seed weight plant⁻¹ (2.71 g) and 1000 seed weight (7.99 g) were on par with irrigation scheduled at pre sowing irrigation *fb* two irrigations at vegetative stage (30-35 DAS) and flowering stage (40-45 DAS). This might be due to improved growth parameters that in turn lead to better translocation of photosynthates from source to sink which helped for attaining higher yield attributing parameters and yield. However, 1000 seed weight of linseed was non-significant with irrigation scheduling in linseed. From the result it was found that, application of pre sowing irrigation *fb* three irrigations at vegetative stage (30-35 DAS), flowering stage (40-45 DAS) and capsule development stage (60-65 DAS) resulted in higher seed yield (1106 kg ha⁻¹) and straw yield (2344 kg ha⁻¹) and this was found to be on par with pre sowing irrigation *fb* two irrigations at vegetative stage (30-35 DAS) and flowering stage (40-45 DAS) with seed (995 kg ha⁻¹) and straw yield (2188 kg ha⁻¹), respectively. Rainfed condition registered significantly lower seed yield (477 kg ha⁻¹) and straw yield (1287 kg ha⁻¹) (Table 2).

The higher seed and straw yields are typically the result of combination of optimal soil and water management, use of high-yielding crop varieties, effective pest and disease control, favourable climatic conditions and good agronomic practices. Implementing these factors collectively can lead to significant improvements in crop yield and overall productivity. Adequate and well-managed soil moisture ensures that plants have enough moisture throughout their growing period, which supports robust growth and high yield. Proper irrigation scheduling can prevent drought / stress and water logging, leading to better crop yields. Soils rich in essential nutrients (nitrogen, phosphorus, potassium, etc.) support strong crop growth and development and higher yield. Proper fertilization practices can enhance nutrient availability. Similar observations were recorded by previous workers

water use effective by considering the above values is totally wrong.), viz., pre sowing irrigation only, pre sowing irrigation fb one irrigation at vegetative stage (30-35 DAS), pre sowing irrigation fb two irrigations at vegetative stage (30-35 DAS) and flowering stage (40-45 DAS), pre sowing irrigation fb three irrigations at vegetative stage (30-35 DAS), flowering stage (40-45 DAS) and capsule development stage (60-65 DAS) and rainfed condition, respectively. Water use efficiency differed significantly due to different scheduling of irrigation at different growth stages of linseed. Water use efficiency can be presented in terms of yield realized per hectare meter of water used. Among all the treatments pre sowing irrigation fb three irrigations at vegetative stage (30-35 DAS), flowering stage (40-45 DAS) and capsule development stage (60-65 DAS) recorded significantly lower water use efficiency of 5.26 kg ha-mm⁻¹. Water use efficiency shows the decreasing trend with increase in amount of water used. The rainfed condition treatment recorded significantly higher water use efficiency of 13.59 kg ha-mm⁻¹ over all other irrigation treatments (Table 3 and Fig. 1).

Fig. 1: Total Water Used and Water Use Efficiency of linseed as influenced by scheduling of irrigation



Excessive irrigation may lead to reduced efficiency, potentially due to issues like waterlogging or less effective water management practices. Water management emphasize the need for carefully managing irrigation schedules to avoid overuse and to optimize water use. Effective water management strategies could balance water application with crop needs to maximize yield per unit of water used. While irrigation is necessary for crop growth, excessive water application can diminish efficiency. Linseed appears to be highly efficient under limited water conditions, suggesting that

targeted irrigation practices and water conservation can lead to overall better water use efficiency. The results are in conformity with findings of [13].

CONCLUSION

From the above study it can be concluded that, application of irrigation water at optimum quantity is essential to improve the crop yields. If ample amount of irrigation water is available, then scheduling of irrigation at pre sowing irrigation *fb* three irrigations at vegetative stage (30-35 DAS), flowering stage (40-45 DAS) and capsule development stage (60-65 DAS) helps in getting higher yield. If water is limited, then irrigation scheduling at pre sowing irrigation *fb* two irrigations at vegetative stage (30-35 DAS) and flowering stage (40-45 DAS) is essential for attaining higher yield. Water use efficiency shows the decreasing trend with increase in amount of water utilised. Rainfed condition recorded significantly higher water use efficiency. Whereas, pre sowing irrigation *fb* three irrigations at vegetative stage (30-35 DAS), flowering stage (40-45 DAS) and capsule development stage (60-65 DAS) recorded significantly lower water use efficiency.

REFERENCES

1. Anonymous, 2017, Annual report of water for sustainable food and agriculture. Food and Agriculture Organization of the United Nations, Rome.
2. Anonymous, 2022, Agricultural statistics at a glance. <https://eands.dacnet.nic.in>.
3. Anonymus, 2022a, Area, production and productivity of linseed in India and Karnataka, Indiaagristat, www.indiastat.com.
4. Ebied, M. and Badawi, M. I., 2023, Effect of different irrigation treatments on yield, quality and water use efficiency for some flax (*Linum usitatissimum* L.) varieties. *Menoufia J. Plant. Prod.*, 8(11): 211-227.
5. Gomez, K. A. and Gomez, A. A., 1984, Statistical Procedure for Agriculture Research, 2nd Ed., John Willey and Sons, New York, p. 680.
6. Gurjar, R., Chimote, A. and Raut, G. L. V., 2017, Impact of irrigation and nutrient management on growth and yield of flaxseed (*Linum usitatissimum* L.). *Trends Biosci.*, 10(27): 5624-5626.
7. Hocking, P. J. and Pinkerton, A., 1991, Response of growth and yield components of linseed to the onset or relief of nitrogen stress at several stages of crop development. *Field Crops Res.*, 27(1-2): 83-102.
8. Hocking, P. J., 1995, Effect of nitrogen supply on the growth, yield components and distribution of nitrogen in linola. *J. Plant Nutr.*, 18(2): 257-275.
9. Kashyap, T. L., Khajani, S. N., Thakur, C. L. and Lakpale, R., 2017, Effect of different tillage practices and irrigation schedule on the growth and yield of linseed in *alfisols* of Chhattisgarh plains. *J. Plant. Dev. Sci.* 9(11): 1039-1044.

10. Lisson, S. N. and Mendham, N. J. 2000, Agronomic studies of flax in a south-eastern Australia. *Australian J. Exp. Agric.*, 40: 1101-1112.
11. Marbate R. S., Uparkar A. I., Choudhari, R. J. and Paga, P. C., 2020, Growth attributes, yield and soil moisture studies of linseed as influenced by different land configuration treatments and irrigation levels. *Int. J. Chem. Stud.*, 8(4): 3593-3596.
12. Matheson, E. M., 1976, Linseed. Vegetable oil Seed crops in Australia, Sydney: *Holt, Rinehart and Winston*, Sydney, 111-121.
13. Sharma, J. C., Tomar, S. S., Shivran, R. K. and Prakash, C., 2012, Water requirement, water use efficiency, consumptive use, yield and quality parameters of linseed (*Linum usitatissimum* L.) varieties as influenced by fertility levels and irrigation scheduling. *Adv. Life Sci.*, 1(2): 180-182.
14. Thanki, R. B., R. M., Solanki1, J. M., Modhavadia, B. S., Gohil. and P. J., Prajapati., 2014, Effect of irrigation scheduling at critical growth stages and fertility levels on growth, yield and quality of summer sesame (*Sesamum indicum* L.) *J. Oilseeds Res.*, 31(1): 41-45.
15. Vaisey-Genser, M. and H. Morris., 2016, Introduction: history of the cultivation and uses of flaxseed. 1-21.
16. Wilson, D. R., 1987, New approaches to understanding the growth and yield of pea crops. In W. A. Jermyn and G. S. Wratt (eds). Peas: Management for Quality. Agronomy Society of New Zealand. *Spl. Publication* 6: 23-28.
17. Wood, I. M., 1997, Fibre crops - New opportunities for Australian agriculture. Department of primary industries, Brisbane. pp.18-24.