

# First global host report and new distribution record of *Bactrocera penecorrecta* Drew (Diptera: Tephritidae) from Bihar, India

---

## ABSTRACT

Fruit fly, *Bactrocera penecorrecta* Drew (Diptera: Tephritidae) is here reported to reared from infested brinjal fruit (*Solanum melongena* Linnaeus) from Bihar, India. We report here brinjal as the first global host record of *B. penecorrecta*. Also the fruit fly *B. penecorrecta* reported for the first time from Bihar State and North India. Earlier report of *B. penecorrecta* is from Kerala, South India. In the present investigation, *B. penecorrecta* recorded after 22 years of its discovery from Kerala, South India during 2002 by R. A. I. Drew and S. Raghu. The record of brinjal as first host of *Bactrocera penecorrecta* warrant indepth investigation of level of infestation and sanitary and phyto-sanitary protocol development for the export as many recent invasion by different *Bactrocera* species in various part of the world beyond their native distributional records.

*Keywords: Fruit fly, Bactrocera, Tephritidae, Brinjal, India, Host Plant*

## 1. INTRODUCTION

The fruit fly, *Bactrocera penecorrecta*, first described by R. A. I. Drew [1] in the year 2002 and reported by Drew and Raghu 2002 from New Amarambalam Forest, Kerala, India. *B. penecorrecta* belongs to the tribe Dacini of family Tephritidae. The tribe Dacini is a highly diverse tribe within family Tephritidae and contains around 1000 species, a fifth of all known species in the family [2,3,4,5]. All tephritid fruit flies fall under the tribe Dacini are frugivorous or florivorous and about 10% of the 932 currently recognized species are economic pests of fruits and vegetables around the world [4,5,6,7,8,9,10,11]. These flies are also internationally significant pests of fruits and vegetables as adults lay their eggs into sound fruit on-plant, where the subsequent larval feeding causes fruit loss [11,12]. Knowledge of potential host plants of fruit flies is important in areas where fruit crops are grown commercially because damage to these crops may be caused by fruit fly migration from nearby host plants [13].


## 2. MATERIAL AND METHODS

The collection of fruit flies infested fruits of Brinjal (only two fruits with maggots were found in the field) from the Dumraon area, Buxar, Bihar, India (Latitude: 25.56° N and Longitude 84.14° E) carried out during May 2024 and brought to the laboratory of Department of Entomology, Veer Kunwar Singh College of Agriculture, Dumraon, Bihar India. The infested

fruits were kept in a transparent plastic jar of 1000 cc capacity with 5 cm layer of sawdust in the base of the jar for pupation of the matured third instar maggots. After ten days of fruit collection, one one adult emerged from the pupa. The emerged adult fly was preserved in 96% ethyl alcohol for identification. The identification of the adult specimen of fruit fly was carried out with the published key and description available in Drew and Raghu [1], Drew and Romig [14, 15]. For the photography of the specimen, fruit fly were pinned and used for photography (Make: Panasonic, Model: DMC-TZ80 under macro setting). The specimen is preserved in the Insect Museum, Veer Kunwar Singh College of Agriculture, Dumraon, Bihar, India, (Cat. No.: FFBPc-01)

### 3. RESULTS AND DISCUSSION

On the basis of morphological taxonomic keys and description available in Drew and Raghu [1] and Drew and Romig [14, 15], the fruit fly species reared from infested brinjal was identified as *Bactrocera penecorrecta*. The photographs of different part of *B. penecorrecta* are presented in Figure 1. The fruit fly *B. penecorrecta* has first described by Drew in 2002 [1].



**Figure1. Photograph of different parts of *B. penecorrecta* reard from Brinjal**

Till date the host plant of *B. penecorrecta* is unknown and has not been reported in published literature. We report here the brinjal as the first global host record of *B. penecorrecta* from Dumraon, Bihar, India. Also the earlier geographical distribution record of *B. penecorrecta* is limited to South India Drew & Raghu [1]; Drew & Romig [14]. In the present investigation, *B. penecorrecta* reported from Dumraon, Bihar (a north and east state of India) is a new geographical distribution record for the species. The present location of collection of *B. penecorrecta* is about 2000 km far away from its first site of collection from New Amarambalam forest, Kerala, India. Other tephritids recorded from brinjal and other solanaceous plant species in India are *Bactrocera latifrons* (Hendel), *Bactrocera prabhakari* Maneesh et al. and *Zeugodacus tau* (Walker) [16, 17, 18]. More information on biology, other

host plants and level of infestation needs to be generated for further enhancement of the knowledge about *B. penecorrecta*. Also as other tephritid fruit flies are invasive and economic pest of many fruit and vegetable crops, close watch is warranted for the entire state for any economic losses from the species along with the development of sanitary and phyto-sanitary protocol development for the export as many recent invasion by different *Bactrocera* species in various part of the world beyond their native distributional records.

## ETHICAL APPROVAL

This study does not contain any studies with human participants or large animals performed by any of the authors. No approval of research ethics committees was required to accomplish the goals of this study.

## REFERENCES

- [1] Drew RAI, Raghu S. The fruit fly fauna (Diptera: Tephritidae: Dacinae) of the rainforest habitat of the Western Ghats, India. Raffles B Zool. 2002; 50(2):327-52. <https://www.cabidigitallibrary.org/doi/book/10.1079/9781780640358.0000>
- [2] Norrbom AL, Carroll LE, Thompson FC, White IM, Freidberg A. Systematic Database of Names, In: F C, Thompson (Ed.) Fruit fly expert identification system and systematic information database. Myia 1999; 9: 65–51.
- [3] Pape T, Blagoderov V, Mostovski MB. Order DIPTERA Linnaeus1758. Zootaxa. 2011; 3148:222–29. <https://www.mapress.com/zootaxa/2011/f/zt03148p229.pdf>
- [4] Freidberg A, Kovac D, Shiao S. A revision of Ichneumonopsis Hardy 1973. (Diptera: Tephritidae: Dacinae: Gastrozonini ) Oriental bamboo-shoot fruit flies. Eur J Taxon. 2017; 317: 1–23. <https://doi.org/10.5852/ejt.2017.317>
- [5] Doorenweerd C, Leblanc L, Norrbom AL, Jose MS, Rubinoff D. A global checklist of the 932 fruit fly species in the tribe Dacini (Diptera, Tephritidae). ZooKeys. 2018;730:19–56. <https://doi.org/10.3897/zookeys.730.21786>
- [6] Fletcher B. The biology of Dacine fruit flies. Annu Rev Entomol. 1987; 32:115-44. <https://doi.org/10.1146/annurev.ento.32.1.115>
- [7] White IM, Elson-Harris MM. Fruit Flies of Economic Significance. Wallingford: CABI; 1992.
- [8] Clarke AR, Armstrong KF, Carmichael AE, Milne JR, Raghu, S, Roderick, GK Yeates, DK. Invasive phytophagous pests arising through a recent tropical evolutionary radiation: the *Bactrocera dorsalis* complex of fruit flies. Annu Rev Entomol. 2005;50:293–19. <https://doi.org/10.1146/annurev.ento.50.071803.130428>
- [9] Prabhakar CS. Biodiversity of fruit flies (Tephritidae: Diptera) and utilization of gut bacteria in their management. Ph.D. thesis, CSK Himachal Pradesh Agricultural University,

- [10] DeMeyer M, Delatte H, Mwatawala M, Quilici S, Vayssières JF, Virgilio M. A review of the current knowledge on *Zeugodacus cucurbitae* (Coquillett) (Diptera, Tephritidae) in Africa, with a list of species included in *Zeugodacus*. *ZooKeys*. 2015;540:539–57. + Supplementary material 1:(4). <https://doi.org/10.3897/zookeys.540.9672>
- [11] Vargas RI, Piñero JC, Leblanc L. An overview of pest species of *Bactrocera* fruit flies (Diptera: Tephritidae) and the integration of biopesticides with other biological approaches for their management with a focus on the Pacific region. *Insects*. 2015; 6:297–18. <https://pubmed.ncbi.nlm.nih.gov/26463186/>
- [12] Ekesi S, DeMeyer M, Mohamed SA, Virgilio M, Borgemeister C. Taxonomy ecology and management of native and exotic fruit fly species in Africa. *Annu Rev Entomol*. 2016; 61: 219-38. <https://doi.org/10.1146/annurev-ento-010715-023603>
- [13] Souza SAS, Resende ALS, Strikis PC, Costa JR, Ricci MSF, Aguiar-Menezes EL. Infestação ao natural de moscas frugívoras (Diptera: tephritoidea) em café arábica, sob cultivo orgânico e arborizado e a pleno sol, em Valença. RJ. *Neotrop Entomol*. 2005; 34:639-48. Spanish. <https://www.scielo.br/j/ne/a/rXTvS9CtCRNJTwjNdSCf3JD/?lang=pt>
- [14] Drew RAI, Romig MC. Tropical fruit flies of South-East Asia. Wallingford: CABI; 2013. <https://www.cabidigitallibrary.org/doi/book/10.1079/9781780644196.0000>
- [15] Drew RAI, Romig MC. Keys to the Tropical Fruit Flies of South-East Asia. Wallingford: CABI; 2016. <https://www.cabidigitallibrary.org/doi/book/10.1079/9781780644196.0000>
- [16] Akhila MU, Jiji T. Record of solanum fruit fly *Bactrocera latifrons* (Hendel) (Diptera: Tephritidae) infesting solanaceous vegetables in Kerala. *Curr Biotica*. 2015;9(1):78-81. <https://www.cabidigitallibrary.org/doi/pdf/10.5555/20153371510>
- [17] Kumar S, Agarwal ML, Bharati RC. Studies on the infestation of fruit fly, *Bactrocera latifrons* (Hendel) in berries of *Solanum indicum* Willd. *RAU J Res*. 2006; 16:66-69.
- [18] Singh MP, Gupta D, Hancock DL, Sharma I, Sharma PI. A new species of *Bactrocera* Macquart reared from *Solanum khasianum* Clarke in northern India and a new synonym. *Zootaxa*, 2023; 5380(6):526–40. <https://doi.org/10.11646/zootaxa.5380.6.2>