

Impact of intercropping systems on the management of defoliators in Groundnut (*Arachis hypogaea* L.)

Formatted: Font: Italic

Abstract: Groundnut (*Arachis hypogaea* L.), a crucial oilseed and protein crop, encounters substantial yield and quality losses due to insect pests. This study explores the efficacy of various intercrop systems in managing major insect pests, particularly ~~defoliators-defoliator~~ populations. ~~Study Conducted~~conducted during the ~~kharif seasons of~~ 2022-23 at the Main Agricultural Research Station, University of Agricultural Sciences, Dharwad, ~~the~~The research employed a randomized block design (RBD) to evaluate different intercrop combinations. Results indicate that among the various intercrops studied, the combination of groundnut + pearl millet was found to be effective in suppressing the *Spodoptera litura* ~~F.~~ by recording the lowest average larval population (0.76 larvae/~~m²~~meter row length), followed by groundnut + sorghum (1.17 larvae/~~m²~~meter row length) and the same combination *i.e* groundnut + pearl millet was also proved to be effective in minimizing the leaf miner incidence too (1.49 larvae/plant) and had least percent of leaf damage ~~where-as~~whereas groundnut intercropped with cluster beans found to be least effective and was found on par with sole groundnut. The combination of these millets in groundnut ~~crop-crops providing-provides~~ sustainable ~~Cancer-control of pest-management solutions~~solutions population in ~~for~~groundnut cultivation.

Formatted: Font: Italic

Formatted: Font: Italic

Introduction:

Groundnut is a major player in India's edible oil sector, ranking third in vegetable protein sources and fourth in oilseed crops. Its seeds contain 43-55% oil and 25-28% protein, along with vitamins E, K, and B (Blummel et al., 2005). The oil, rich in oleic (75-80%) and linoleic acids, has 5.6 calories per gram (Padgham et al., 1990). The oil cake is a high-protein animal feed, and the haulm provides quality fodder. In India, groundnut is grown on 4.73 million hectares, producing 6.72 million tonnes with a productivity of 1,422 kg/ha. Major producing states Gujarat, Andhra Pradesh, Tamil Nadu, Karnataka, and Maharashtra account for 80% of the area and 84% of production. Karnataka contributes 0.51 million hectares and 0.39 million tonnes with a productivity of 759 kg/ha (Anon., 2019).

Productivity is affected by biotic and abiotic stresses, and insect pests, which cause losses of Rs. 238 crores (Dutta and Bandopadhyay, 2020). Over 100 insect species damage groundnut, with major pests including the red hairy caterpillar (*Amsacta albistriga* Walker), tobacco caterpillar (*Spodoptera litura* Fabricius), gram caterpillar (*Helicoverpa armigera* Hübner), leaf miner (*Aproaerema modicella* Deventer), and sucking pests such as thrips (*Scirtothrips dorsalis* Hood), aphids (*Aphis craccivora* Koch), and leafhoppers (*Empoasca kerri* Pruthi) (Amin and Mohammad, 1980, Dodiya et al., 2023). Yield declines are due to low inputs, non-adoption of high-yield varieties, rainfed cultivation, pest and disease issues, and extreme climate conditions. Chemical insecticides, while effective, are costly and ecologically harmful. This study investigates the use of intercrops to manage defoliators in groundnut, aiming to develop cost-effective, environmentally friendly, and sustainable pest management strategies.

Materials and methods

The present investigation on the impact of various intercrops on the incidence of major insect pests and natural enemies in groundnut was carried out during *kharif* 2022-23 and 2023-24 at Main Agricultural Research Station, University of Agricultural Sciences Dharwad. The experiment was laid out in a randomized block design (RBD) with three replications having a plot size of 5.0 x 3.0 m leaving a gang way of one meter around the treatment plots. The groundnut variety, Dh-256 was sown at a spacing of 30 x 10 cm by following the recommended package of practices along with various intercrops at ratios specified in treatment details. All the agronomic practices were adopted and no plant protection measures were taken throughout the season. The data obtained was analyzed by following Analysis of Variance (ANOVA) and the treatments were differentiated by using the Duncan Multiple Range Test (DMRT).

List 1. Treatment details

Sl.No	Intercrop combination	Variety or hybrid of intercrops	Intercropping ratio
T1	Groundnut + Pearl millet	Kaveri Super Boss	4:1
T2	Groundnut + Maize	CP-818	4:1
T3	Groundnut + Foxtail millet	Dhft109-3	4:1
T4	Groundnut + Sorghum	CSH-30	4:1
T5	Groundnut + Cowpea	DC-15	4:1
T6	Groundnut + Clusterbean	Pusa Navabahar	4:1
T7	Groundnut + Sunflower	RSFH1887	4:1
T8	Groundnut sole crop	-	-

Observations on the incidence of major defoliators like groundnut leaf miner (No. of mines per plant), *S. litura* (No. of larvae per meter row length), and ~~per-cent~~ leaf damage ~~was-were~~ calculated by using the following formula:

$$\text{Per-cent} \text{ Percent of leaf damage} = \frac{\text{Number of damaged leaflets per plant}}{\text{Total number of leaves per plant}} \times 100$$

The data on major natural enemies like coccinellids (No./plant) and spiders (No./ plant) were recorded from five randomly selected plants from each treatment at weekly ~~interval~~ intervals from 25 to 90 days after sowing.

Results and Discussion

During the kharif season of 2022-23, the population of *Spodoptera litura* larvae was monitored weekly from 25 days after sowing (DAS) through eight weeks. The incidence of *S. litura* was observed from 30 to 75 DAS under various intercropping systems. Notably, no larvae were detected in the groundnut + pearl millet intercrop ~~at-in~~ the first week, whereas the groundnut + foxtail millet, groundnut + sorghum, and groundnut + cowpea intercrops each had 0.33 larvae per meter row length (mrl). These systems were significantly better compared to the groundnut + sunflower (0.97 larvae/mrl) and groundnut + cluster beans (1.00 larvae/mrl), with sole groundnut crops having the highest larval population of 1.33 larvae/mrl.

The larval population generally increased from the second to the fifth week, peaking in the fourth week before declining to its lowest by the eighth week. ~~At-In~~ the second week, the lowest larval population was recorded in the groundnut + pearl millet intercrop (0.33 larvae/mrl), which was statistically similar to groundnut + sorghum and groundnut + cowpea (0.67 larvae/mrl each), and significantly lower than other treatments. The sole groundnut crop had the highest larval population at 2.00 larvae/mrl. By the third week, the groundnut + pearl millet intercrop had the least larvae (1.00 larvae/mrl), comparable to groundnut + sorghum and groundnut + maize (1.33 larvae/mrl). Other treatments, including groundnut + cowpea (1.67 larvae/mrl), showed moderate effectiveness but were still significantly lower than the sole groundnut crop (3.33 larvae/mrl).

In the fourth and fifth weeks, the peak incidence of *S. litura* larvae was observed. Groundnut + pearl millet had the lowest larval counts (1.33 larvae/mrl and 1.39 larvae/mrl, respectively) and was statistically similar to groundnut + maize and groundnut + sorghum. Sole groundnut crops recorded the highest larval populations (3.67 larvae/mrl and 4.00 larvae/mrl) during these weeks. Groundnut + cluster bean, groundnut + sunflower, and groundnut + cowpea also had moderately high larval populations. By the sixth to eighth weeks, the larval population decreased, with groundnut + cowpea, groundnut + sunflower, and groundnut + cluster bean showing 1.00, 1.00, and 1.33 larvae/mrl, respectively, compared to the sole groundnut crop, which had 2.67 larvae/mrl.

Overall, the groundnut + bajra intercrop recorded the lowest average larval population (0.76 larvae/mrl), followed by groundnut + sorghum (1.17 larvae/mrl). Other intercropping systems managed *S. litura* populations within the range of 1.42 to 2.04 larvae/mrl, with the untreated control showing the highest larval count (2.95 larvae/mrl). These results are consistent with Girija

et al. (2015) and Agasimani et al. (1993), who noted reduced *S. litura* incidence in intercrop systems like groundnut + jowar. Nath and Singh (1998) and Rashmi et al. (2011) also observed reduced pest populations in groundnut intercropped with bajra and foxtail millet.

Leaf miner incidence was zero in all treatments at 25 DAS but increased by the second week. Groundnut + foxtail millet (1.52) and groundnut + pearl millet (1.49) had the lowest populations, statistically similar to each other. Other treatments had moderate leaf miner numbers, with groundnut + sunflower showing the highest count (2.34 larvae/plant), similar to the sole crop. The percent reduction in leaf miner population was highest in groundnut + foxtail millet and groundnut + pearl millet (44.00%), followed by groundnut + sorghum and groundnut + maize (38.00%). Groundnut as a sole crop had a mean larval load of 1.70 per plant. Phytochemicals in millets and cowpea may have influenced oviposition by leaf miner moths, leading to lower populations in these intercrops. Intercropping might also create a microclimate that benefits [the](#) natural enemies of the leaf miner.

Leaf damage due to defoliators was significantly lower in groundnut intercropped with bajra and foxtail millet (12.29% and 12.50%, respectively) compared to the sole groundnut crop, which had the highest damage (16.86%). Groundnut + maize and groundnut + sorghum also showed lower damage (13.93% and 16.00%, respectively). Other intercropping systems, including groundnut + cowpea and groundnut + sunflower, had moderate levels of damage, with groundnut + cluster bean showing the highest damage (27.87%), comparable to the sole crop (30.55%). The reduced leaf damage in certain intercropping systems supports findings from Girija et al. (2015) and Rashmi et al. (2011) regarding lower damage in groundnut + foxtail millet systems. Dodiya et al. (2024) reported the incidence of *S. litura* in field 29th to 44th SMW. Rango Rao et al. (1995) and Patil (1993) similarly reported reduced damage in groundnut + sunflower and groundnut + sorghum intercrops. Nath and Singh (1998) also observed lower damage in groundnut + bajra systems, aligning with this study's results.

Table 1. Effect of intercropping on population of *Spodoptera litua* during *kharif* 2022-23 at Dharwad.

Formatted: Font: Italic

Treatments		No. of larvae /meter row length								Cumulative mean
		1 week	2 weeks	3 weeks	4 weeks	5 weeks	6 weeks	7 weeks	8 weeks	
T1	Groundnut + Pearl millet	0 (0.71) c	0.33 (0.9) c	1 (1.22) c	1.33 (1.34) c	1.39 (1.37) c	1 (1.46) c	0.67 (1.07) d	0.33 (0.9) d	0.76
T2	Groundnut + Maize	1.33 (1.34) a	1 (1.22) b	1.33 (1.34) c	2 (1.56) bc	2.33 (1.68) bc	1.67 (1.58) bc	1 (1.22) cd	0.67 (1.07) cd	1.42
T3	Groundnut + Foxtail millet	0.33 (0.9) bc	1 (1.22) b	2.33 (1.68) ab	2 (1.58) bc	2.67 (1.77) ab	2 (1.34) b	0.67 (1.07) d	0.67 (1.07) cd	1.46
T4	Groundnut + Sorghum	0.33 (0.9) bc	0.67 (1.07) bc	1.33 (1.34) c	2 (1.56) bc	2 (1.58) bc	1.33 (1.46) c	1 (1.22) cd	0.67 (1.07) cd	1.17
T5	Groundnut + Cowpea	1 (1.22) a	0.67 (1.05) bc	1.67 (1.46) bc	3 (1.87) ab	2.33 (1.68) bc	1.67 (1.66) bc	1.83 (1.51) bc	1 (1.22) bc	1.65
T6	Groundnut + Cluster bean	1 (1.22) a	1.33 (1.34) ab	2.33 (1.68) ab	2.67 (1.77) ab	2.67 (1.74) ab	2.33 (1.56) c	2.67 (1.77) ab	1.33 (1.34) b	2.04
T7	Groundnut + Sunflower	0.97 (1.21) a	1.33 (1.34) ab	2.67 (1.77) ab	2.33 (1.68) abc	2.5 (1.73) bc	2 (2.04) b	1.33 (1.34) cd	1 (1.22) bc	1.77
T8	Groundnut sole crop	1.33 (1.34) a	2 (1.58) a	3.33 (1.95) a	3.63 (2.03) a	4 (2.12) a	3.67 (4.02) a	3 (1.86) a	2.67 (1.77) a	2.95
	S.Em. ±	0.08	0.1	0.1	0.12	0.13	0.14	0.1	0.07	
	C.D. (5%)	0.24	0.30	0.31	0.36	0.38	0.42	0.32	0.22	
	C. V. (%)	12.24	13.99	11.45	12.21	12.77	12.76	13.07	10.55	

Table 2. Influence of intercropping on leaf damage in groundnut by defoliators during kharif 2022-23.

Treatments		Per cent leaf damage at								Cumulative mean
		1 week	2 weeks	3 weeks	4 weeks	5 weeks	6 weeks	7 weeks	8 weeks	
T1	Groundnut + Pearl millet	12.29 c (20.52)	16.89 b (24.26)	18 c (25.1)	22.67 c (28.36)	25.14 c (30.06)	26.94 b (31.24)	26.94 de (31.24)	22.95 bc (28.62)	21.48
T2	Groundnut + Maize	13.93 abc (21.9)	16.71 b (24.12)	21 bc (27.26)	27.63 abc (31.7)	34 a (35.67)	35.63 a (36.65)	34.88 a (36.19)	33.67 a (35.46)	27.18
T3	Groundnut + Foxtail millet	12.5 c (20.7)	12.67 c (20.85)	20.2 bc (26.7)	24.67 bc (29.7)	34.78 a (36.12)	34.5 a (35.97)	35.62 a (36.64)	33.99 a (35.62)	26.12
T4	Groundnut + Sorghum	16 ab (23.56)	18.33 ab (25.35)	24.04 ab (29.28)	30.05 ab (33.22)	32.78 ab (34.92)	31.49 ab (34.13)	30.14 cd (33.29)	28.67 ab (32.33)	26.43
T5	Groundnut + Cowpea	14.58 abc (22.45)	17.6 ab (24.8)	25 ab (30)	27.4 abc (31.55)	26.11 bc (30.72)	26.11 b (30.72)	24.96 e (29.97)	20.33 c (26.8)	22.76
T6	Groundnut + Cluster bean	13.6 bc (21.64)	16.96 b (24.32)	23.27 ab (28.83)	27.94 abc (31.9)	36.75 a (37.31)	35.85 a (36.78)	35.58 a (36.62)	33 a (35.06)	27.87
T7	Groundnut + Sunflower	15.53 ab (23.2)	17.97 a (25.06)	21.97 abc (27.92)	29.16 ab (32.68)	31.49 abc (34.13)	30.82 ab (33.7)	30.56 bc (33.55)	27 abc (31.28)	25.56
T8	Groundnut sole crop	16.86 a (24.14)	19.89 a (26.31)	27 a (31.22)	33.97 a (35.62)	36.88 a (37.26)	36.78 a (37.21)	37.66 a (37.85)	35.33 a (36.32)	30.55
	S.Em. ±	0.78	1	1.22	1.35	1.48	1.42	0.73	1.76	
	C.D. (5%)	2.37	3.03	3.69	4.10	4.50	4.32	2.20	5.34	
	C. V. (%)	6.07	7.1	7.45	7.35	7.45	7.14	3.66	9.33	

Table 3 Effect of intercropping on population of leaf miner during kharif 2022-23 at Dharwad.

Treatments		No. of larvae per plant								Cumulative mean
		1 week	2 weeks	3 weeks	4 weeks	5 weeks	6 weeks	7 weeks	8 weeks	
T1	Groundnut + Pearl millet	0 (0.71)	a 1.49 (1.41)	c 1.38 (1.37)	c 1.1 (1.27)	c 0.84 (1.16)	b 0.63 (1.11)	b 1.05 (1.24)	bc 1.12 (1.27)	0.95
T2	Groundnut + Maize	0 (0.71)	a 1.88 (1.54)	bc 1.87 (1.54)	abc 1.18 (1.30)	c 0.91 (1.191)	b 0.62 (1.13)	b 1.01 (1.23)	bc 1.01 (1.23)	1.06
T3	Groundnut + Foxtail millet	0 (0.71)	a 1.52 (1.4)	c 1.5 (1.41)	c 1.16 (1.29)	c 0.88 (1.17)	b 0.65 (1.1)	b 0.93 (1.19)	c 0.96 (1.21)	0.95
T4	Groundnut + Sorghum	0 (0.71)	a 1.66 (1.47)	bc 1.61 (1.42)	c 1.19 (1.3)	c 0.92 (1.19)	b 0.71 (1.14)	b 1.06 (1.25)	bc 1.06 (1.25)	1.03
T5	Groundnut + Cowpea	0 (0.71)	a 1.99 (1.58)	abc 2.22 (1.65)	abc 2.09 (1.61)	a 1.02 (1.23)	b 0.8 (1.06)	b 1.08 (1.26)	bc 1.08 (1.26)	1.28
T6	Groundnut + Cluster bean	0 (0.71)	a 1.92 (1.55)	abc 1.85 (1.53)	bc 1.22 (1.31)	bc 0.95 (1.2)	b 0.73 (1.07)	b 1.16 (1.29)	abc 1.22 (1.31)	1.13
T7	Groundnut + Sunflower	0 (0.71)	a 2.34 (1.69)	ab 2.45 (1.72)	ab 2.18 (1.64)	a 1.14 (1.28)	ab 0.77 (1.17)	b 1.18 (1.3)	ab 1.66 (1.47)	1.47
T8	Groundnut sole crop	0 (0.71)	a 2.8 (1.8)	a 2.87 (1.82)	a 2.24 (1.62)	a 1.46 (1.39)	a 1 (2.53)	a 1.37 (1.37)	a 1.83 (1.51)	1.7
	S.Em. ±	0	0.08	0.09	0.08	0.05	0.09	0.03	0.07	
	C.D. (5%)	0.00	0.25	0.28	0.26	0.14	0.28	0.10	0.22	
	C. V. (%)	0	9.2	10.24	10.37	6.51	12.5	4.32	9.37	

References

- Agasimani, C. A., Ravishankar, G., Mannikeri, I. M., Patil, R. K. and Giriraj, K., 1993, Groundnut based intercropping systems. Paper presented In: national seminar on oilseed research and development in India: status and strategies, pp 161.
- Amin P W and Mohammad, 1980, Groundnut pest research at ICRISAT. *Proceedings of the International Workshop on Groundnut*, ICRISAT Center, Patancheru, Andhra Pradesh, India, pp. 158-166.
- Anonymous, 2019, District – wise and crop – wise area coverage during summer season, Karnataka state department on agriculture. www.raitamitra.kar.nic.in
- Balakrishnan, L, N., Murali Baskaran, N. K. and Mahadevan, N. R., 2010, Influence of intercrops/trap crops on the preference of major pests of cotton in different IPM modules under rainfed condition. *J. Biop.*, 3 (1): 373 – 378.
- Blummel M, Ramakrishna Reddy C, Ravi D, Nigam S N and Upadhyaya H D, 2005, Food-fodder traits in groundnut. *Journal of SAT Agricultural Research*, 1(1): 1-3.
- Dodiya R D, Barad A H, Italiya J V, Pathan N P and Bhatt N A, 2023, Host suitability of tobacco leaf eating caterpillar (*Spodoptera litura* F.) under laboratory condition. *International Journal of Bio-resource and Stress Management*, 14(8): 1186-1195.
- Dodiya R D, Barad A H, Italiya J V and Prajapati H N, 2024, Impact of weather parameters on population dynamics of tobacco leaf eating caterpillar, *Spodoptera litura* (F.) infesting groundnut. *Environment and Ecology*, 42(1A): 301-306.
- Dutta D and Bandyopadhyay P, 2020, Production potential of intercropping of groundnut (*Arachis hypogea*) with pigeonpea (*Cajanus cajana*) and maize under various proportions in rainfed. *Indian Journal of Agronomy*, 51(2):103-106.
- Girija T, Mahabaleshwar Hegde and Virupaksha Prabhu, 2015, Impact of intercrops on pests of groundnut. *Journal of Experimental Zoology*, 18: 133-138.

- Kennedy, F. J. S., Rajamanickam. K. and Raveendran. T. S., 1990, Effect of intercropping on insect pests of groundnut and their natural enemies. *J. Biol. Control*, 4 (1): 63-64.
- Murali Baskaran, R. and Thangavelu, S., 1990, Influence of intercrops on the incidence of leaf miner (*Aproaecema modicella* Deventer) in groundnut *J. Oilseed Res.*, 7:143-146.
- Nath, P. and Singh, A. K., 1998, Effect of intercropping of groundnut with millets and pigeon pea on the relative incidence of insect pest. *Ann. Pl. Protec.*, 6: 151-154.
- Padgham D E, Kimmins F M and Rao G R, 1990, Resistance in groundnut, *Arachis hypogaea* L. to *Aphis craccivora* Koch. *Annals of Applied Biology*, 17(2): 285-295
- Patil, R. K., 1993, Progress of research in groundnut pest management at Dharwad. Paper presented In: *Prospective in Entomological Research for Sustainable Agriculture in Northern Karnataka.*, UAS Dharwad, pp. 132.
- Ranga Rao G V and Wightman J A, 1997, Techniques for screening groundnut genotypes for resistance to pests, *Spodoptera litura* (F.) in India. *Procedures of National Scientist Forum Spodoptera litura*, ICRISAT, Patancharu. pp. 68-75.
- Rashmi, Y., Hegde, M. G., Patil, R. K. and Balikai, R. A., 2011, Evaluation of IPM modules for management of defoliators in groundnut. *Ann. Pl. Protec. Sci.*, 19 (2): 321-323.
- Singh, T. V. K., Singh, K. M. and Singh, R. N., 1991, Influence of intercropping on the incidence of major pests in groundnut (*Arachis hypogaea*) Linn. *Indian J. Entomol.*, 53: 18-44.