

# ORGANIC GRAIN PROTECTANTS FOR ENHANCING STORABILITY OF RICE SEEDS

## ABSTRACT

An experiment was conducted at the Department of Seed Science and Technology, College of Agriculture, Vellayani to study the influence of organic grain protectants in enhancing storage life of rice seeds. The experiment consists of six treatments with different organic grain protectants viz Arka (*Calotropis gigantea*) leaf powder, Neem (*Azadirachta indica*) leaf powder, Pongamia (*Pongamia pinnata*) leaf powder, Sweet flag (*Acorus calamus*) rhizome powder, Arappu (*Albizia amara*) leaf powder and control. All the treatments were effective in control of storage pest and mycoflora infestation without any seed damage and weight loss compared to control with 62.333, 16.253 per cent seed damage and weight loss. The treatment T<sub>4</sub> [Sweet flag (*Acorus calamus*) rhizome powder] recorded superior among all treatments, germination percentage (82.00%), germination energy (81.00%), seedling vigour index (1029.20), seedling shoot length (13.03 cm), seedling root length (12.40 cm). However, among all treatments T<sub>4</sub> [Sweet flag (*Acorus calamus*) rhizome powder] recorded least Seed damage percentage, seed weight loss percentage, total number of adult insects and mycoflora infestation compared to control.

*Keywords: Rice seeds, Organic grain protectants, Seed storage, Storage pest, Mycoflora infestation and Seed germination.*

## 1. Introduction

Seed is the fertilized, matured ovule and a carrier of genetic potential for sustainable crop production. It is the basic and crucial input of agriculture around which all other input acts. Rice (*Oryza sativa*) is a staple food for half of the world's population but it is severely threatened by diseases. The population may meet the requirements of only 60% of the population (FAO, 2018). China is the leading producer of rice (142.3 million tonnes) followed by India (110.4 million tonnes, thereby plays a major role in meeting the rice demand. Seed storage is an essential segment of seed industry. In storage, viability and vigour of the seeds is regulated by many physico-chemical factors like moisture content of the seed atmospheric humidity, temperature, initial seed quality, physical and chemical composition of seed, gaseous exchange, storage structure and packaging materials. The production and supply of quality seed is one of the important factors responsible for the increased productivity and production of any crop. The highest quality of the seed is attained under that complex of conditions evoking the most favourable interactions Storage pest infestation, Seed germination and mycoflora between genetic makeup of a seed and the environment under which it is produced, harvested, processed and stored. The seed potentiation is mainly achieved by treating the seeds with various chemicals and botanicals can reduce the infestation and maintain the quality of the seed in term of viability and vigour for longer period in storage. Maintenance of seed quality during storage period is important not only for successful crop production but also for maintaining the quality and integrity of the seed that are inconstant threat of genetic erosion. The use of naturally occurring seed protectants and usage of bio-control agents is powerful alternative. Use of locally available plant materials to protect seeds from insect pests is a common age-old practice and the extracts from different plants have been known to possess insecticidal properties against wide range of pests (Isman, 2008). Standardize of seed treatment with organic grain protectants would be of great advantage to reduce the problems in maintenance of seed quality during storage. Leaf and fruit powders of herbal plants are widely used for seed treatment (Bashyam, 1999). The performance of dry seed treatment with crude plant materials in powder form have been found to significantly slow down the deterioration of seeds under various ageing conditions (De *et al.*, 2004 in

wheat; Rudrapal and Basu, 2004 in French bean; Sengupta *et al.*, 2005 in onion; Kundagrami *et al.*, 2008 in rice). The pest-controlling efficacy of many plant derivatives has already been proved against several storage pests and they can improve seed quality. A number of synthetic and natural components have been suggested as potential candidate for seed treatment. The almost non-existent and therefore could be beneficial to our agricultural sector. However, their efficiency as well as effect on seed quality remains to be observed (Kumar *et al.*, 2007).

## 2. Materials and methods

The experiment was conducted in Department of Seed Science and Technology, College of Agriculture, Vellayani, Thiruvananthapuram to study the effects of organic grain protectants in enhancing storage life of rice seeds. The experiment consists of six treatments viz T<sub>1</sub>: Arka (*Calotropis gigantea*) leaf powder @ 12.5g, T<sub>2</sub>: Neem (*Azadirachta indica*) leaf powder @ 12.5g, T<sub>3</sub>: Pongamia (*Pongamia pinnata*) leaf powder @ 12.5g, T<sub>4</sub>: Sweet flag (*Acorus calamus*) rhizome powder @ 5g, T<sub>5</sub>: Arappu (*Albizia amara*) leaf powder @ 12.5g, T<sub>6</sub>: Control. The experiment was conducted in Completely Randomized Block Design (CRD) in three replications. 500 grams of seed was used in each replication and packed in cloth bags stored for ten months at room temperature. The bimonthly observations on various parameters were carried out and statistically analysed. The percentage seed damage was determined by collecting a sample of 100 seeds from each three replications of each treatment at bimonthly intervals proposed by Adams and Schulten (1978). Seed weight loss percentage was calculated from number of damaged and undamaged seeds, weight of damaged and undamaged seeds as given by Adams and Schulten (1978). Total number of adult insects was also calculated at bimonthly intervals. Mycoflora infestation was assessed by agar plate method. The germination test was carried out with 100 seeds in four replications with rolled paper towel method for 8 days (ISTA, 2013). Normal healthy seeds were taken for germination test. Other germination parameters like germination energy, seedling shoot and root length, Seedling vigour index (Abdul- Baki and Anderson, 1973) was calculated at bimonthly intervals.

$$\text{Seed damage percentage} = \frac{\text{Number of damaged seeds}}{\text{Total number of seeds taken}} \times 100$$

$$\text{Seed weight loss percentage} = \frac{U(Nd) - D(Nu)}{U(Nu + Nd)} \times 100$$

Nd- Number of damaged seeds

Nu- Number of undamaged seeds

D- Weight of damaged seeds

U- Weight of undamaged seeds

$$\text{Germination percentage} = \frac{\text{Number of seeds germinated}}{\text{Total number of seeds taken}} \times 100$$

$$\text{Germination energy} = \frac{\text{Number of seeds germinated on day 4 count}}{\text{Total number of seeds tested}} \times 100$$

Seedling vigour index = Per cent seed germination x Seedling length

## 3. Results and discussion

### Storage pest infestation assessment

Among the different treatments all the organic grain protectants used were significantly superior to control. Seed damage was not observed in treatments T<sub>1</sub> [*Calotropis gigantea* leaf powder @ 12.5g/500g], T<sub>2</sub> [*Azadirachta indica* leaf powder @ 12.5g/500g], T<sub>3</sub> [*Pongamia pinnata* leaf powder @ 12.5g/500g] for up to four months of storage and T<sub>5</sub> [*Albizia amara* leaf powder @ 12.5g/500g] for up to two months of storage. No

seed damage and weight loss were observed in T<sub>4</sub> [*Acorus calamus* rhizome powder @ 5g/500g] for up to six months of storage. In contrast, seed infestation was observed from initial month up to ten months in control. At the end of storage period, seed damage percentage was highest in control (62.333%) which was then followed by T<sub>5</sub> (20.333%), T<sub>2</sub> (13.333%), T<sub>1</sub> (12.333%). Treatment T<sub>4</sub> recorded damage percentage 5.667% which was on par with T<sub>3</sub> (8.667%). Seed weight loss was highest in control (16.253%) lowest in T<sub>4</sub> (7.537%). The results were in agreement the findings of Shukla *et al.* (2009) concluded that the dose of 5mg rhizome powder of *Acorus calamus* per gram proved fatal, causing 100 per cent mortality of *C. chinensis* in chickpea and completely inhibiting F1 emergence. In the study, three major storage pests were found infesting the seeds. various treatments were lesser grain borer (*Rhyzopertha dominica*), rice weevil (*Sitophilus oryzae*) and Angoumois grain moth (*Sitotroga cerealella*) during different periods of storage. In control lesser grain borer (*Rhyzopertha dominica*), rice weevil (*Sitophilus oryzae*) and Angoumois grain moth (*Sitotroga cerealella*) was observed from two, four and six months of storage. Where as in T<sub>4</sub> adult insects were observed from sixth month- lesser grain borer (*Rhyzopertha dominica*) and rice weevil (*Sitophilus oryzae*) from eighth month of storage onwards. No incidence of Angoumois grain moth (*Sitotroga cerealella*) throughout the storage period. The total number of adult insects at the end of ten months storage in control was lesser grain borer (*Rhyzopertha dominica*) with 7.667%, rice weevil (*Sitophilus oryzae*) with 7.333% and Angoumois grain moth (*Sitotroga cerealella*) with 2.333%. In contrast in T<sub>4</sub> lesser grain borer (*Rhyzopertha dominica*) with 1.667%, rice weevil (*Sitophilus oryzae*) with 0.667%. Paul *et al.*, 1965 and Jillani and Haq, 1984 reported that *Acorus calamus* rhizome powder repelled the population of *S. oryzae* and *R. dominica*. Similarly, in another test conducted against Angoumois grain moth, (*Sitotroga cerealella* O.), it was found that 100% mortality was resulted when rhizome dust of *A. calamus* was used at 10g/kg of rice grain (Timsina *et al.*, 2007). The results of the present findings are agreement with the results of Kudachi and Balikai (2009) who reported that *A. calamus* rhizome powder @ 1 percent recorded percent adult mortality of *R. dominica* with zero percent seed damage, minimum percent seed weight loss of zero percent.

**Table 1: Effect of Organic grain protectants on seed damage percentage and seed weight loss percentage**

Treatment	Seed damage percentage			Seed weight loss percentage		
	2 MAS	6 MAS	10 MAS	2 MAS	6 MAS	10 MAS
T <sub>1</sub> <i>Calotropis gigantea</i> leaf powder @ 12.5g/500g	0.000	1.333	12.333	0.000	2.370	10.993
T <sub>2</sub> <i>Azadirachta indica</i> leaf powder @ 12.5g/500g	0.000	1.667	13.333	0.000	2.333	10.320
T <sub>3</sub> <i>Pongamia pinnata</i> leaf powder @ 12.5g/500g	0.000	1.000	8.667	0.000	2.327	10.037
T <sub>4</sub> <i>Acorus calamus</i> rhizome powder @ 5g/500g	0.000	0.333	5.667	0.000	1.033	7.537
T <sub>5</sub> <i>Albizia amara</i> leaf powder @ 12.5g/500g	0.333	2.333	20.333	0.333	3.033	11.217
T <sub>6</sub> Control	1.333	13.667	62.333	3.000	9.467	16.253
SE (m)	0.19	0.87	2.76	0.43	0.58	0.81
CD (5%)	NS	2.685	8.532	NS	1.809	2.500

**Table 2: Effect of organic grain protectants on total number of adult insects**

Treatment	Lesser grain borer ( <i>Rhyzopertha dominica</i> )			Rice weevil ( <i>Sitophilus oryzae</i> )			Angoumois grain moth ( <i>Sitotroga cerealella</i> )		
	2 MAS	6 MAS	10 MAS	2 MAS	6 MAS	10 MAS	2 MAS	6 MAS	10 MAS
T <sub>1</sub> <i>Calotropis gigantea</i> leaf powder @ 12.5g/500g	0.000	0.667	3.333	0.000	0.333	4.333	0.000	0.000	0.667
T <sub>2</sub> <i>Azadirachta indica</i> leaf powder @ 12.5g/500g	0.000	0.667	2.667	0.000	0.000	3.333	0.000	0.000	0.000
T <sub>3</sub> <i>Pongamia pinnata</i> leaf powder @ 12.5g/500g	0.000	0.333	1.667	0.000	0.000	1.333	0.000	0.000	0.000
T <sub>4</sub> <i>Albizia amara</i> leaf powder @ 12.5g/500g	0.000	0.333	1.667	0.000	0.000	0.667	0.000	0.000	0.000
T <sub>5</sub> <i>Albizia amara</i> leaf powder @ 12.5g/500g	0.000	1.333	6.333	0.000	0.667	5.333	0.000	0.000	1.333
T <sub>6</sub> Control	0.667	4.333	7.667	0.000	1.667	7.333	0.000	0.667	2.333
SE (m)	0.27	0.62	1.24	0.00	0.40	1.20	0.00	0.13	0.62
CD (5%)	NS	1.922	3.843	NS	NS	3.703	NS	NS	NS

### Mycoflora infestation assessment

In the assessment of mycoflora infestation was highest in control (69.33%) and least in T<sub>4</sub> (10.16%) which was recorded from eighth month of storage onwards, whereas in control from fourth month onwards. The infested mycoflora in storage was *Aspergillus flavus*. *A. calamus* shows growth inhibitory effect against *Aspergillus flavus* and *Aspergillus niger*. Since the plant is easily available to farmers, it could serve as an alternate approach to chemical fungicides which are costly and highly toxic (Bisht, 2011).

**Table 3: Effect of organic grain protectants on mycoflora infestation**

Treatment	Mycoflora infestation in seeds (%)		
	2 MAS	6 MAS	10 MAS
T <sub>1</sub> <i>Calotropis gigantea</i> leaf powder @ 12.5g/500g	0.00	6.67	20.00
T <sub>2</sub> <i>Azadirachta indica</i> leaf powder @ 12.5g/500g	0.00	2.67	25.12
T <sub>3</sub> <i>Pongamia pinnata</i> leaf powder @ 12.5g/500g	0.00	2.67	17.13
T <sub>4</sub> <i>Acorus calamus</i> rhizome powder @ 5g/500g	0.00	0.00	10.16
T <sub>5</sub> <i>Albizia amara</i> leaf powder @ 12.5g/500g	0.00	4.66	37.13
T <sub>6</sub> Control	0.00	42.66	69.33
SE (m)	0.00	2.09	5.86
CD (5%)	NS	6.442	18.065

## Seed germination parameters

Organic grain protectants improved the seed germination parameters compared to control. After ten months of storage seed treated with T<sub>4</sub> Sweet flag (*Acorus calamus*) rhizome powder recorded highest in seed germination percentage, vigour index and seedling parameters. The results were in accordance with (Anandi R, 2001; Parameswari, 2002) seed treated with *Acorus calamus* rhizome powder excelled other botanicals in the seed quality maintenance. At the end of ten months of storage period T<sub>4</sub> recorded highest germination per cent (82.00%) which was on par with T<sub>3</sub> (80.00%). Also, T<sub>6</sub> recorded the lowest germination per cent (79.00%) among all treatments. Satisfactory germination (80%) as per the Indian minimum seed certification standards for rice was maintained for 12 months and even more period with sweet flag rhizome powder seed treatment. With respect to sweet flag rhizome powder, higher germination noticed may be related to its antioxidant property of  $\beta$  asarone which acts as a scavenger of free radicals and inhibit seed deterioration. These results are in accord with the findings of Bhargude *et al.* (2021) in sorghum. The treatment T<sub>4</sub> recorded maximum germination energy, seedling vigour index, seedling shoot and root length among all treatments with 81.00%, 1020.20, 13.03 cm and 12.40 cm. In contrast control was recorded lowest germination energy, seedling vigour index, seedling shoot and root length with 75.33%, 942.00, 10.26cm and 11.80 cm.

**Table 4: Effect of organic grain protectants on germination parameters**

Treatment	Germination percentage			Germination energy		
	2 MAS	6 MAS	10 MAS	2 MAS	6 MAS	10 MAS
T <sub>1</sub> <i>Calotropis gigantea</i> leaf powder @ 12.5g/500g	95.33	89.66	79.66	86.67	82.33	77.33
T <sub>2</sub> <i>Azadirachta indica</i> leaf powder @ 12.5g/500g	96.33	89.00	80.67	88.67	83.00	77.00
T <sub>3</sub> <i>Pongamia pinnata</i> leaf powder @ 12.5g/500g	96.00	89.66	80.00	89.00	84.00	78.67
T <sub>4</sub> <i>Acorus calamus</i> rhizome powder @ 5g/500g	97.66	91.00	82.00	89.67	86.00	81.00
T <sub>5</sub> <i>Albizia amara</i> leaf powder @ 12.5g/500g	96.00	89.33	79.66	88.00	82.67	77.67
T <sub>6</sub> Control	95.33	87.60	74.00	87.66	82.00	75.33
SE (m)	0.72	0.79	0.94	0.79	1.21	0.86
CD (5%)	NS	2.445	2.905	NS	NS	2.652

The increased rate of imbibition, where the tiny particles in the coating act as a "wick" or moisture-attracting material, or possibly to improve germination, was the cause of the seed treated with *Acorus calamus* rhizome powder's improved vigour metrics (Renugadevi *et al.*, 2006). Gadewar *et al.* (2017) observed sorghum seeds treated with sweet flag powder (2.5 per cent) showed significantly higher 100 seed weight, germination percentage, seedling vigour, field emergence percentage compared to other seed treatment and control during storage. Regarding seed treatments, the seeds treated with sweet flag rhizome powder at a rate of 20 g kg<sup>-1</sup> had the greatest root length (13.40 cm), shoot (8.98 cm), seedling length (22.39 cm), and seedling vigour index I (2037) after 12 months of storage. (Radha *et al.*, 2024). Improved germination rates are a result of maintaining seed integrity and reducing insect infestation (Latha & Naganagoud, 2015). The powder contains bio active compounds that can stimulate seed metabolism and enhance the overall vigour of the seeds. Treatment with sweet flag rhizome

powder helps maintain seed viability during storage by reducing seed damage and preserving seed quality. The powder may also improve the availability of essential nutrients to the seeds, promoting better germination and early seedling growth.

**Table 5: Effect of organic grain protectants on seedling vigour index, seedling shoot length and seedling root length**

Treatment	Seedling vigour index			Seedling shoot length			Seedling root length		
	2 MAS	6 MAS	10 MAS	2 MAS	6 MAS	10 MAS	2 MAS	6 MAS	10 MAS
T <sub>1</sub> <i>Calotropis gigantea</i> leaf powder @ 12.5g/500g	1218.3	1144.2	959.7	12.33	11.73	10.93	12.66	12.63	11.90
T <sub>2</sub> <i>Azadirachta indica</i> leaf powder @ 12.5g/500g	1207.0	1152.2	991.4	11.63	11.70	10.30	12.40	12.80	12.16
T <sub>3</sub> <i>Pongamia pinnata</i> leaf powder @ 12.5g/500g	1349.7	1178.1	988.0	13.50	12.93	12.00	13.93	13.00	12.20
T <sub>4</sub> <i>Acorus calamus</i> rhizome powder @ 5g/500g	1381.2	1221.2	1029.0	14.06	13.66	13.03	14.00	13.26	12.40
T <sub>5</sub> <i>Albizia amara</i> leaf powder @ 12.5g/500g	1184.6	1130.4	961.0	11.40	11.66	10.33	12.23	12.53	11.90
T <sub>6</sub> Control	1341.7	1100.0	942.00	12.53	11.66	10.20	13.90	12.40	11.80
SE (m)	83.45	91.61	58.64	0.28	0.30	0.26	0.90	0.99	0.68
CD (5%)	NS	282.28	180.70	0.887	0.934	0.812	NS	NS	2.112

## CONCLUSION

Out of all the organic grain protectants used in this investigation, *Acorus calamus* was able to sustain germination above IMSCS and provided total seed protection by inhibiting *Sitophilus oryzae*-induced seed damage. *A. calamus* has a well-established insecticidal effect on a variety of insect pests. The current study suggests using *Acorus calamus* rhizomes to suppress weevils. Because *A. calamus* products are used as medications all over the world, they may be less dangerous to people than the majority of traditional pesticides. Herbal plants that are readily available and virtually harmless to the environment and human health are used as organic grain protectants. We agree that some traditionally used botanicals can be an excellent alternative to chemical pesticides in stored grains if used in the correct dose and method. The ease in availability and simplicity in these botanicals are most likely to be accepted and adopted by the farmers.

## REFERENCES

1. Isman, M.B., 2008. Botanical insecticides: for richer, for poorer. *Pest Management Science: formerly Pesticide Science*, 64(1), pp.8-11.
2. Bashyam, S., 1999. Seed quality testing and management procedures in three important condiments of coriander, fennel and fenugreek. M. Sc. (Ag.) Thesis, Tamil Nadu Agricultural University, Coimbatore.

3. De, B.K., Mandal, A.K. and Basu, R.N., 2004. Compatibility of fungicides with pre-storage dry and mid-storage wet physiological seed treatments for storability and field performance of wheat (*Triticum aestivum* L.).
4. Rudrapal, D. and Basu, R.N., 2004. The role of initial seed vigour status in the expression of invigoration effect of wet and dry seed treatments in French bean (*Phaseolus vulgaris* L.). *Seed Sci. and Technol.*, 32(1), pp.205-212.
5. Sengupta, A.K., De, B.K. and Mandal, A.K., 2005. Pre-storage seed invigoration treatments for the maintenance of vigour, viability and field performance of high-vigour onion seed (*Allium cepa* L.). *Seed Sci. and Technol.*, 33(3), pp.753-760.
6. Kundagrami, S., De, B.K. and Manual, A.K., 2008. Post-harvest seed invigoration treatments for improved storability and field performance of rice (*Oryza sativa* L.).
7. Kumar, D. and Kalita, P., 2017. Reducing postharvest losses during storage of grain crops to strengthen food security in developing countries. *Foods*, 6(1), p.8.
8. Adams JM and Schulten GGM. Losses caused by insects, mites and microorganisms. PostHarvest Grain Loss Assessment Methods. American Association of Cereal Chemists, St. Paul, Minnesota. USA, 1978; pp.98-100
9. Abdul-Baki AA and Anderson JD. Physiological and biochemical deterioration of seed. In: Kozlowski, T.T., Seed Biology (2nd eds). Academic Press, New York, London, 1973; pp. 283-315.
10. ISTA International Seed Testing Association International rules for seed testing. *Seed Sci. and Technol.* 2013; 27: 25-30
11. Bisht, D., Pal, A., Chanotiya, C.S., Mishra, D. and Pandey, K.N., 2011. Terpenoid composition and antifungal activity of three commercially important essential oils against *Aspergillus flavus* and *Aspergillus niger*. *Nat. product res.*, 25(20) pp.1993-1998.
12. Radha J, Shakuntala NM, Basavegouda, Sangeetha M, Mahantashivayogayya K, Masthanreddy BG. Effect of packaging materials and seed treatments on seed quality parameters of rice during seed storage. *Int J Res Agron* 2024;7(9):526-532
13. Bhargude, A.R., Patil, S.K. and Kadam, D.R., 2021. Eco-friendly management of rice weevil (*Sitophilus oryzae* Linnaeus) on Sorghum during storage. *J Entomol Zool Stud*, 9(1), pp.1647-1652.
14. H., C., Latha., A., Naganagoud. (2015). Effect of different packaging materials on the efficacy of sweet flag rhizome powder (*Acorus calamus* L.) treated sorghum against *Sitophilus oryzae*. *J. of Appl. and Nat. Sci.*, 7(2), 922-926.
15. Renugadevi, J., Natarajan, N., Rajasekaran, R. and Srimathi, P., 2006. Bio-efficacy of promising botanicals against insect infesting cowpea. cv. CO 4. *Indian J. of Agric. Res.*, 40(4), pp.262-266.
16. Parameswari, K., 2002. Strategies to improve seed production and shelf life of pigeonpea *cajanus cajan* I Millsp Hybrid coph 2.
17. Anandi R. Seed technological studies in cowpea CO 5 (*Vigna unguiculata* L. wallp), M.Sc (Ag.) Thesis., Tamil Nadu Agricultural University, Coimbatore, India 2001
18. Timsina, G.P., Bajracharya, A.S.R., Sah, Y.P., Mandal, D.L. and Mehata, S.N., 2007. Assessment of storage losses of maize by angouimois grain moth (*Sitotroga crealella* Oliver) and weevils (*Sitophilus spp* Linn.) in farmers' storage condition.
19. Kudachi, D.C. and Balikai, R.A., 2009. Efficacy of botanicals for the management of lesser grain borer, *Rhyzopertha dominica* Fab. in sorghum during storage.

20. Gadewar, R., Babhulkar, V., Lambat, P., Lambat, A., Parate, R. and Charjan, S., 2017. The influence of some botanicals against rice weevil during storage in rabi sorghum. *Int. J. of Res. Biol. Agric. and Technol.*, 5(1), pp.28-30.
21. Shukla, A.C., Shahi, S.K. and Anupam Dikshit, A.D., 2002. *Eucalyptus pauciflora* potential source of sustainable, ecofriendly storage pesticide.
22. Paul, C.F., Agarwal, P.N. and Ausat, A., 1965. Toxicity of solvent extract of *Acorus calamus* Linn, to some grain pests and termites.
23. Jilani, G. and Haq, H.S., 1984. Studies on some indigenous plant materials as grain protectant against insect pests of stored grains. Pak. *Entomol*, 6(24).

UNDER PEER REVIEW