

## **Moringa oleifera: The Miracle Tree - Nutritional Powerhouse and Medicinal Marvel**

### **Abstract**

The "Miracle Tree," *Moringa oleifera*, is a nutrient-dense plant with many health advantages. It has strong anti-inflammatory, antidiabetic, anti-cancer, and antioxidant qualities. It is also rich in vitamins, minerals, and bioactive compounds. This review examines its nutritional worth, therapeutic applications, and industrial uses, such as food fortification and water purification. With potential for wider application in international health and business, moringa is an essential tool in the fight against malnutrition and the advancement of sustainable agriculture. More research is required to maximize its cultivation and increase its advantages.

**Keywords:** *Moringa oleifera*, Traditional medicine, Anticancer potential, Phytochemicals, Sustainable agriculture

### **Introduction**

*Moringa oleifera*, often referred to as the "Miracle Tree" or "Drumstick Tree," is one of the most versatile and nutritious plants known to humanity. Native to India, this fast-growing, drought-resistant tree thrives in tropical and subtropical regions worldwide, where it has been used for centuries as a food source and a traditional medicine. In recent years, the global interest in *Moringa oleifera* has surged due to its exceptional nutritional profile and potential health benefits, particularly in addressing malnutrition and chronic diseases (Gopalakrishnan *et al.*, 2016).

The popularity of *Moringa oleifera* is not only due to its nutritional and medicinal properties but also because of its adaptability to various environmental conditions. This adaptability makes it a valuable crop in regions prone to food insecurity and harsh climates. Moreover, the growing body of scientific evidence supporting the health benefits of *Moringa oleifera* has further fueled its recognition as a superfood with a wide range of applications, from combating malnutrition to managing chronic diseases such as diabetes and cancer (Mbikay, 2012; Tiloke *et al.*, 2013).

### **Nutritional Composition of *Moringa oleifera***

#### **Overview of Nutrients**

*Moringa oleifera* is renowned for its remarkable nutritional profile, which includes a rich array of vitamins, minerals, proteins, and essential phytochemicals. The leaves, seeds, pods, and flowers of the tree are all edible and contain significant nutritional value, making *Moringa oleifera* a critical resource in combating malnutrition, especially in developing countries (Fuglie, 2005).

The leaves of *Moringa oleifera* are particularly nutrient-dense. They contain seven times more vitamin C than oranges, ten times more vitamin A than carrots, 17 times more calcium than milk, nine times more protein than yogurt, 15 times more potassium than bananas, and 25 times more iron than spinach (Gopalakrishnan *et al.*, 2016). The leaves are also a complete source of protein, containing all essential amino acids, which is unusual for a plant-based food source. This makes *Moringa oleifera* leaves an excellent dietary supplement for individuals with limited access to animal proteins (Yang *et al.*, 2006).

*Moringa oleifera* seeds have nutritional value as well. They are rich in proteins, a variety of vitamins and minerals, and oleic acid, a type of monounsaturated fat. The seeds are especially high in calcium, potassium, and zinc, all of which are necessary for supporting the immune system, preserving bone health, and cardiovascular health (Mbikay, 2012). Although they are less frequently eaten than the leaves and seeds of *Moringa oleifera*, the pods and flowers still have a lot of nutritional value. The pods are high in essential fatty acids, such as linoleic acid, which is beneficial for heart health, and dietary fibre. Because of their high nectar content, the flowers are an important source of nutrition and energy for both people and animals (Sánchez-Machado *et al.*, 2010).

### **Impact of Environmental Factors on Nutrient Content**

The nutrient content of *Moringa oleifera* can vary significantly depending on environmental factors such as soil quality, climate, and cultivation practices. For example, studies have shown that *Moringa* plants grown in different regions can have different levels of vitamins and minerals, which may affect their overall nutritional value (Asante *et al.*, 2014). In regions with high temperatures and low rainfall, *Moringa* plants tend to have higher concentrations of certain nutrients, such as vitamin C and beta-carotene, compared to plants grown in more temperate climates. This is likely due to the plant's ability to adapt to environmental stress by producing higher levels of antioxidants to protect itself from oxidative damage (Moyo *et al.*, 2011). Soil quality also plays a crucial role in determining the nutrient content of *Moringa oleifera*. Plants grown in nutrient-rich soils with adequate levels of nitrogen, phosphorus, and potassium tend to have higher concentrations of proteins, vitamins, and minerals. Conversely, plants grown in poor soils may have lower nutrient levels, which can affect their overall health benefits (Dania *et al.*, 2014). To optimize the nutritional content of *Moringa oleifera*, it is important to consider the specific environmental conditions in which the plant is grown. This may involve selecting the appropriate soil type, adjusting irrigation practices, and using organic fertilizers to enhance soil fertility. By optimizing these factors, it is possible to produce *Moringa* plants with the highest possible nutritional value, making them an even more valuable resource for addressing malnutrition and promoting health (Gopalakrishnan *et al.*, 2016).

### **Medicinal Properties and Applications**

*Moringa oleifera* has been used in traditional medicine for centuries to treat a wide range of ailments, from digestive disorders to infectious diseases. In recent years, scientific research has confirmed many of the traditional uses of *Moringa oleifera* and has identified new potential health benefits, particularly in the prevention and management of chronic diseases (Mbikay, 2012).

#### **Antioxidant Properties**

One of the most well-documented medicinal properties of *Moringa oleifera* is its potent antioxidant activity. Antioxidants are compounds that help neutralize harmful free radicals in the body, which can cause oxidative stress and damage to cells and tissues. Oxidative stress is a major contributing factor to the development of chronic diseases such as cancer, cardiovascular disease, and neurodegenerative disorders (Berkovich *et al.*, 2013). *Moringa oleifera* is rich in a variety of antioxidants, including quercetin, chlorogenic acid, and beta-carotene. These compounds help protect the body against oxidative damage and may reduce the risk of developing chronic diseases. For example, quercetin has been shown to have anti-inflammatory, antiviral, and anticancer properties, making it a valuable component of the *Moringa* plant.

(Chumark *et al.*, 2008). In addition to its antioxidant properties, *Moringa oleifera* has also been shown to enhance the body's natural antioxidant defense system by increasing the activity of enzymes such as superoxide dismutase (SOD) and catalase. These enzymes play a crucial role in neutralizing free radicals and preventing oxidative damage to cells and tissues (Tiloke *et al.*, 2013).

### **Anticancer Properties**

The anticancer potential of *Moringa oleifera* is another area of growing interest in the scientific community. Studies have shown that extracts from the leaves, seeds, and pods of *Moringa* have potent anti-proliferative effects on cancer cells, meaning they can inhibit the growth and spread of cancerous cells in the body (Tiloke *et al.*, 2013).

The anticancer activity of *Moringa oleifera* is thought to be due to its high concentration of glucosinolates, benzyl isothiocyanate (BITC), and niazimicin. These compounds have been shown to induce apoptosis (programmed cell death) in cancer cells while sparing healthy cells, making *Moringa* a promising natural treatment for cancer (Hermawan *et al.*, 2012).

For example, research has shown that BITC, a compound found in *Moringa* seeds, can inhibit the growth of breast cancer cells by inducing oxidative stress and triggering apoptosis. Similarly, niazimicin, a compound found in *Moringa* leaves, has been shown to inhibit the development of skin tumors in animal models, suggesting that it may have potential as a topical treatment for skin cancer (Nakamura *et al.*, 2002; Miyoshi *et al.*, 2004).

While the anticancer properties of *Moringa oleifera* are promising, more research is needed to fully understand the mechanisms by which these compounds exert their effects and to determine the most effective ways to use *Moringa* in cancer prevention and treatment (Liou & Storz, 2010).

### **Antidiabetic Properties**

Diabetes is a major global health concern, affecting millions of people worldwide. *Moringa oleifera* has shown promise as a natural treatment for diabetes, particularly in its ability to lower blood sugar levels and improve insulin sensitivity (Mbikay, 2012).

The antidiabetic properties of *Moringa oleifera* are thought to be due to its high concentration of polyphenols, flavonoids, and other bioactive compounds that have been shown to improve glucose metabolism and reduce oxidative stress in the body (Cerf, 2013). For example, studies have shown that the aqueous extracts of *Moringa* leaves and seeds can reduce fasting blood glucose levels and improve insulin sensitivity in animal models of diabetes (Divi *et al.*, 2012).

In addition to its blood sugar-lowering effects, *Moringa oleifera* has also been shown to protect against the complications of diabetes, such as retinopathy, nephropathy, and atherosclerosis. These complications are often the result of oxidative stress and inflammation, which *Moringa* can help mitigate through its antioxidant and anti-inflammatory properties (Chumark *et al.*, 2008).

The potential of *Moringa oleifera* as a natural treatment for diabetes is particularly important in developing countries, where access to conventional diabetes medications may be limited. By incorporating *Moringa*

into the diet, individuals with diabetes may be able to manage their condition more effectively and reduce their risk of complications (Cerf, 2013).

### **Anti-inflammatory and Antimicrobial Properties**

*Moringa oleifera* has long been used in traditional medicine as an anti-inflammatory and antimicrobial agent. The leaves, seeds, and roots of the plant contain a variety of bioactive compounds that have been shown to reduce inflammation and inhibit the growth of harmful bacteria and viruses (Viera *et al.*, 2010).

The anti-inflammatory properties of *Moringa oleifera* are primarily due to its high concentration of flavonoids, phenolic acids, and other polyphenols. These compounds have been shown to inhibit the production of pro-inflammatory cytokines and enzymes, such as tumor necrosis factor-alpha (TNF- $\alpha$ ) and cyclooxygenase-2 (COX-2), which play a key role in the inflammatory response (Mahajan & Mehta, 2009).

In addition to its anti-inflammatory effects, *Moringa oleifera* has also been shown to have broad-spectrum antimicrobial activity against a variety of pathogens, including bacteria, viruses, and fungi. For example, studies have shown that extracts from *Moringa* leaves and seeds can inhibit the growth of *Staphylococcus aureus*, *Escherichia coli*, and *Candida albicans*, among other pathogens (Viera *et al.*, 2010).

The antimicrobial properties of *Moringa oleifera* are thought to be due to the presence of compounds such as pterygospermin, benzyl isothiocyanate, and moringine, which have been shown to disrupt the cell membranes of bacteria and fungi, leading to their death (Hermawan *et al.*, 2012).

### **Commercial and Industrial Applications**

*Moringa oleifera* is not only valued for its nutritional and medicinal properties but also for its wide range of commercial and industrial applications. The versatility of the plant makes it an important resource in various industries, from agriculture to cosmetics and pharmaceuticals.

#### **Use in Water Purification**

One of the most significant commercial applications of *Moringa oleifera* is in water purification. The seeds of *Moringa* contain a natural coagulant, a cationic protein that can bind to impurities in water and cause them to settle out, making the water clear and safe to drink (Gopalakrishnan *et al.*, 2016). This natural coagulant is particularly valuable in regions where access to clean water is limited, as it provides a low-cost, sustainable solution for water treatment.

The effectiveness of *Moringa* seeds as a water purifier has been well-documented in scientific studies. For example, Suhartini *et al.* (2011) developed a two-stage clarifier using coconut fiber and sand media mixed with powdered *Moringa* seeds to treat tapioca starch wastewater. The system was highly effective at removing impurities and stabilizing pH, demonstrating the potential of *Moringa* seeds as a natural coagulant in wastewater treatment.

In addition to its use in water purification, *Moringa* seed extract has also been shown to have antimicrobial properties, making it effective at eliminating waterborne pathogens such as bacteria and viruses. This dual functionality makes *Moringa* seeds an ideal solution for improving water quality and preventing waterborne diseases in rural communities (Viera *et al.*, 2010).

## **Oil Extraction and Uses**

The seeds of *Moringa oleifera* are also used to extract oil, known as Ben oil, which is highly valued for its nutritional and cosmetic properties. Ben oil is rich in oleic acid, a type of monounsaturated fat that is beneficial for heart health. It also contains high levels of tocopherols (vitamin E) and sterols, which have antioxidant properties and can protect the skin from oxidative damage (Lalas & Tsaknis, 2002).

Ben oil is commonly used in cooking as a substitute for olive oil, as it has a similar fatty acid profile and a mild, pleasant flavor. It is also used in the cosmetic industry as an ingredient in skin and hair care products due to its moisturizing and anti-aging properties (Lalas & Tsaknis, 2002).

In addition to its nutritional and cosmetic uses, Ben oil has potential applications in the production of biodiesel. The high oleic acid content of the oil makes it an excellent feedstock for biodiesel production, providing a renewable and sustainable source of energy (Lalas & Tsaknis, 2002).

## **Food Fortification and Nutraceuticals**

*Moringa oleifera* is increasingly being used to fortify food products and develop nutraceuticals, particularly in regions where malnutrition is a significant concern. The high nutrient content of *Moringa* leaves, seeds, and pods makes them an ideal ingredient for enhancing the nutritional value of foods such as snacks, cereals, and beverages (Abou-Zaid & Nadir, 2014).

For example, *Moringa* leaves have been used to fortify cookies, bread, and cereal gruels, significantly increasing their protein, vitamin, and mineral content. Studies have shown that *Moringa*-fortified snacks are well accepted by consumers and can help address nutrient deficiencies in vulnerable populations (Owusu *et al.*, 2014).

In addition to food fortification, *Moringa oleifera* is also being used to develop nutraceutical products, such as supplements and functional foods, that provide health benefits beyond basic nutrition. These products are particularly popular in the wellness industry, where *Moringa* is marketed as a superfood with a wide range of health benefits (Abou-Zaid & Nadir, 2014).

## **Cultivation and Agricultural Practices**

*Moringa oleifera* is a hardy and resilient plant that can be cultivated in a wide range of environments. However, to maximize yield and nutritional content, it is important to follow specific agricultural practices that optimize growing conditions.

### **Ideal Growing Conditions**

*Moringa oleifera* thrives in tropical and subtropical regions with temperatures between 25–35°C. The plant prefers well-drained, sandy or loamy soils with a slightly acidic to slightly alkaline pH (Gopalakrishnan *et al.*, 2016). While *Moringa* can tolerate drought conditions, it requires adequate rainfall (250–3000 mm per year) for optimal growth and productivity (Dania *et al.*, 2014).

To ensure healthy growth, it is important to provide *Moringa* plants with sufficient sunlight, as they require full sun exposure for photosynthesis. In areas with low light conditions, growth may be stunted, and nutrient content may be reduced (Gopalakrishnan *et al.*, 2016).

## Propagation Methods

*Moringa oleifera* can be propagated from seeds or cuttings, depending on the desired characteristics of the plants. Seeds are typically used for propagation because they have high germination rates and produce plants with strong root systems (Sánchez-Machado *et al.*, 2010). Seeds should be sown at a depth of 2 cm in well-drained soil and kept moist until germination, which typically occurs within 5–12 days (Dania *et al.*, 2014).

Alternatively, *Moringa* can be propagated from cuttings, which involves planting a branch from an existing tree. Cuttings should be 1 m in length and 4–5 cm in diameter and should be planted in sandy or loamy soil. However, plants grown from cuttings may have weaker root systems and may be more vulnerable to drought and wind damage (Gopalakrishnan *et al.*, 2016).

## Soil and Nutrient Management

Soil quality is a critical factor in determining the yield and nutritional content of *Moringa oleifera*. Plants grown in nutrient-rich soils tend to have higher concentrations of proteins, vitamins, and minerals, while those grown in poor soils may have lower nutrient levels (Dania *et al.*, 2014).

To improve soil fertility, organic fertilizers such as compost or poultry manure can be applied to the soil. These fertilizers provide essential nutrients, such as nitrogen, phosphorus, and potassium, that support healthy plant growth and improve nutrient content (Dania *et al.*, 2014). In addition to organic fertilizers, inorganic fertilizers may also be used, but it is important to avoid over-application, as this can lead to nutrient imbalances and reduced plant health (Moyo *et al.*, 2011).

## Spacing and Pruning

Proper spacing and pruning are essential for maximizing the yield and quality of *Moringa* plants. In commercial cultivation, plants should be spaced 3–5 meters apart to allow for adequate sunlight and airflow. This spacing also makes it easier to manage the plants and harvest the leaves, seeds, and pods (Gopalakrishnan *et al.*, 2016).

Pruning is important for encouraging the growth of new shoots and increasing leaf production. It also helps prevent the plants from becoming too tall and difficult to manage. Pruning should be done regularly, starting when the plants are young, to promote a bushy growth habit and improve overall productivity (Moyo *et al.*, 2011).

Table 1. Nutritional value and medicinal properties of plants parts

Component	Nutritional Value	Medicinal Properties	Applications
Leaves	- Rich in Vitamins A, C, E	- Antioxidant	- Dietary supplement
	- High in Calcium, Iron, Protein	- Antidiabetic	- Fortification of foods
		- Anti-inflammatory	
Seeds	- High in Oleic Acid	- Anticancer	- Oil extraction (Ben oil)
	- Rich in Protein, Calcium, Potassium	- Antimicrobial	- Natural coagulant
		- Water purification agent	

<b>Pods</b>	- Good source of Dietary Fiber and Essential Fatty Acids	- Antimicrobial	- Used in food and beverages
		- Nutrient-rich food source	
<b>Flowers</b>	- Contains Nectar	- Antioxidant	- Traditional medicine
	- Rich in Calcium and Potassium	- Anti-inflammatory	- Nutritional supplements
<b>Roots &amp; Bark</b>	- Contains Alkaloids and Essential Minerals	- Antimicrobial	- Used in traditional remedies
		- Anti-inflammatory	
		- Digestive aid	

## Future Prospects and Research Directions

### Expanding Cultivation and Use

One of the most promising areas for future research is the expansion of *Moringa oleifera* cultivation in regions where it is not yet widely grown. This could help address food insecurity and malnutrition in vulnerable populations, particularly in areas with harsh climates where traditional crops may struggle to survive. Research into the best practices for cultivating *Moringa* in diverse environments, including studies on soil improvement techniques and climate resilience, could lead to more widespread adoption of this valuable crop (Gopalakrishnan *et al.*, 2016).

Additionally, expanding the use of *Moringa* in various industries, such as food processing, pharmaceuticals, and cosmetics, could create new economic opportunities for farmers and entrepreneurs. By developing new products that utilize *Moringa*'s nutritional and medicinal properties, businesses can tap into the growing demand for natural and sustainable ingredients. This could also promote greater awareness of *Moringa*'s benefits and encourage more people to incorporate it into their diets and daily routines (Abou-Zaid & Nadir, 2014).

### Enhancing Nutritional and Medicinal Properties

Another important area for future research is the enhancement of *Moringa oleifera*'s nutritional and medicinal properties through selective breeding and biotechnology. By identifying and promoting specific traits, such as higher concentrations of vitamins, minerals, or bioactive compounds, scientists can develop new varieties of *Moringa* that offer even greater health benefits. This could be particularly valuable in addressing specific nutrient deficiencies or targeting certain health conditions, such as diabetes or cancer (Tiloke *et al.*, 2013).

Biotechnology also offers the potential to enhance *Moringa*'s resistance to pests, diseases, and environmental stresses, making it an even more resilient crop. Genetic modification techniques could be used to introduce new traits or improve existing ones, potentially leading to the development of *Moringa* varieties that are better suited to different climates and growing conditions (Mbikay, 2012).

### Investigating Health Benefits

While *Moringa oleifera* has been shown to have a wide range of health benefits, there is still much to learn about the mechanisms behind these effects. Further research is needed to understand how *Moringa*'s bioactive compounds interact with the human body and contribute to disease prevention and treatment. For example, more studies are needed to determine the optimal dosages and delivery methods for using *Moringa* as a therapeutic agent for conditions such as diabetes, cancer, and cardiovascular disease (Cerf, 2013).

Clinical trials involving human participants are particularly important for validating the health claims associated with *Moringa*. While many studies have been conducted using animal models or in vitro experiments, there is a need for more research that directly examines the effects of *Moringa* on human health. This could help establish *Moringa* as a credible and effective treatment option in the medical community (Berkovich *et al.*, 2013).

### **Exploring Environmental and Socio-Economic Impacts**

As *Moringa oleifera* becomes more widely cultivated and utilized, it is important to consider the environmental and socio-economic impacts of its production. Research into sustainable farming practices, such as agroforestry and organic farming, can help ensure that *Moringa* is grown in a way that benefits both the environment and local communities. This includes studying the effects of *Moringa* cultivation on soil health, biodiversity, and water resources, as well as evaluating the potential for *Moringa* to improve food security and economic development in rural areas (Gopalakrishnan *et al.*, 2016).

Additionally, research into the market dynamics of *Moringa* products, including consumer demand and pricing, can help inform strategies for promoting *Moringa* as a viable and profitable crop. By understanding the economic factors that influence the *Moringa* industry, policymakers and stakeholders can develop initiatives that support smallholder farmers and encourage the growth of *Moringa* businesses (Abou-Zaid & Nadir, 2014).

### **Conclusion**

*Moringa oleifera* stands out as one of the most versatile and beneficial plants available, offering a wealth of nutritional, medicinal, and commercial opportunities. Its rich nutritional profile, including high levels of essential vitamins, minerals, and proteins, makes it an invaluable resource for combating malnutrition, particularly in regions with limited access to diverse food sources. Additionally, its wide range of medicinal properties, including antioxidant, anticancer, antidiabetic, anti-inflammatory, and antimicrobial effects, highlights its potential as a natural remedy for many chronic diseases.

Beyond its health benefits, *Moringa oleifera* is also a valuable resource for sustainable development. Its adaptability to different environmental conditions and its use in water purification, oil extraction, and food fortification make it a key player in efforts to address global challenges such as food security, clean water access, and economic development. The potential of *Moringa* to support rural livelihoods and promote sustainable agricultural practices further underscores its importance in both local and global contexts.

However, despite the extensive research already conducted on *Moringa oleifera*, there is still much to discover about this remarkable plant. Future research should focus on expanding *Moringa* cultivation, enhancing its nutritional and medicinal properties, investigating its health benefits, and exploring its environmental and socio-economic impacts. By continuing to study and promote *Moringa oleifera*, we can

unlock its full potential and harness its power to improve health, sustainability, and well-being around the world.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

#### **CONSENT**

It is not applicable.

#### **ETHICAL APPROVAL**

It is not applicable.

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

#### **References**

1. Gopalakrishnan, L., Doriya, K., & Kumar, D. S. (2016). *Moringa oleifera*: A review on nutritive importance and its medicinal application. *Food Science and Human Wellness*, 5(2), 49-56. <https://doi.org/10.1016/j.fshw.2016.04.001>
2. Mbikay, M. (2012). Therapeutic potential of *Moringa oleifera* leaves in chronic hyperglycemia and dyslipidemia: A review. *Frontiers in Pharmacology*, 3, 1-12. <https://doi.org/10.3389/fphar.2012.00024>
3. Fuglie, L. J. (2005). The *Moringa* Tree: A local solution to malnutrition. *Church World Service in Senegal*.

4. Yang, R., Chang, L., Hsu, J., Weng, B. B. C., Palada, C., Chadha, M. L., & Levasseur, V. (2006). Nutritional and functional properties of *Moringa* leaves from germplasm, to plant, to food, to health. *American Chemical Society*.
5. Chumark, P., Khunawat, P., Sanvarinda, Y., Phornchirasilp, S., Morales, N. P., Phivthongngam, L., ... & Pongrapeeporn, K. U. (2008). The in vitro and ex vivo antioxidant properties, hypolipidaemic and antiatherosclerotic activities of water extract of *Moringa oleifera* Lam. leaves. *Journal of Ethnopharmacology*, *116*(3), 439-446. <https://doi.org/10.1016/j.jep.2007.12.010>
6. Tiloke, C., Phulukdaree, A., & Chuturgoon, A. A. (2013). The antiproliferative effect of *Moringa oleifera* crude aqueous leaf extract on cancerous human alveolar epithelial cells. *BMC Complementary and Alternative Medicine*, *13*, 226-233. <https://doi.org/10.1186/1472-6882-13-226>
7. Lalas, S., & Tsaknis, J. (2002). Characterization of *Moringa oleifera* seed oil variety Periyakulam-1. *Journal of Food Composition and Analysis*, *15*(1), 65-77.
8. Sánchez-Machado, D. I., Núñez-Gastélum, J. A., Reyes-Moreno, C., Ramírez-Wong, B., & López-Cervantes, J. (2010). Nutritional quality of edible parts of *Moringa oleifera*. *Food Analytical Methods*, *3*(3), 175-180.
9. Dania, S. O., Akpansubi, P., & Eghagara, O. O. (2014). Comparative effects of different fertilizer sources on the growth and nutrient content of *Moringa* (*Moringa oleifera*) seedlings in a greenhouse trial. *Pharmaceutical and Clinical Research*, *5*(2), 67-72.
10. Moyo, B., Masika, P., Hugo, A., & Muchenje, V. (2011). Nutritional characterization of *Moringa* (*Moringa oleifera* Lam.) leaves, *African Journal of Biotechnology*, *10*(60), 12925-12933.
11. Hermawan, A., Nur, K. A., Sarmoko, D., Dewi, P., & Meiyanto, E. (2012). Ethanolic extract of *Moringa oleifera* increased cytotoxic effect of doxorubicin on HeLa cancer cells. *Journal of Natural Remedies*, *12*(2), 108-114.
12. Nakamura, Y., Kawakami, M., Yoshihiro, A., Miyoshi, N., Ohigashi, H., Kawai, K., et al. (2002). Involvement of the mitochondrial death pathway in chemopreventive benzyl isothiocyanate-induced apoptosis. *Journal of Biological Chemistry*, *277*(10), 8492-8499.
13. Miyoshi, N., Uchida, K., Osawa, T., & Nakamura, Y. (2004). A link between benzyl isothiocyanate-induced cell cycle arrest and apoptosis: involvement of mitogen-activated protein kinases in the Bcl-2 phosphorylation. *Cancer Research*, *64*(6), 2134-2142.
14. Liou, G. Y., & Storz, P. (2010). Reactive oxygen species in cancer. *Free Radical Research*, *44*(5), 479-496.
15. Cerf, M. E. (2013). Beta cell dysfunction and insulin resistance. *Frontiers in Endocrinology*, *4*, 1-12. <https://doi.org/10.3389/fendo.2013.00037>
16. Divi, S. M., Bellamkonda, R., & Dasireddy, S. K. (2012). Evaluation of antidiabetic and antihyperlipidemic potential of aqueous extract of *Moringa oleifera* in fructose-fed insulin-resistant and

STZ-induced diabetic Wistar rats: a comparative study. *Asian Journal of Pharmaceutical and Clinical Research*, 5(2), 67-72.

17. Mahajan, G. S., & Mehta, A. A. (2009). Anti-arthritis activity of hydroalcoholic extract of flowers of *Moringa oleifera* Lam. in Wistar rats. *Journal of Herbs, Spices & Medicinal Plants*, 15(2), 149-163.

18. Viera, G. H. F., Mourão, J. A., Ângelo, Â. M., Costa, R. A., & Vieira, R. H. S. F. (2010). Antibacterial effect (in vitro) of *Moringa oleifera* and *Annona muricata* against Gram-positive and Gram-negative bacteria. *Revista do Instituto de Medicina Tropical de São Paulo*, 52(3), 129-132.

19. Suhartini, S., Hidayat, N., & Nurhayati, O. D. (2011). The effect of powdered *Moringa oleifera* seeds and sand filtration in treating tapioca starch wastewater. *International Journal of Environmental Science and Development*, 2(1), 12-16.

20. Abou-Zaid, A. A., & Nadir, A. S. (2014). Quality evaluation of nutritious chocolate and halawa tahinia produced with *Moringa (Moringa oleifera)* leaves powder. *Middle East Journal of Applied Sciences*, 4(4), 1007-1015.

21. Owusu, D., Ellis, W. O., Oduro, I., & Afutu, E. (2014). Development of butter and cream crackers using *Moringa oleifera* and *Ipomoea batatas* flours. *Food Science & Nutrition*, 2(5), 464-470. <https://doi.org/10.1002/fsn3.120>

22. Lalas, S., & Tsaknis, J. (2002). Characterization of *Moringa oleifera* seed oil variety Periyakulam-1. *Journal of Food Composition and Analysis*, 15(1), 65-77. <https://doi.org/10.1006/jfca.2001.1049>

23. Moyo, B., Masika, P. J., Hugo, A., & Muchenje, V. (2011). Nutritional characterization of *Moringa (Moringa oleifera* Lam.) leaves. *African Journal of Biotechnology*, 10(60), 12925-12933.