

# Study The Ecological of TheBanggaiCardinalfish in the Sea Waters of BanggaiLaut Regency, Indonesia

## ABSTRACT

Indonesia is one of the countries with a high level of biodiversity. This is evidenced by the many species of fish and coral in Indonesian waters. The abundance of biodiversity also allows Indonesia to have endemic species, one of which is the BanggaiCardinalfish. BanggaiCardinalfish (*Pterapogonkauderni*) is one of the endemic species found in the Banggai Islands-Central Sulawesi-Indonesia. This study aims to see the bio-ecological conditions of the BanggaiCardinalfish. This study was conducted in January-March 2014 in North Banggai District, BanggaiLaut Regency, Central Sulawesi Province. The study was conducted using the observation method and monitoring the population of *Pterapogonkauderni* at predetermined stations and measuring the quality of marine waters at each station. Population monitoring was carried out using the Belt transect method. The data obtained were then analyzed using population density analysis. The water quality data obtained were then compared with the seawater quality standards according to PP No. 22 of 2021. Based on the research results, it is known that the population of BanggaiCardinalfish and associated biota from each location is generally low, only at station III which has the highest population. Water quality conditions are generally optimal only at salinity parameters that tend to be very high.

**Keywords:** Ecology; EndemicSpecies; BanggaiCardinalfish; Populastion; Water Quality

## 1. INTRODUCTION

Indonesia is part of *The Coral Triangle region*, which is a region that has a very high level of biodiversity and is the center of marine biota diversity [1]–[3]. This high level of biodiversity allows for endemic marine biota species only found in Indonesian waters. BanggaiCardinalfish (*Pterapogonkauderni* Koumans, 1933) is a species endemic to the Banggai Islands-Central Sulawesi and one of Indonesia's Endemic Coral Fish Species [4], [5]. This species has a distribution area of 5,000 km<sup>2</sup> with a potential habitat area of 20-24 km<sup>2</sup>, and its primary habitat is coastal areas containing coral reefs and seagrass beds with a depth of 0-5 m [6].

The current existence of *Pterapogonkauderni* has experienced a decline in population numbers due to human activities, such as both due to fishing, land use and so on. The United States and the European Union claim that the exploitation of BanggaiCardinalfish for export has been over-harvested and unsustainable [7]. In addition, there are concerns about the taking of associated biota from BanggaiCardinalfish by coastal communities, as a food source [8]. In 2016, 17th CITES CoP, Appedix II, categorized *Pterapogonkauderni* as an

**endangered species.** This decision requires Indonesia to ensure the preservation of this species by carrying out conservation and sustainable management [7]–[9] .

The Indonesian government, through the Decree of the Minister of Maritime Affairs and Fisheries of the Republic of Indonesia, has issued decision **Number 49/KEPMEN-KP/2018** concerning the Determination of the Limited Protection Status of the BanggaiCapungan Fish (*Pterapogonkauderni*). Along with the Decree of the Minister of Maritime Affairs and Fisheries Number 53/KEPMEN-KP/2019 concerning coastal conservation and the Banggai Small Islands, Banggai Sea, Banggai Islands and the surrounding waters of Central Sulawesi Province. To maintain the preservation of *Pterapogonkauderni* in its natural habit.

Based on monitoring in 2019, the population status of *Pterapogonkauderni* at eight monitoring locations in the Bokan Islands showed a decline in the population of *Pterapogonkauderni* and its associated biota (sea urchins and sea anemones) [10] . Based on the results of previous population monitoring, it is necessary to re-monitor the *Pterapogonkauderni* population in different locations in the Banggai Islands. The waters of North Banggai District, BanggaiLaut Regency, are one of the areas where *Pterapogonkauderni* lives, and have been designated by the Decree of the Minister of Maritime Affairs and Fisheries as part of the coastal and small island conservation area in Central Sulawesi Province. This area is also a beach tourist attraction in BanggaiLaut Regency.

The existence of aquatic biota populations cannot be separated from the quality conditions of the waters. Water quality can be broadly interpreted as physical and chemical factors that influence the life of fish and other aquatic organisms, either directly or indirectly [11] . Monitoring water quality aims to maintain ecosystems and aquatic habitats because decreased water quality quickly affects these two components[12] . Therefore, this research aims to determine the population density of *Pterapogonkauderni* and water quality in BanggaiUtara District, BanggaiLaut Regency.

## **2. MATERIALS AND METHODS**

### **2.1 TIME AND PLACE**

The research was carried out from January to March 2024 in Banggai Utara District, BanggaiLaut Regency, Central Sulawesi Province. With four data collection station points, namely; (1) Station I Popisi Village: E123°30.908' and S01°29.956'; (2) Station II Tolisetubono Village: E123°29.966' and S01°31.520'; (3) Station III Bonebaru Village: E123°29.643' and S01°31.916'; (4) Station IV Bonebaru/Bongo Village: E123°28.911' and S1°32.422'. This area is included in the conservation area designated by the KKP in PERMEN KP Number 53/KEPMEN-KP/2019. The data collection location can be seen in Figure 1.




Figure 1. Research Location

## 2.2 TOOLS AND MATERIALS

The tools and materials used in this study were Stationery for data recording, TR-LE-731 Portable for collecting Temperature, pH and DO data, Refracto meter for measuring Salinity, Meter Roll for transect measurements.

## 2.3 RESEARCH METHODS

The research was conducted using survey methods and direct observation at predetermined locations. Data was collection by monitoring the population number of *Pterpogonkauderniat* predetermined stations and measuring the quality of sea waters at each station. The technique for collecting data on population numbers and water quality can be seen below.

- a. Number of Populations of *Pterpogonkaudernirefers* to [15]
  - 1) Observations were carried out by placing transects parallel to the coastline at a depth of 1.5-3 m where BanggaiCardinalfish are generally found. However, it will have to be adjusted to tidal conditions.
  - 2) Observations were made using the *belt transect method*. The size of the transect is 20 m long and 5 m wide (transect area **100 m<sup>2</sup>**), namely 2.5 m wide on the right and 2.5 m on the left (figure 2).
  - 3) Two divers collected data, each dividing the task of collectingfish data. Each on the left and right of the transect.
  - 4) Each sampling station was conducted 6 times. The BanggaiCardinalfish found were recorded based on the fish size category(Recruit < 1,8 cm, Juvenile 1,8 to 3,5 cm , Adult>3,5 cm).




Figure 2. Population survey of *Pterapogon kauderni* using the belt transect method [15]

b. Water quality

- 1) Water quality measurement is carried out in-situ including temperature parameters (°C), pH, DO (ppm), salinity (ppt).
- 2) Sampling is carried out once for each parameter. Water sampling is carried out in a composite manner between the bottom of the water and 30 cm from the water surface.

## 2.4 DATA ANALYSIS

Data from the *Pterapogon kauderni* population survey and associated Biota were analyzed to determine the density of fish at each location and size category using a density formula based on the following monitoring guidelines from the Ministry of Marine Affairs and Fisheries:

$$\text{Density Population (individu/m}^2\text{)} = \frac{\text{total fish individuals}}{100} \text{ [15]}$$

Water quality measurement data is analyzed by comparing with quality standards based on PP No. 22 of 2021 concerning sea water quality standards.

## 3 . RESULTS AND DISCUSSION

### 3.1 Population Conditions of *Pterapogon kauderni*

Figure 3 presents the *Pterapogon kauderni* population survey results. The most numerous *Pterapogon kauderni* population is at station III, which is dominated by the very high presence of Juveniles, reaching 12.60 Individuals/m<sup>2</sup>, then Adults 2.18 Individuals/m<sup>2</sup> and Recruitment 1.14 Individuals/m<sup>2</sup>.




Figure 3. Population Density of *Pteropogonkauderni*

The high population at station III is likely due to the many associated biota and coral reefs there. *Pteropogonkauderni* can be found in seagrass and coral reef ecosystems, including fringing reefs and reef flats [7]. Coral reefs provide a place for fish to shelter, spawn and feed.

The lowest population density was found at station II, with the number of recruit 0.06 individuals/m<sup>2</sup>, juveniles 0.44 individuals/m<sup>2</sup>, and adults 0.96 individuals/m<sup>2</sup>. The low population of *Pteropogonkauderni* at station II is likely due to the lack of associated biota at that location. The absence of an ecosystem supporting this species and its associated biota can affect its population density. *Pteropogonkauderni* does not have a self-defense mechanism, so it needs associated biota to avoid predators. It is known that for most of its life, it coexists with biota; its leading associations are sea urchins, soft corals, and sea anemones [7], [10], [14].

### 3.2 Condition of Association Biota




Figure 4. Association Biota Population

Based on Figure 4. The population of associated biota (sea urchins and anemones) is most abundant at station III, with a density of sea urchins of 0.91 individuals/m<sup>2</sup> and anemones of 0.08 individuals/m<sup>2</sup>. Meanwhile, the lowest was at station IV with a density of both associated biota, namely 0.01 individuals/m<sup>2</sup>.

It is strongly suspected that the associated biota at Station IV exists because it is captured for consumption by the local community. Tourist activities can also be the cause of the reduction in sea urchins in the area. As is known, this area is a tourist spot for local people on Banggai Island. The loss of associated biota is caused by habitat degradation caused by human activities and climate change [7], [10].

### 3.4 Water Quality

Water quality conditions at each research location can be seen in Table 2.

Table 2. Water quality at the research location

Parameter	Location				Quality Standards (PP No. 22 of 2021)	Note.
	Station I	Station II	Station III	Station IV		
Temperature (°C)	29.8	29.9	29.3	31.8	28 to 32 °C	Optimal
pH	8.06	8.08	8.01	8.19	7 to 8.5	Optimal
DO (ppm)	7.9	11.04	8.1	11.04	> 5 ppm	Optimal
Salinity (ppt)	35	34	41	36	33 to 34 ppm	Bad

Based on the research results, it is known that the water quality conditions for temperature, pH, and DO at each station are still optimal for seawater biota. The water temperature at the research location ranges from 29.3 to 31.8 °C. Optimal water conditions will have a good impact on marine biota. These conditions will support the metabolic processes of fish so that they can support growth, survival and reproduction.

The pH of the water ranges between 8.01 to 8.19, which is still in optimal conditions. Water pH can affect the physiology of marine biota, including inhibiting growth, biota will be very sensitive to bacteria and parasites, and the water will be toxic to fish. Changes in pH that are very acidic or alkaline will disrupt the survival of aquatic organisms because they disrupt the respiration process [15].

The DO condition of the research location waters ranges from 7.9 to 11.04 ppm, which is very good. The DO of waters greatly determines the survival of marine biota. Water conditions that contain dissolved oxygen. Oxygen levels can affect the decomposition, reproduction and growth processes. The need for dissolved oxygen in fish is influenced by age, activity and water conditions. The oxygen content influences organic and inorganic materials' oxidation and reduction processes [16].

Salinity at the research location ranges from 34 to 41 ppt, this value is quite high and can be dangerous for the existence of marine biota. Seawater salinity influences the distribution, abundance and growth of aquatic biota as well as their density in a body of water [17]. Suboptimal salinity can cause osmoregulation disorders for biota. The water's high salinity is likely due to the very high evaporation level. The cause of the high salinity is also thought to be due to the data collection being carried out in sunny and cloudless conditions. One of the factors that influences salinity is evaporation. The greater the level of seawater evaporation in an area, the higher the salinity, and vice versa in areas with low levels of seawater evaporation [18].

#### 4. CONCLUSION

[Based on the research results, it is known that the population of Banggai Cardinalfish from each location is generally low, only at station III which has the highest population. The population of associated biota is also low, only at station III which has a high population. Water quality conditions are generally optimal only when the salinity parameter tends to be very high.]

Efforts to carry out in-situ conservation need to pay attention to environmental conditions, especially salinity, so that conservation can be carried out optimally.]

#### Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

#### REFERENCES

- [1] A. D. B. ADS, *State of the Coral Triangle: Indonesia*. Philippines: Asian Development Bank, 2014.
- [2] S. J. Peterson and T. A. Conant, "Status Review Report Banggai Cardinalfish, *Pterapogon kauderni*," *Natl. Mar. Fish. Serv. Natl. Ocean. Atmos. Adm.*, no. November, 2015.
- [3] D. I. K. Mujiono and J. Oktaviani, "Segitiga Terumbu Karang Dunia (the Coral Triangle): Manfaat, Masalah Dan Upaya," *J. Din. Glob.*, vol. 6, no. 01, pp. 1–19,

2021, doi: 10.36859/jdg.v6i01.405.

- [4] Hasim, M. R. F. H. Sunani, and Mulis, "Interaction of salinity and natural feed on growth and survival rate of the banggai cardinalfish (*Pterapogon kauderni*) for ex-situ conservation development," *Egypt. J. Aquat. Biol. Fish.*, vol. 25, no. 5, pp. 241–252, 2021, doi: 10.21608/ejabf.2021.198591.
- [5] A. Moore, S. Ndobe, A. I. M. Salanggon, and D. Wahyudi, "Biodiversitas dan Isu-Isu Pengelolaan Coral Reefs of Sonit Island , Banggai Archipelago : Biodiversity and Management Issues," in *Simposium Nasional Kelautan dan Perikanan, Makassar II, 9 Mei 2015*, 2015, pp. 1–8.
- [6] S. Ndobe, I. Yasir, A. M. Moore, M. V Biondo, and S. J. Foster, "A study to assess the impact of international trade on the conservation status of *Pterapogon kauderni* (Banggai cardinalfish)," *This Inf. Doc. has been Submitt. by Secr. Req. Int. Union Conserv. Nat. Relat. to agenda item 21\**, p. 65, 2018, [Online]. Available: <https://cites.org/sites/default/files/eng/com/ac/30/Inf/E-AC30-Inf-16.pdf>.
- [7] A. Mardiasuti and R. T. Soehartono, "BANGGAI CARDINALFISH (*Pterapogon kauderni*) AND THE ATTEMPT FOR THE INCLUSION OF CITES APPENDIX II," *J. Ilmu dan Teknol. Kelaut. Trop.*, vol. 12, no. 2, pp. 597–607, 2020, doi: 10.29244/jitkt.v12i2.30944.
- [8] A. Moore, S. Ndobe, A.-I. Salanggon, Ederyan, and A. Rahman, "Banggai Cardinalfish Ornamental Fishery: The Importance of Microhabitat," *Proc. 12th Int. Coral Reef Symp. Cairns, Aust. 9-13 July 2012*, no. July 2012, p. 13C, 2012, [Online]. Available: [http://www.icrs2012.com/proceedings/manuscripts/ICRS2012\\_13C\\_1.pdf](http://www.icrs2012.com/proceedings/manuscripts/ICRS2012_13C_1.pdf).
- [9] U. Y. Arbi, S. Ndobe, and Dirhamsyah, *Ikan Capungan Banggai (Pterapogon kauderni) sebuah catatan bioekologi dan introduksi*. Jakarta: Balai Pustaka, 2022.
- [10] N. Pompon, S. Ndobe, K. Mansyur, and M. Tis'In, "Growth and survival of juvenile Banggai cardinalfish (*Pterapogon kauderni*) reared under different salinities in recirculating aquaria equipped with protein skimmers," *IOP Conf. Ser. Earth Environ. Sci.*, vol. 253, no. 1, 2019, doi: 10.1088/1755-1315/253/1/012002.
- [11] N. N. Wiadnyana *et al.*, "Population trends of Banggai cardinalfish in the Banggai Islands, Central Sulawesi, Indonesia," *IOP Conf. Ser. Earth Environ. Sci.*, vol. 420, no. 1, 2020, doi: 10.1088/1755-1315/420/1/012033.
- [12] S. Ndobe *et al.*, "Monitoring the endemic ornamental fish *Pterapogon kauderni* in Bokan Kepulauan, Banggai marine protected area, Indonesia," *Depik*, vol. 9, no. 1, pp. 18–31, 2020, doi: 10.13170/depik.9.1.15363.
- [13] C. Megawati, M. Yusuf, and L. Maslukah, "Sebaran Kualitas Perairan Ditinjau dari Zat Hara, Oksigen Terlarut dan pH di Perairan Selat Bali Bagian Selatan," *J. Oseanografi*, vol. 3, no. 2, pp. 142–150, 2014, [Online]. Available: <http://ejournal-s1.undip.ac.id/index.php/jose.50275Telp/Fax>.
- [14] N. L. G. R. A. Saraswati, I. W. Arthana, and I. G. Hendrawan, "Analisis Kualitas Perairan Pada Wilayah Perairan Pulau Serangan Bagian Utara Berdasarkan Baku Mutu Air Laut," *J. Mar. Aquat. Sci.*, vol. 3, no. 2, p. 163, 2017, doi: 10.24843/jmas.2017.v3.i02.163-170.

- [15] D. K. dan K. H. L. DKKHL, "Pedoman Monitoring Banggai CardinalFish (Pterapogon kauderni)." Kementerian Kelautan dan Perikanan, Jakarta, 2019.
- [16] U. Y. Arbi and A. Faricha, "New host record of microhabitat preferences of the Banggai cardinalfish (Pterapogon kauderni) in the introduced habitat in Luwuk waters, Sulawesi," *IOP Conf. Ser. Earth Environ. Sci.*, vol. 944, no. 1, 2021, doi: 10.1088/1755-1315/944/1/012018.
- [17] D. Irawan, S. P. Sari, E. Prasetyono, and A. F. Syarif, "GROWTH PERFORMANCE AND SURVIVAL RATE OF BRILLIANT RASBORA (*Rasbora einthovenii*) AT DIFFERENT PH TREATMENTS," *J. Aquatropica Asia*, vol. 4, no. 2, pp. 15–21, 2019, doi: 10.33019/aquatropica.v4i2.2221.
- [18] L. Riadhi, M. Rivai, and F. Budiman, "Pengaturan Oksigen Terlarut Menggunakan," *J. Tek. ITS*, vol. 6, no. 2, pp. 5–9, 2017.
- [19] T. M. Faisal, S. Bahri, A. Putriningtias, and A. Harahap, "Kualitas perairan di daerah pesisir Pulau Ujung Perling, Kota Langsa, Aceh," *Habitus Aquat.*, vol. 2, no. 2, pp. 95–99, 2022, doi: 10.29244/haj.2.1.95.
- [20] D. N. Rema and K. Umroh, "Analysis Pollution of Coastal Water in Bedukang, Deniang Village, Bangka Regency," *J. Trop. Mar. Sci.*, vol. 2, no. April, pp. 1–10, 2019.