

Investigating the Impact of Sprouting on the Nutritional and Anti-Nutritional Properties of *Ogi* Made from Yellow and White Maize Varieties

Abstract

Ogi, a popular traditional fermented maize gruel in Africa, is a staple food for millions of people for its nutritional value and versatility in culinary applications. This study investigated the effects of sprouting and fermentation on the nutritive quality of *ogi* produced from yellow and white maize varieties using standard methods analysis. The results showed significant improvements in the nutritional profile of sprouted *ogi* samples compared to unsprouted ones. The pro-vitamin A ranged from 26.17 ± 0.02 to 31.24 ± 0.03 in unsprouted yellow *ogi* (UYO and SYO) respectively. The corresponding values for the unsprouted white *ogi* (UWO) and sprouted white *ogi* (SWO) are 9.28 ± 0.01 to 11.81 ± 0.03 . The protein content of SYO and SWO are 13.73 ± 0.07 and 12.02 ± 0.06 respectively compared to UYO and UWO samples (12.33 ± 0.56 and 10.22 ± 0.07). Sample SYO has the highest vitamin A content of $31.24 \mu\text{g}/100\text{g}$ followed by sample UYO which indicated higher amount of vitamin A content in yellow maize products (Samples UYO and SYO) could be attributed to the maize variety (yellow) containing higher amount of carotenoid compound, a pro-vitamin A. Notably, yellow maize *ogi* samples exhibited superior nutritional quality compared to white maize *ogi* samples. Sprouting and fermentation enhanced the bioavailability of nutrients, making *ogi* a more nutritious and wholesome food option. This study highlights the potential of simple processing techniques to improve the nutritional value of traditional foods, contributing to better nutrition and health outcomes in Africa.

Keywords: *Ogi*, sprouting, fermentation, nutritive quality, yellow maize, white maize.

1. Introduction

Cereal-based food, such as *ogi*, plays a vital role in the diets of millions worldwide, particularly in regions where maize is a staple crop. *Ogi*, traditional fermented cereal gruel, is popular across Africa for its nutritional value and versatility in culinary applications. *Ogi* is a popular fermented semi-solid food of various colour depending on the maize types (white or yellow) used in its preparation¹. Maize (*Zea mays* L.) is one of the most widely consumed cereal crops in Nigeria. It is rich in carbohydrate, supplying over 360 kcal/100 g of calories and contains adequate quantity of protein. It can be cooked, boiled, roasted, made into gruel for *Ogi* and useful in the production of animal feeds¹.

Ogi is the most popular fermented health food and its consumption cuts across the various segments of the society, including all age groups in many West African countries². The three major tribes in Nigeria refer to the fermented gruel (pap) by their different local names: Yourba (*ogi*), Igbo (*akamu*), and Hausa (*koko*). The popularity of this food product is mainly due to its availability, high digestibility and the fact that it supplies the necessary calories required³. However, research has shown that a diet based on maize may be deficient in micro-nutrients such as potassium, iron, B complex vitamins, amongst others²⁻³. It was reported that 100 g of maize foods consumed could to some extent, contribute in satisfying dietary reference intakes of children and women in energy, proteins, carbohydrates, magnesium, zinc, vitamins B1 and B6. However, it provides very low supply of fats, calcium, sodium, selenium, vitamins C, A and E⁴. The processing of maize ultimately increases its shelf life but, a significant loss of micronutrients may occur via heat degradation and/or leaching during processing and this may be associated with under-nutrition and hidden hunger⁵.

Sprouting and fermentation are both complex biological processes involving many biochemical and physiological reactions, result to significant changes in the nutrient composition and physicochemical properties of grains. Controlled sprouting and fermentation processes can lead to a final product with improved nutritional value, sensor properties, safety, stability and functional qualities⁶⁻⁷. While previous research has explored the effects of sprouting on various grains, including maize, there is paucity of information on the impact of sprouting on the nutritional and ant-nutritional properties of seed based food especially *ogi* made from different variety of maize. Therefore, the aim of this study is to conduct a comparative analysis of the effect of sprouting on the nutritional and anti-nutritional properties of *ogi* produced from yellow and white maize varieties. By systematically evaluating changes in nutrient content, bioavailability, and anti-nutrient levels following sprouting and fermentation, the study seek to provide valuable insights into the potential of sprouting as a sustainable approach to enhance the nutritional value of *ogi*.

2. Methodology

2.1 Preparation of Maize Grains to *Ogi* Powder

Yellow and white sweet maize grains obtained for the study were sorted and divided into two portions. The first portion was soaked for 24 hours after which the water was decanted and the grains were spread on trays lined with cloth and kept wet by frequent spraying of water at every morning and evening for 2 days. The sprouted maize grains were thereafter re-soaked and washed with clean water. The second portion was steeped in clean water at room temperature for 48 h. The water was decanted from the two portions and the fermented grains were washed with clean water and wet milled using an attrition mill. The wet bran was removed with a muslin cloth and the filtrate was allowed to settle for 24 h to form the starchy sediment *ogi* slurry⁸. The sediment was de-watered in a jute sack using a hydraulic jack. The de-watered mash was pulverized in a granulating machine, dried in a cabinet dryer (55±5°C), and dry milled to pass a mesh sieve of 0.5 mm, as reported⁴. The *ogi* powder thus prepared were labeled as follows: UYO: Unsprouted Yellow Maize *Ogi*, SYO: Sprouted Yellow Maize *Ogi*, UWO: Unsprouted White Maize *Ogi* and SWO: Sprouted White Maize *Ogi* respectively.



Figure 1: Pictures of Yellow (A) and White (B) Maize variety

2.2 Determination of proximate composition

The proximate composition of the samples was determined using the standard methods of the Association of Official Analytical Chemists international⁹. Crude protein was determined using micro-Kjeldahl method (Method No 988.05), crude fat was determined by the Soxhlet extraction method (Method No 920.39) and moisture content was determined using air oven at 105°C to a constant weight according to Method No 991.02 (Plus 11 SanyoGallenkamp UK). The ash content was determined by ashing the sample in a muffle furnace (Gallenkamp 3) at 550°C for 4 hours (Method No 935.13), crude fibre was determined by (Method No 978.10). The total carbohydrate was determined by difference according to Equation:

$$\text{Total carbohydrate} = 100 - (\% \text{ moisture} + \% \text{ fat} + \% \text{ protein} + \% \text{ fibre} + \% \text{ ash})$$



Figure 2: Pictures of Yellow ogi (C) and white ogi (D) powder

2.3 Pro-vitamin A: Carotenoids Analysis

Approximately 15 g of each sample added to 3 g of *Celite 454* (Tedia, Ohio, USA), were weighed and successive additions of 25 ml of acetone were added to obtain a paste, which was then transferred to a sintered funnel (5 µm) coupled to a 250 ml

Buchner flask and filtered under vacuum. This procedure was repeated three times until the sample became colourless after which the extract was transferred to a 500 ml separation funnel containing 40 ml of petroleum ether. The acetone was removed by the slow addition of ultrapure water (*Millipore*) to prevent emulsion formation. The aqueous phase was discarded and this procedure was repeated four times until no residual solvent remained (ultrapure water). The extract was then transferred through a funnel containing 15 g of anhydrous sodium sulphate (drying agent) and made up a volume of 50 ml with petroleum ether. Additionally, for the identification and quantification of the pro-vitamin A (carotenoid), 2 ml was removed from the extract and dried in an amber flask under nitrogen flow. The sample was diluted in 100 μ l of acetone under shaking in a vortex mixer (*Genie 2Scientific Industries*) and transferred to a 2-ml amber flask for high-performance liquid chromatography (HPLC) analysis. The concentration of the pro-vitamin A carotenoid was determined using the equation reported¹⁰.

$$\text{Carotenoid} \left(\frac{\mu\text{g}}{\text{g}} \right) = \frac{Ax * Cs (\mu\text{g ml}) * V \text{ ml}}{Ax * P \text{ g}}$$

Where Ax = carotenoid peak area, Cs = standard concentration, As = standard area, V = total extract volume, and P = sample weight.

2.4 Determination of Anti-nutrients

2.4.1 Oxalate Determination

This determination involved three major steps: digestion, oxalate precipitation, and permanganate titration.

- a. *Digestion*: Digestion was effected by suspending 2 g of the sample in 190 mL of distilled water in a 250 mL volumetric flask after which 10 mL of 6M HCl was added and the suspension digested at 100 °C for 1 hour. The solution was cooled, and made up to 250 mL mark in the flask before filtration.
- b. *Precipitation*: Duplicate portions of the filtrate were measured into breakers and four drops of methyl red indicator added. Ammonia solution was added in drops until the test solution changed from pink to faint yellow colour (pH 4.0-4.5). Each portion was then heated to 90 °C, cooled and then filtered to remove the precipitate containing ferrous ion. The filtrate was again heated to 90 °C and 10ml of 5% CaCl₂ solution was added while being stirred constantly. The solution was then heated and left overnight at 25 °C, and centrifuged at 2000rpm for 5minutes. The supernatant was decanted and the precipitate completely dissolved in 20ml of 25% (v/v) H₂SO₄ solution and filtered.
- c. *Titration*: The total filtrate resulting from the digestion of 2 g of sample was made up to a volume of 300 mL. Aliquots of 125 mL of the filtrate was heated until near boiling and then titrated against 0.02 M standardized KMnO₄ solution to a faint pink color which persisted for 30 seconds. The calcium oxalate content was calculated using the formula:

$$\text{Oxalate Content} = \frac{T \times V_{me} \times D_f}{M_e \times M_f} \left(\frac{\text{Mg}}{100\text{g}} \right)$$

Where T is the titre of KMnO_4 (mL), V_{me} is the volume-mass equivalent, D_f is the Dilution factor = V_t/A Where V_t is the total volume of filtrate (300 mL) and A is the aliquot used i.e. 125 mL, M_e is the molar equivalent of KMnO_4 in oxalate and M_f is the mass of sample used.

2.4.2 Phytate Determination

The sample (5 mL) was mixed, cured for 5 hrs and filtered. Aliquots of 250 mL of the filtrate in a conical flask was added to 5.00 mL of 0.30 % ammonium thiocyanate, the mixture was titrated with standard iron (III) chloride solution to a persistent brownish yellow coloration that persisted for 4 mins. The amount of phytates was calculated with the equation below:

$$\text{Phytic acid} = \frac{\text{Titre value} \times 0.00195 \times 1.9 \times 100}{2}$$

2.4.3 Determination of Trypsin Inhibitors

The determination of trypsin inhibitor was carried out according to the procedure outlined by *Kakade and Evans* (1965). This involves weighing of 0.2g of the samples into a screw cap centrifuge tube. 10ml of 0.1M phosphate buffer was added and the contents shaken at room temperature for 1hour on a UDY shaker. The suspension obtained was centrifuged at 5000 rpm for 5minutes and filtered through *Whatman No. 42* filter paper. The volume of each was adjusted to 2ml with phosphate buffer. The test tube was placed in water bath, maintained at 37°C. Six millilitres of 5% TCA solution was added to one of the tubes to serve as a blank. 2ml of casein solution was added to all the tubes, which were previously kept at 37 °C. These were incubated for 20 mins. The reaction was stopped after 20 mins by adding 6ml of TCA solution to the experimental tubes and the tubes were shaken. The reaction was allowed to proceed for 1hour at room temperature. The mixture was filtered through *Whatman No. 42* filter paper. Absorbance of filtered sample and trypsin standard solutions was read at 280 nm. The trypsin inhibitor in mg/g was calculated using the formula:

$$\text{T. I. mg/g} = \frac{A_{\text{standard}} - A_{\text{sample}} \times \text{Dilution factor}}{0.1\text{g} \times \text{sample wt in g} \times 1000 \times \text{sample size}}$$

2.5 Mineral analysis

The ash of each sample (obtained in a crucible, by igniting 2 g of the samples in a muffle furnace) was digested by adding 5 ml of 2 M HCl, heated slowly for 20 minutes. After heating, it was filtered into a 100 ml volumetric flask. The filtrate was used for the determination of the concentration of each mineral element (potassium, sodium and iron while phosphorus was determined by molbdate method, a colorimetric technique). Buck Scientific - 200 Atomic Absorption Spectrophotometer -AAS (Buck Scientific,

Norwalk) was used. The composition of minerals was determined according to the standard method of the Association of Official Analytical Chemists⁹.

3.0 Results and Discussion

Table 1: Proximate Analysis of Ogi Samples

| Sample | Moisture (%) | Crude protein (%) | Crude ash (%) | Crude fat (%) | Crude fibre (%) | Carbohydrate (%) |
|------------|-------------------------|--------------------------|-------------------------|--------------------------|-------------------------|--------------------------|
| UYO | 3.49 ^a ±0.02 | 12.33 ^a ±0.56 | 2.86 ^a ±0.02 | 12.33 ^a ±0.01 | 1.55 ^b ±0.02 | 67.24 ^a ±0.09 |
| SYO | 5.61 ^a ±0.02 | 13.73 ^b ±0.07 | 3.94 ^a ±0.01 | 12.51 ^b ±0.02 | 1.44 ^a ±0.01 | 64.21 ^b ±0.12 |
| SWO | 4.22 ^b ±0.04 | 12.02 ^c ±0.06 | 2.69 ^b ±0.02 | 11.16 ^c ±0.01 | 1.42 ^c ±0.02 | 68.46 ^c ±0.05 |
| UWO | 5.45 ^c ±0.04 | 10.22 ^d ±0.07 | 3.64 ^b ±0.01 | 10.37 ^d ±0.02 | 1.37 ^c ±0.02 | 68.95 ^a ±0.19 |

Values are represented as means of triplicate determinations ± SD. Means with the same superscript within the same column indicate no significant difference ($p < 0.05$).

Proximate composition of the enriched and control samples

The results of the proximate analysis of the samples presented in Table 1 shows that the crude protein content of sample **SYO** (sprouted yellow ogi) (13.73 %) was significantly higher ($p < 0.05$) than that of the other samples while **UWO** sample (unsprouted white maize ogi) has the least amount of crude protein (10.22 %). This same trend was observed for the crude fat. The carbohydrates content of samples **SWO** and **UWO** are respectively 68.46% and 68.95%, which significantly higher than the samples **UYO** and **SYO** respectively. This is in line with the notion that yellow maize type has higher protein and lower carbohydrate content than the white maize of powdered ogi⁹.

The ash content of the sprouted ogi samples (**SYO** and **SWO**) were higher than the unsprouted ogi samples (**UYO** and **UWO**). This could be due to the endogenous enzymes hydrolyzing the complex carbohydrate to release fibre. The **SYO** sample had the highest ash content (3.94 %) and the least carbohydrate content (64.21 %) while the sample **UYO** had the ash content of 2.86 % and the carbohydrate content of 67.24 % which is lower than the value recorded for **UWO** (68.95). The findings of the present study clearly showed that the higher the protein content of the sample, the lower the carbohydrate content of the sample. This trend is in line with the previous studies¹¹⁻¹²

Table2: Pro-vitamin A (Carotenoid) and Calcium Analysis

| SAMPLE | PROVITAMIN A | CALCIUM (mg/100g) |
|--------|--------------------------|-------------------------|
| UYO | 26.17 ^b ±0.02 | 7.22 ^b ±0.02 |
| SYO | 31.24 ^a ±0.03 | 9.74 ^a ±0.01 |
| SWO | 11.81 ^c ±0.03 | 6.38 ^c ±0.01 |
| UWO | 9.28 ^d ±0.01 | 5.77 ^d ±0.02 |

The results revealed that sample SYO (Sprouted yellow *ogi*) has the highest vitamin A content of 31.24 µg/100g followed by sample UYO (unsprouted yellow *ogi*). The higher amount of vitamin A content in yellow maize products (Samples UYO and SYO) could be attributed to the maize variety (yellow) containing higher amount of carotenoid compound, a pro-vitamin A. The corresponding low content of pro-vitamin A in sample SWO and UWO respectively can be attributed to the lower carotenoid content of the maize from which the products (*ogi*) were made. Vitamins are major micronutrients needed in our body system to carry out normal body physiological functions. Vitamin A is important for eye functioning (vision)¹³.

The sample trend was observed in the result of calcium content of the samples investigated. Samples SYO and SWO which are made from the sprouted maize (yellow and white) have the higher amount of calcium content (9.74 and 6.38 g/100g respectively). These observations reinforced the assertion that processing method employed impact on the nutritional quality of the products. The slightly higher calcium content noted in *ogi* made from sprouted white maize could be as a result of enzymatic action that occurred during the sprouting process¹⁴.

Table 3: Mineral content of Sprouted and Unsprouted *ogi* from yellow and white maize variety

| Samples | Na | K | Ca | Mg | P |
|---------|----------|-----------|-----------|------------|-----------|
| UYO | 6.9±0.02 | 15.5±0.05 | 7.22±0.01 | 1.26±0.01 | 0.81±0.02 |
| SYO | 1.3±0.01 | 13.8±0.06 | 9.74±0.01 | 9.52±0.02 | 0.65±0.02 |
| SWO | 7.1±0.02 | 14.4±0.05 | 6.38±0.01 | 9.18±0.02 | 0.57±0.01 |
| UWO | 6.8±0.01 | 15.9±0.06 | 5.77±0.02 | 10.39±0.02 | 0.71±0.02 |

The results of the mineral composition of the is presented in Figure 1. The sodium content ranged from 6.81 ± 0.23 in UWO to 11.31±0.02 in SYO. The same trends were observed for potassium, calcium and magnesium. Generally, it was observed that mineral content of unsprouted samples were higher than the sprouted samples. Some of these minerals must have being used

up during enzymatic reactions taken placed during sprouting leading to decrease in concentrations of theses minerals. The significant higher concentration of phosphorous in unsprouted samples is an indication of bioavailability of these minerals to the consumers of the products (Ogi). Relatively higher concentrations of ant nutrients: the phytates, tannins and trypsin inhibitors in the unsprouted samples indicate that the available minerals in the food substance might not be bioavailable to the body¹⁵. Phytate is a stores form of phosphorus in seeds and has strong ability to complex with mineral elements thus making the elements non bioavailable to the animal. Therefore, the significant reductions of these ant-nutrients in the sprouted yellow and white maize samples is clear indication that the available minerals will be readily available to the consumers of the products made from the sprouted maize variety and




Figure3: Anti-nutrients in Ogi samples

Conclusion

The study revealed that the quality of food products depends on the quality of raw materials used in its preparation as well as the processing method adopted. Yellow maize contains higher nutritional quality than the white maize species. However, sprouting, an enzymatic biochemical reaction improved the nutritional quality of *ogi* irrespective of the maize types employed. Sprouting method should therefore be adopted in the processing of *ogi*, a common staple food in the west Africa. The sprouting will reduce the anti-nutritive factors in the seeds and thereby improve the bioavailability of the mineral elements present in the food formulation.

References

- [1] Rouf-Shah, T. Prasad, K. Kumar, P (2016). Maize-a potential source of human nutrition and health: a review, *Cogent. Food Agric.* 2(1) (2016) 1166995. <https://doi.org/10.1080/23311932.2016.1166995>.
- [2] Adejuwon, K.P., Osundahunsi, O. F. Akinola, S.A. (2021). Effect of fermentation on nutritional quality, growth and hematological parameters of rats fed sorghum-soybean-orange flesh sweet potato complementary diet, *Food Sci.*

- Nutri. 9(2) (2021) 639-650. <https://doi.org/10.1002/fsn3.2013>.
- [3] Akin-Osanaiye, B. C., & Oladele, T. F. (2017). Evaluation of the microorganisms associated with untreated and salt-treated ogi (corn caramel) stored at room temperature. *Direct Research Journal of Biology and Biotechnology*, 3, 60-66.
- [4] Hussain, M. Saeed, F. Niaz, F.B. et al. (2021). Anjum, Biochemical and nutritional profile of maize bran-enriched flour in relation to its end-use quality, *Food Sci. Nutri.* 9(6) (2021) 3336-3345. <https://doi.org/10.1080/10942912.2021.1973026>.
- [5] Adelekan, A.O., Alamu, E.A. Daramola, B.E (2021). Effect of enrichment with turmeric and ginger on some quality characteristics of fermented maize Ogi, *Croat. J. Food Sci. Technol.* 13(2) (2021) 210-220. <https://doi.org/10.17508/CJFST.2021.13.2.11>.
- [6] Galani Y.J.H., Orfila C., Gong Y. Y (2022). A review of micronutrient deficiencies and analysis of maize contribution to nutrient requirements of women and children in Eastern and Southern Africa: Critical Review, *Food Science and Nutrition*.:62(6):1568-1591. doi: 10.1080/10408398.2020.1844636.
- [7] Abdulsudi Issa-Z. and Gudila B. M (2024). Compliance Level and Stability of Micronutrients in Fortified Maize Flour in Tanzania. *Hindawi International Journal of Food Science Article ID 7746750, 11 pages* <https://doi.org/10.1155/2024/7746750>
- [8] Fenfeng Wu and Xueming Xu (2019), 7-Sprouted grain-based fermented products. *Sprouted Grains: Nutritional value, Production and Applications*. AACC International Press pg 143-173. ISBN 9780128115251
- [9] Association of Official Analytical Chemist, Official Methods of Analysis of the Analytical Chemist International, 18th ed. Gathersburg, MD USA (2012).
- [10] Carvalho, L.M.J., Oliveira, A.R.G., Ronoel L. O., Godoy, R.L.O., Pacheco, S., Nutti, M.R., De Carvalho, J.L.V., Pereira, E.J. and Fukuda, W.G (2012). Retention of total carotenoid and β -carotene in yellow sweet cassava (*Manihot esculenta Crantz*) after domestic cooking, *Food and Nutrition Research*; 56: 15788-15797.
- [11] Ilori, A. O., Ogungbemi, K., Balogun, B., Oke, O. A., Balogun, D. A., Odeniyi, T. A., Ajisafe, S. S. and Ogunremi, O. B. (2022). Evaluation of the nutritional composition and acceptability of powdered ogi enriched with date palm fruits. *Research Journal of Food Science and Nutrition*, Volume 7(1); pg.25-29. <https://doi.org/10.31248/RJFSN202134>
- [12] Ijarotimi O. S., Oluwajuyitan T. D., Olugbuyi A. O., Makanjuola S. B. (2022). Comparative study on nutrient composition, functional property and glycaemic index of "Ogi" in healthy rats prepared from selected cereal grains. *Journal of Future Foods* 2(4); pg. 380-387
- [13] Farinde, E. O. (2015). Chemical and sensory properties of sieved and unsieved fortified 'ogi'. *Nature and Science*, 13(1), 49-53.
- [14] Makanjuola, O. M., Adepegba, A. O., Ajayi, A., & Ajoja, S. S. (2017). Proximate composition and sensory qualities of powdered ogi (fermented maize meal) fortified with powdered unfermented locust bean seeds (*Pakiabiglobosa*). *Journal of Global Biosciences*, 6(10), 5289-5295.
- [15] Noah, A.A. Alagamba, E.A (2020). Microbial, proximate, and sensory quality of pito beverage locally prepared and hawked in Ogun State, Nigeria, *Europ. J. Biotechnol. Biosci.* 8(6) (2020) 35-39. <http://eprints.federalpolyilaro.edu.ng/id/eprint/1609>.
- [16] Ojo, D.O. Enujiugha, V.N (2016). Chemical composition, physico-chemical properties and acceptability of instant 'Ogi' from blends of fermented maize, Conophor. Nut. and Melon. Seeds *J. Food Process. Technol.* 7(630) (2016) 2. <https://doi.org/10.4172/2157-7110.1000630>.
- [17] Okoye, J.I. Ene, G.I (2018). Effects of processing on the nutrient and anti-nutrient contents of tiger nut (*Cyperus esculentus Lativum*), *J. Food Tech. Food Chemi.* 1-101.
- [18] Handa, V. Kumar, V. Panghal, A (2017). Effect of soaking and germination on physicochemical and functional attributes of horsegram flour, *J. Food Sci. Technol.* 54(13); 4229-4239. <https://doi.org/10.1007/s13197-017-2892-1>.

UNDER PEER REVIEW