

A COMPARATIVE ANALYSIS OF THE INFLUENCE OF VARIOUS PROCESSING TECHNIQUES ON QUALITY ATTRIBUTES OF JACKFRUIT SEED FLOUR (JSF) FROM DIFFERENT CULTIVARS

ABSTRACT

The study was aimed at investigating the effect of processing on the functional properties, proximate composition and storage stability of jackfruit seed flour from different cultivars. The jackfruit seeds of cultivars *Koozha* and *Varikka* were selected for the study. The water absorption capacity (WAC), oil absorption capacity (OAC), solubility index (SI), swelling power (SP) and bulk density (BD) of the JSF ranged between 194.6-200.1ml/100g, 84.19-85.06ml/100g, 1.12-1.20%, 3.12-3.51g/g and 0.67-0.71g/cm³, respectively. The moisture, ash, crude fat, protein, crude fiber and carbohydrate contents varied from 4.86-5.11%, 2.40-2.60%, 1.68-1.76%, 12.22-12.79g/100g, 2.82-2.98% and 72.01- 72.45g/100g, respectively. Microbial analysis of jackfruit seed flour was performed to determine its shelf life and showed favorable results. Jackfruit seed flour produced from different processing techniques is found to have good storage stability. The jackfruit seed flour thus processed can be employed as a key ingredient to create a wide range of nutrient-dense food products.

Keywords

Jackfruit seed, *Koozha*, *varikka*, flour, tray drying, roasting, functional properties

INTRODUCTION

Research in recent years has emphasized the need to identify lesser-known and underutilised crops, many of which are useful in human nutrition. The jackfruit, *Artocarpus heterophyllus* is native to India and is a member of the mulberry family. It is widely distributed in the Western Ghats, which has a variety of wildlife [14]. Despite comprising 10% to 15% of the fruit's weight and providing significant nutritional value, jackfruit seeds are frequently disregarded and undervalued [11]. Due to a lack of knowledge about effective ways of usage, a lack of post-harvest technology, and weaknesses in the supply chain networks, jackfruit is said to be an underutilised fruit. The seeds of the jackfruit are an excellent source of protein, fiber, and carbohydrates. Additionally, jackfruit is rich in various minerals such as N, P, K, Ca, Mg, S, Zn, and Cu [18]. Due to the difficulties in processing and storing seeds, enormous amounts of seeds are wasted every year. The storage life of the seeds can be refined by storing at cool, moist conditions and welding processing techniques such as roasting. The dried seeds can be ground into flour and incorporated to various food products to increase their value and enhance the shelf life. The objective of this study therefore is to determine the effects of various processing methods on the functional properties, proximate composition and storage stability of jackfruit seed flour.

2 MATERIALS AND METHOD

2.1 Raw materials

Jackfruit cultivars *Koozha* (k) and *Varikka* (v) were selected for the study. In order to gather seeds, mature fruits were chosen. Ripened jackfruits were purchased from Instructional Farm, College of Agriculture, Vellayani and also from the local markets.

2.1.1 Processing of Jackfruit seed flour

The seeds of both cultivars were manually cleansed, the white arils (seed coat) were peeled off, and the brown spermoderm was removed using a stainless-steel knife. The seeds were rinsed and washed with

Commented [A1]: When starting an abstract for a research paper, it's often helpful to begin with a brief introduction or background statement to set the context of your study. Just a sentence or two can give the reader a sense of the research problem or broader topic, helping them understand the context before you dive into the specific objectives.

A final sentence that briefly mentions the implications or future research directions could add depth and completeness of the abstract.

Commented [A2]: This statement lacks citations or data to substantiate the claim about the extent of waste. Including a reference to research or data that quantifies the waste or outlines the specific challenges in processing and storing jackfruit seeds would strengthen the statement and enhance the credibility of your research.

Commented [A3]: Clarify the criteria used for selecting these specific cultivars.

Commented [A4]: Include how you determined that the fruits were mature and ripened. This can add credibility to your methodology.

Commented [A5]: It would be helpful to provide more details about where the jackfruits were purchased, such as the specific locations of the local markets or the Instructional Farm. This gives a clearer picture of where the materials came from and adds context to your research.


running water. Then it was gelatinized at 65°C for 10 minutes. After gelatinization the seeds were sliced and dehydrated using the following techniques.

D₁ – Tray drying

D₂- Roasting

Seeds from both cultivars were dried using the above-mentioned techniques. For Tray drying the drying time and temperature were standardized as 12 hours at 110 °C and for Roasting 1 hour at – 100 °C. After dehydration seeds from both cultivars were ground to powder and sieved through a fine wire mesh (150 micron), packed in polyethylene pouches and used for further analysis.

Fig. 1 Preparation of Jackfruit seed flour (Flow chart)



Commented [A6]: To ensure grammatical consistency, the steps should all start with either verbs or nouns. This approach makes the sequence clear and uniform.

2.2 Analytical methods

2.2 Determination of Functional properties of Jackfruit seed flour

Functional properties are very critical to the quality of food. The elements of food, particularly the carbohydrates, proteins, fats and oils, moisture, fiber, and ash, as well as the structures of these elements, have an impact on the functional qualities of foods and flours [5]. Water absorption capacity and oil absorption capacity of the jackfruit seed flour was determined by the method of [26] and [6]. Solubility index and swelling power were determined according to the method of [2] and [17]. Bulk density was determined by the method suggested by [23].

2.3 Determination of Proximate composition of Jackfruit Seed Flour

Moisture, ash and crude protein and crude fat of the samples were determined by the AOAC [3] method. Crude fiber content was estimated by the method of AOAC [4]. The carbohydrate content was calculated by difference, (100- sum of the values for moisture, crude protein, crude fiber, crude fat and crude ash).

2.4 Determination of Storage Stability of Jackfruit Seed Flour

Microbial population analysis in developed food products is significant since it determines food quality and safety. Food safety is guaranteed by eliminating pathogenic microorganisms and inhibiting their proliferation using all conceivable measures. The moisture intake, Total viable count (TVC), Total fungal count (TFC), and Total coliform count (TCC) of the jackfruit seed flour were measured at 0 days (initial) and after three months (90 days) of storage to assess its storage properties. The bacterial, fungal, and coliform growth was examined using Nutrient agar, Rose Bengal agar, and Eosin methylene blue (EMB), respectively. The evaluation was done by serial dilution of samples followed by pour plating technique suggested by [15]. The technique was used to detect the presence of microbes.

2.5 Statistical analysis

All data obtained from various analyses were pooled and subjected to Completely Randomized Design (One-way ANOVA) using KAU-GRAPES software.

3 RESULTS AND DISCUSSION

3.1 Functional properties of Jackfruit seed flour

The functional properties of the jackfruit seed flour samples are presented in Table 1.

Commented [A7]: Instead of just mentioning that carbohydrates were calculated by difference, you could briefly explain why this approach is commonly used in proximate analysis.

The water absorption capacity of a food material reflects the ability of the material to associate with water under conditions of limited water, and hence its ability to bind the water [27]. In the present investigation water absorption capacity of the jackfruit seed flour ranged between 194.6-200.1 ml/100g. Water absorption capacity was significantly higher for the cultivar *Varikka* (VD₁-200.1 ml/100g and VD₂- 198.6 ml/100g). The findings abide by the study done by Reddy et al [25] and Borghi and Bharathi [8], who reported the water absorption capacity of jackfruit seed flour as 184.27 ml/100g and 201.37 ml/100g, respectively.

Commented [A8]: Add a brief explanation for this result.

Oil absorption is an important property in food formulations because fat improve the flavour and mouth feel of food. Oil absorption capability is mostly due to the physical trapping of oils. It represents the rate at which protein binds to fat in food compositions [27]. The oil absorption capacity obtained for the seed flour samples in this study ranged from 84.19- 85.06 ml/100g. Oil absorption capacity was found to be higher for the seed flour samples processed using the roasting method (VD₂- 85.06 ml/100 g and KD₂- 84.88 ml/100 g). The results are comparable with the values obtained by Islam et al [12] and Borghi and Bharathi [8], who reported oil absorption capacity of jackfruit seed flour as 86.00 ml/100g and 89.93 ml/100g, respectively. Nabubuya et al. [20] reported oil absorption capacity of jackfruit seed flour as 129.5ml/100g, which is higher than the obtained values.

Commented [A9]: To provide a clearer picture, it would be helpful to explore why Nabubuya et al. reported a higher value. For instance, differences in how the seeds were roasted or prepared might explain the variation in results.

Solubility in foods is a chemical and functional quality that refers to a food substance's capacity to dissolve in a solvent, most commonly water or oil. A substance's solubility is essentially determined by the chemical and physical properties of the solvent and solute, as well as pressure, pH, temperature, and the presence of additional substances in the solution [9]. The solubility index of the jackfruit seed flour samples ranged from 1.12 – 1.20%. The highest solubility index was observed for the cultivar *Varikka* (VD₁ -1.20% and VD₂ 1.17%). The solubility index of the cultivar *Koozha* seed flour was found to be on par across different processing techniques. The obtained values are comparable with the values reported by Akter and Haque [1] 1.80% and Islam et al [12] 2.31%.

The swelling power is the measure of the starch ability to absorb water and swell, and also reflects the extent of associative forces in the starch granules. It is an indication of the non-covalent bonding between the molecules of starch granules and also one of the factors of the α amylose and amylopectin ratios [13]. The swelling power obtained for the seed flour samples in this investigation ranged from 3.12 g/g – 3.51g/g. It was observed that the cultivar *Varikka* exhibited highest swelling power (VD₁ – 3.51 g/g and VD₂ – 3.41 g/g) than the cultivar *Koozha*. The obtained values are similar with the values reported by Akter and Haque [1] 3.62g/g and Ocloo et al [21] 4.77g/g. The swelling power of jackfruit seed flour were reported as 39.0g/g by [20] which is highest than the obtained values.

Commented [A10]: Mention potential reasons for this result.

Bulk density represents the relative volume or capacity of the needed packaging material. The higher the bulk density of the flour, the denser the amount of packaging material required. It reflects the porosity of a food product, which effects the design of the package and can be used to determine the type of packing material necessary [13]. The bulk density of jackfruit seed flour samples ranged between 0.67- 0.71 g/cm³, it was discovered that the cultivar *Varikka* had a higher bulk density (VD₁- 0.71 g/cm³ and VD₂- 0.70 g/cm³) than the cultivar *Koozha*. The findings abide by the study done by Odoemelum [22] 0.61 g/cm³ and Reddy et al [25] 0.73 g/cm³.

Commented [A11]: Mention potential reasons for this result.

Table. 1 Functional properties of Jackfruit seed flour

Treatments	WAC (ml/100g)	OAC (ml/100g)	SI (%)	SP (g/g)	BD (g/cm ³)
VD ₁	200.1 ^a	84.51 ^b	1.20 ^a	3.51 ^a	0.71 ^a
KD ₁	196.8 ^c	84.19 ^c	1.12 ^c	3.12 ^d	0.68 ^c

VD₂	198.6 ^b	85.06 ^a	1.17 ^b	3.41 ^b	0.70 ^b
KD₂	194.6 ^d	84.88 ^a	1.12 ^c	3.13 ^c	0.67 ^d
±SE(m)	0.083	0.078	0.004	0.004	0.003
CV %	0.073	0.16	0.612	0.215	0.72

Values are means of triplicates. Values with different superscripts (a,b,c,d) within the same column are significantly different ($P \leq 0.05$). VD₁- Varikka Tray dried jackfruit seed flour, KD₁- Koozha Tray dried jackfruit Seed flour, VD₂- Varikka Roasted jackfruit seed flour, KD₂-Koozha Roasted jackfruit Seed flour. WAC= Water absorption capacity, OAC= Oil absorption capacity, SI= Solubility index, SP= Swelling power, BD = Bulk density.

3.2 Proximate composition of Jackfruit seed flour

The proximate composition of jackfruit seed flour samples is depicted in Table.2.

In the present investigation ash content of jackfruit seed flour ranged from 2.40 – 2.60%. It was discovered that the ash content was found to be significantly higher for tray-dried seed flour samples (VD₁-2.60% and KD₁-2.57%). According to Eke-Ejiofor et al. [10], the ash content of dried jackfruit seed flour is 2.46%, whereas that of roasted jackfruit seed flour is 2.45%. Another study conducted by Nabubuya et al [20] reported the ash content of jackfruit seed flour as 2.60%. The obtained values correspond to the reported values.

In the current study the fat content of jackfruit seed flour ranged from 1.68 – 1.76%. The fat content was found to be substantially higher in the cultivar *Varikka* (VD₁- 1.71% and VD₂- 1.76%). Jackfruit seed flour contains a negligible amount of fat, which prevents rancidity issues and prolongs the flour's shelf life. The results are consistent with the findings of [12] and [24], who reported the fat content of jackfruit seed flour as 1.77% and 1.44%, respectively.

Protein is one of the most important nutrients required by the body to carry out a wide range of functions essential for the maintenance of life. Protein content of jackfruit seed flour ranged from 12.22g-12.79g/100g. The protein content was found to be significantly higher in the cultivar *Varikka* (VD₁- 12.67 g/100g and VD₂- 12.79 g/100g). The results are comparable with the values obtained by [21] 13.50g/100g and [24] 13.9g/100g.

The crude fiber content of the jackfruit seed flour samples ranged between 2.82% - 2.98%. The crude fiber content of seed flour samples was found to be substantially higher in cultivar *Varikka* (VD₁-2.98% and VD₂-2.91%). The findings abide by the study done by Ocloo et al. [21], who reported the crude fiber content of jackfruit seed flour as 3.19. The crude fiber content of autoclaved jackfruit seed flour was found to be higher, at 6.17%, according to [10].

Carbohydrates are one of the most abundant and widespread organic substance in nature. The carbohydrate content of the jackfruit seed flour samples ranged between 72.01 – 72.45 g/100g. It was found that based on the processing techniques, tray dried seed flour samples had more carbohydrates than roasted seed flour samples (VD₁-72.45 g/100 g and KD₁-72.11 g/100 g). The findings are comparable with the values reported by Islam et al. [12] and Juárez-Barrientos et al [16], who reported 71.46 g/100 g and 73.44 g/100 g of carbohydrate content in jackfruit seed flour, respectively.

Commented [A12]: Use consistent units and formats when presenting data. Here's how you can revise it: "The protein content of jackfruit seed flour ranged from 12.22 g/100g to 12.79 g/100g."

Commented [A13]: "The carbohydrate content of the jackfruit seed flour samples ranged from 72.01 g/100g to 72.45 g/100g."

Table. 2 Proximate composition of Jackfruit seed flour

Treatments	Ash (%)	Fat (%)	Protein (g/100g)	Crude fiber (%)	Carbohydrate (g/100g)
VD ₁	2.60 ^a	1.71 ^b	12.67 ^b	2.98 ^a	72.45 ^a
KD ₁	2.57 ^b	1.69 ^c	12.22 ^d	2.82 ^d	72.11 ^b
VD ₂	2.45 ^c	1.76 ^a	12.79 ^a	2.91 ^b	72.01 ^d
KD ₂	2.40 ^d	1.68 ^d	12.27 ^c	2.84 ^c	72.04 ^c
±SE(m)	0.003	0.003	0.007	0.006	0.004
CV%	0.23	0.337	0.095	0.346	0.01

Values are means of triplicates. Values with different superscripts (a,b,c,d) within the same column are significantly different ($P \leq 0.05$). VD₁- Varikka Tray dried jackfruit seed flour, KD₁- Koozha Tray dried jackfruit Seed flour, VD₂- Varikka Roasted jackfruit seed flour, KD₂-Koozha Roasted jackfruit Seed flour.

3.3 Storage stability of Jackfruit seed flour

The microbial analysis of jackfruit seed flour samples is presented in the Table 3.

The microbial analysis report of jackfruit seed flour is depicted in the Table.3. Moisture gives an indication of how much water and total solid matter are present in the seed flour samples. It is also a measure of the flour's storability. Reduced moisture content implies better shelf life and stability. Initially, the moisture content of jackfruit seed flour ranged between 4.86 – 5.11%. With respect to processing methods, tray drying had significantly higher moisture content. After a period of three months, it was noted that the jackfruit seed flour moisture content had risen constantly while remaining within the acceptable range of 5.21 to 5.71 percent. The findings are consistent with the study of Morshed et al. [19], who reported that the moisture content of jackfruit seed flour during the storage period was between 5.43 and 5.70%.

The study examined the distribution and number of microorganisms in jackfruit seed flour over a three-month (90-days) storage period. The results demonstrated that the storage time had a substantial impact on the total viable count (TVC) in the seed flour, with the count increasing from day 30 to day 90. However, the counts were within the permissible limits. Initially, there was no presence of fungal colonies for up to two months (60 days) in the processed seed flour. After two months (60 days), the existence of fungal colonies was detected in the seed flour; nonetheless, the total fungal count remained below acceptable limits. There were no coliform colonies found in the jackfruit seed flour during the storage period. A study conducted by Borgis et al [7] reported that pan roasted and unprocessed jackfruit seed flour contain significantly lower bacterial and fungal colonies. Morshed et al. [19] reported that for the first 96 days of storage, jackfruit seed flour was free of microbes.

Commented [A14]: There's a bit of redundancy in the sentences. You can streamline this by combining them into one clear statement.

Commented [A15]: Expand on how increased moisture content specifically impacts storage stability.

Commented [A16]: Explain why tray drying resulted in higher moisture content compared to other methods. This adds depth to the discussion a

Table. 3 Storage Stability of Jackfruit seed flour

Treatments	Storage period	Moisture (%)	TVC (CFU/gram of sample)	TFC (CFU/gram of sample)	TCC (CFU/gram of sample)
VD ₁	Initial count	5.04 ^k	2.0×10^2	ND	ND
	30 days	5.19 ^h	4.0×10^2	ND	ND
	60 days	5.46 ^c	8.0×10^2	ND	ND
	90 days	5.71 ^a	12.1×10^2	1.8×10^2	ND
KD ₁	Initial count	5.11 ⁱ	2.1×10^2	ND	ND
	30 days	5.25 ^f	4.5×10^2	ND	ND

	60 days	5.44 ^d	8.3 × 10 ²	ND	ND
	90 days	5.67 ^b	12.0 × 10 ²	2.0 × 10 ²	ND
VD ₂	Initial count	4.86 ^m	1.5 × 10 ²	ND	ND
	30 days	4.97 ⁱ	3.5 × 10 ²	ND	ND
	60 days	5.09 ^j	7.6 × 10 ²	ND	ND
	90 days	5.21 ^g	12.5 × 10 ²	2.2 × 10 ²	ND
KD ₂	Initial count	4.97 ⁱ	1.6 × 10 ²	ND	ND
	30 days	5.12 ^j	3.6 × 10 ²	ND	ND
	60 days	5.19 ^h	7.6 × 10 ²	ND	ND
	90 days	5.31 ^e	12.6 × 10 ²	2.1 × 10 ²	ND
±SE(m)		0.004	0.024	0.048	–
CV (%)		0.143	0.621	15.984	–

Values are means of triplicates. Values with different superscripts (a,b,c,d) within the same column are significantly different ($P \leq 0.05$). TVC- Total viable count, TFC- Total fungal count, TCC- Total coliform count, ND – Not Detected. VD₁- Varikka Tray dried jackfruit seed flour, KD₁- Koozha Tray dried jackfruit Seed flour, VD₂- Varikka Roasted jackfruit seed flour, KD₂-Koozha Roasted jackfruit Seed flour.

Conclusion

The purpose of the study was to assess the impact of processing methods on functional qualities, proximate composition and storage stability of jackfruit seed flour from different cultivars. This study used tray drying and roasting as the primary processing techniques for the production of the jackfruit seed flour. There was a significant difference in the functional properties and proximate composition of jackfruit seed flour from different cultivars. Quality analysis revealed that jackfruit seed flour is excellent source of carbohydrate and protein and they are also good source of fiber. Jackfruit seed flour had negligible amount of fat. Less fat allows for the prevention of rancidity issues and longer shelf life of the flour. From microbial analysis it is evident that the jackfruit seed flour can be stored up to three months (90 days) without any microbial spoilage. This method allows you to preserve jackfruit seeds by converting them into flour and extending their shelf life. The findings revealed that jackfruit seed flour can be employed as a key ingredient to create a wide range of nutrient-dense food products.

Reference

1. Akter B, Haque MA. Utilization of jackfruit (*Artocarpus heterophyllus*) seed's flour in food processing: A review. The Agriculturists. 2018;16(2):131-142.
2. Anderson RA. Gelatinization of corn grits by roll-and extrusion-cooking. J. Cereal. Sci. 1969; 14(2):4-7.
3. AOAC Official methods of analysis: 12th Edition. Association of Official Analytical Chemists. Washington, DC. 1990;114.
4. AOAC. Official Methods of Analysis: 18th Edition. Association of Official Analytical Chemists. Washington, DC. 2005;210.
5. Awuchi CG, Igwe VS, Echeta CK. The functional properties of foods and flours. International Journal of Advanced Academic Research. 2019;5(11):139-160.

Commented [A17]: Specify the nature of the "significant differences" observed.

Commented [A18]: While older references are valuable, citing recent studies—those published within the last 10 years—is equally important to ensure your work reflects the latest advancements and current knowledge in the field. For example, the research study titled "Production, Proximate Analysis, and Functional Properties of Dragon Fruit Peel Powder" closely parallels your study and offers relevant discussions and insights that may be beneficial for your work.

6. Beuchat LR. Functional and electrophoretic characteristics of succinylated peanut flour protein. *Journal of Agricultural and Food chemistry*. 1977;25(2):258-261.
7. Borgis S, Bharati P, Shirnalli G. Effect of processing on storage and microbial quality of jackfruit (*Artocarpus heterophyllus* Lam.) seed flour. *Int. J. Curr. Microbiol. Appl. Sci*. 2018; 7:3058-3066.
8. Borgis S, Bharati P. Processing characteristics and acceptability of jackfruit (*Artocarpus heterophyllus* Lam.) seeds, physical and functional properties of its flour. *EPRA International Journal of Research and Development*. 2020;5(10):193-202.
9. David O, Arthur E, Kwadwo SO, Badu E, Sakyi P. Proximate composition and some functional properties of soft wheat flour. *International Journal of Innovative Research in Science, Engineering and Technology*. 2015;4(2):753-758.
10. Eke-Ejiofor J, Beleya EA, Onyenorah NI. The effect of processing methods on the functional and compositional properties of jackfruit seed flour. *Int. J. Nutr. Food Sci*. 2014;3(3):166-173.
11. Hossain MT, Hossain MM, Sarker M, Shuvo AN, Alam MM, Rahman MS. Development and quality evaluation of bread supplemented with jackfruit seed flour. *International Journal of Nutrition and Food Sciences*. 2014;3(5):484.
12. Islam MS, Begum R, Khatun M, Dey KC. A study on nutritional and functional properties analysis of jackfruit seed flour and value addition to biscuits. *Int J Eng Res Technol*. 2015;4(12):139-147.
13. Iwe MO, Onyeukwu U, Agiriga AN. Proximate, functional and pasting properties of FARO 44 rice, African yam bean and brown cowpea seeds composite flour. *Cogent Food & Agriculture*. 2016; 31(1):1142409.
14. Jagadeesh SL, Reddy BS, Basavaraj N, Swamy GS, Gorbak K, Hegde L, Raghavan GS, Kajjidoni ST. Inter tree variability for fruit quality in jackfruit selections of Western Ghats of India. *Scientia Horticulturae*. 2007;112(4):382-387.
15. Johnson ML, Curl T. Standard methods of biochemical analysis. Kalyani publishers, New Delhi. 1973;62-65.
16. Juárez-Barrientos JM, Hernández-Santos B, Herman-Lara E, Martínez-Sánchez CE, Torruco-Uco JG, Ramírez-Rivera EJ, Pineda-Pineda JM, Rodríguez-Miranda J. Effects of boiling on the functional, thermal and compositional properties of the Mexican jackfruit (*Artocarpus heterophyllus*) seed jackfruit seed meal properties. *Emirates Journal of Food and Agriculture*. 2017;29(1):1.
17. Leach HW. Structure of starch granule I. Swelling and solubility patterns of various starches. *J. Cereal Chem*. 1959; 36:534-544.
18. Maurya P, Mogra R. Assessment of consumption practices of jackfruit (*Artocarpus heterophyllus* Lam) in the villages of Jalalpur block, district Ambedkar Nagar (Uttar Pradesh) India. *Advances in Life Sciences*. 2016;5(5):1638-1644.
19. Morshed MH, Ibrahim M, Helali MO, Alam AK, Amin R. Storage life and quality characteristics of nutritious flour from ripe jackfruit seed. *Journal of Engineering*. 2019;10(2):19-24.

20. Nabubuya A, Mugabi R, Kaggwa A, Ainebyona P, Nalugya R. Influence of roasting on the proximate, functional and sensory properties of Jackfruit seeds and amaranth grain composite complementary flours. *Tanzania Journal of Science*. 2022;48(1):156-169.
21. Ocloo FC, Bansa D, Boatun R, Adom T, Agbemavor WS. Physico-chemical, functional and pasting characteristics of flour produced from Jackfruits (*Artocarpus heterophyllus*) seeds. *Agriculture and biology journal of North America*. 2010;1(5):903-908.
22. Odoemelam SA. Functional properties of raw and heat processed jackfruit (*Artocarpus heterophyllus*) flour. *Pakistan Journal of Nutrition*. 2005;4(6):366-370.
23. Okaka JC, Potter NN. Physico-chemical and functional properties of cowpea powders processed to reduce beany flavor. *Journal of Food science*. 1979;44(4):1235-1240.
24. Palamthodi S, Shimpi S, Tungare K. A study on nutritional composition and functional properties of wheat, ragi and jackfruit seed composite flour. *Food Science and Applied Biotechnology*. 2021; 4(1):63-75.
25. Reddy SS, Devi GN, Lakshmi K, Lakshmi KB. Physico-chemical and functional properties of jackfruit (*Artocarpus heterophyllus*) seed flour. *The Journal of Research ANGRAU*. 2022;50(4):45-56.
26. Sathe SK, Deshpande SS, Salunkhe DK. Functional properties of winged bean [*Psophocarpus tetragonolobus* (L.) DC] proteins. *Journal of Food Science*. 1982;47(2):503-509.
27. Singh U. Functional properties of grain legume flours. *Journal of food science and technology (Mysore)*. 2001;38(3):191-199.