

---

# Effect of Incorporation of Carrot Flour on Quality Characteristics of Cake

**ABSTRACT** Cake is a form of food that is baked and usually sweet in taste. It provides fats, carbohydrates and body building protein. The aim of the present study was to develop carrot blended cake and to assess its quality characteristics during ambient storage conditions. For the development of carrot blended cake, refined wheat flour, whole wheat flour and carrot flour were blended in different proportions of T<sub>1</sub>(100:0::RF:CF), T<sub>2</sub>(90:10::RF:CF), T<sub>3</sub>(80:20::RF:CF), T<sub>4</sub>(70:30::RF:CF), T<sub>5</sub>(100:00::WWF:CF), T<sub>6</sub>(90:10::WWF:CF), T<sub>7</sub>(80:20::WWF:CF) and T<sub>8</sub>(70:30::WWF:CF) and were packed in high density polythene (HDPE) polythene pouches and stored at refrigeration conditions and subjected to quality characteristics at an interval of 7 days for a period of 21 days. The results showed that the phosphorus and iron content of cake decreased with an increase in storage period. No microbial count was detected in the fresh samples, but it increased with an increase in the storage period but was within the safe limits. On the basis of sensory evaluation (colour, texture, taste and flavour) of carrot blended cake, treatment T<sub>3</sub> (**80:20::RWF:CF**) was adjudged to be the best among all the treatments with mean score of colour (8.26), texture (8.15), taste (8.12) and flavour (8.05), respectively.

## 1. INTRODUCTION

Carrot (*Daucus carota* L.) is an important vegetable crop grown all over the world for its enlarged tap roots and belongs to family Apiaceae. It is grown during spring, summer and autumn in temperate climate and during winter in tropical and subtropical climate [1]. Carrots are healthy vegetables, but this agricultural product has low value when they are sold as a raw commodity. To become value-added agricultural products, the value of raw carrots has to be increased through the addition of ingredients or processes that make them more attractive to the buyer. Increasing the added value of the carrots requires the development of food products that considering the voice of the customer [2]. Its maximum retention is of utmost importance for the preservation of the attractive appearance and dietary value of the product. Carrots have a moisture content of 80–90% (wet basis) at the time of harvest. They are seasonal in nature and highly susceptible to moisture loss leading to wilting and loss of fresh appeal [3]. Wheat flour is one of the important cereal grain because of its use for the preparation of many baked products. Bakery products are mostly prepared from refined wheat flour with 60-75% extraction rate, which has still lower nutritive value. However, such products can easily be enriched and fortified at lower cost with proteins, fibre, various vitamins and minerals to meet the specific need of the target groups and vulnerable sections of the population who are undernourished and malnourished [4]. Cakes are the most popular bakery products worldwide especially among children, they love to eat cakes, cupcakes and biscuits. Cake is a complex of fat and water emulsion system containing flour, fat, sugar, baking powder and baking soda. A proper combination of the ingredients can give a high-quality product with desirable flavour and texture [5]. The incorporation of carrot flour in cake may improve its nutritive value, lower the cost and help in enhancing the utilization of carrot.

## 2. MATERIALS AND METHODS

Refined wheat flour and whole wheat flour were substituted with different percentage of carrot flour i.e. 100:0, 90:10, 80:20 and 70:30 for the preparation of blended cake. The batter was thoroughly mixed by adding required amount of all ingredients. After mixing, the batter was filled in moulds containing pan. The cake pan was placed in baking tray smeared with fat and baked at 180°C for 30 min in baking oven/deck oven. After baking, the cakes will be cooled and packed airtight in high density polythene (HDPE) and kept in refrigerator for storage. The stored product was analyzed for quality and organoleptic characteristics at a regular interval of 7 days. The mineral content was estimated as per the procedure described by AOAC [6]. As per the procedure coined by Amerine et al. [7], score of 5.5 and above was reflected as acceptable. The experiment was laid out in factorial CRD with nine treatments

and replicated thrice. The data obtained were statistically analysed as per the procedure of Gomez and Gomez [8] using OPSTAT software.

### 3. RESULTS AND DISCUSSION

#### 3.1 Minerals (Phosphorus and Iron)

During the storage of 21 days, phosphorus and iron content of different treatments revealed that the mean phosphorus and iron content of carrot blended cake significantly decreased from the initial level of 186.13 to 185.04 mg/100g and 2.47 mg/100g to 1.95 mg/100g, respectively (Table 1). Maximum mean phosphorus and iron content of 194.08 mg/100g and 2.60 mg/100g were recorded in T<sub>5</sub> (100:0::WF:CF) whereas minimum of 177.41 mg/100g and 1.75 mg/100g were recorded in the treatment T<sub>4</sub> (70:30::RWF:CF), respectively. The results are in conformity with the findings of Pandey [9] and Singh [10]. The decrease in minerals might be due to its heat sensitivity, even at ambient temperature which leads to destruction of minerals in cake during the storage period. Similar findings were reported by Anu et al. (2008) in banana blended cake, Olatunde et al. [11] in pigeon pea, sweet potato and wheat flour cake and Pandey et al. [12] in carrot powder incorporated bread.

#### 3.2 Microbial Load

A glance of data in Table 2 showed the effect of various treatments and storage on microbial load of cake. However, the microbial loads of the cake samples observed to be on the increase from day 7 to 21 days. The values obtained for cake samples were within the acceptable international commission on microbiological specification (ICMSF) limits and recommendation for products of this nature (10<sup>5</sup>cfu/g) in good manufactured practice [13]. Similar trend of microbial count during storage period was reported by Singh et al. 2022 in pearl millet cake.

Table 1. Effect of treatment and storage period on phosphorus and iron of carrot blended cake Treatments	Phosphorus (mg/100g)				Mean (Treatment)	Iron (mg/100g)			Me (Tr
	Storage period (Days)		Storage period (Days)			Storage period (Days)			
	0	7	14	21	0	7	14		
T1 (100:0::R WF:CF)	185.21	184.89	184.46	184.15	<b>184.68</b>	2.82	2.59	2.45	
T2 (90:10::R WF:CF)	182.79	182.46	182.02	181.67	<b>182.23</b>	2.57	2.30	2.18	
T3 (80:20::R WF:CF)	180.47	180.18	179.74	179.40	<b>179.95</b>	2.32	2.08	1.94	
T4 (70:30::R WF:CF)	177.95	177.63	177.21	176.86	<b>177.41</b>	2.06	1.80	1.63	
T5 (100:0::W WF:CF)	194.62	194.29	193.88	193.54	<b>194.08</b>	2.89	2.65	2.49	
T6 (90:10::W WF:CF)	191.70	191.37	190.95	190.62	<b>191.16</b>	2.63	2.41	2.26	
T7 (80:20::W WF:CF)	189.34	189.06	188.61	188.30	<b>188.83</b>	2.38	2.12	1.98	
T8 (70:30::W WF:CF)	186.97	186.64	186.19	185.82	<b>186.40</b>	2.10	1.87	1.69	
Mean (Storage)	<b>186.13</b>	<b>185.81</b>	<b>185.38</b>	<b>185.04</b>	<b>2.47</b>	<b>2.23</b>	<b>2.08</b>	<b>2.08</b>	
Effects C.D. (p=0.05)	C.D. (p=0.05)		C.D. (p=0.05)		C.D. (p=0.05)		C.D. (p=0.05)		
Treatment (T)	0.06		0.01		0.06		0.01		

Storage (S) 0.04 0.01

T □ S 0.13 0.03

**Table 3. Effect of treatment and storage period on colour and texture of carrot blended cake**

Treatments	Colour					Mean (Treatment)	Texture		
	Storage period (Days)						Storage period (Days)		
	0	7	14	21	28		0	7	14
T1 (100:0::R WF:CF)	7.45	7.27	7.08	6.87	7.17	7.34	7.19	7.02	
T2 (90:10::R WF:CF)	7.83	7.64	7.43	7.25	<b>7.54</b>	7.76	7.57	7.40	
T3 (80:20::R WF:CF)	8.56	8.33	8.17	7.99	<b>8.26</b>	8.41	8.23	8.06	
T4 (70:30::R WF:CF)	8.18	8.02	7.83	7.62	<b>7.91</b>	8.10	7.92	7.74	
T5 (100:0::W WF:CF)	7.29	7.14	6.95	6.78	<b>7.04</b>	7.14	6.91	6.77	
T6 (90:10::W WF:CF)	7.67	7.51	7.32	7.11	<b>7.40</b>	7.55	7.38	7.23	
T7 (80:20::W WF:CF)	8.35	8.20	8.05	7.87	<b>8.12</b>	8.27	8.12	7.94	
T8 (70:30::W WF:CF)	7.92	7.76	7.59	7.38	<b>7.67</b>	7.95	7.78	7.59	
<b>Mean (Storage)</b>	<b>7.91</b>	<b>7.73</b>	<b>7.55</b>	<b>7.36</b>	<b>7.36</b>	<b>7.81</b>	<b>7.64</b>	<b>7.47</b>	

Effects CD (p=0.05) CD(p=0.05)

Treatment (T) 0.03 0.02

Storage (S) 0.02 0.01

T □ S 0.06 0.04

### 3.3 Colour

A significant reduction in colour scores was recorded during 21 days of storage (Table 3). The mean values of colour scores showed a reducing trend from initial level of 7.91 to 7.36 with the advancement in storage. The results are in accordance with the findings of Ghodke and Kale [14], Mittal [15] and Khalaf [16]. Among treatments, highest mean colour score of 8.26 was noticed in treatment T<sub>3</sub> (80:20::RWF:CF) while the lowest score of 7.04 was noticed in T<sub>5</sub> (100:0::WF:CF). The decrease in colour score of the product during storage might be due to browning reaction which could have resulted from non enzymatic oxidation of ascorbic acid. Sahni [17] and Shende [18] also reported decline in colour scores during storage while preparing carrot based cookies and value added sponge cake, respectively.

### 3.4 Texture

From the data in Table 3, it was evident that there was decrease in the texture score of carrot blended cake with the progress in the storage period. Among treatments maximum mean texture score of 8.15 was reported in cake prepared with 80% refined wheat flour and 20% carrot flour (T<sub>3</sub>) whereas minimum score of 6.85 was reported in the T<sub>5</sub> (100:0::WF:CF). The results are in harmony with the findings of Anu et al. (2008), Ghodke and Kale [14] and Baltacıoğlu et al. [19] who reported decreasing trend in texture score in banana blended cake, carrot-brahmibased biscuits and cookies incorporated with carrot powder, respectively. Maximum mean texture score was recorded at beginning of storage (7.81) while the

minimum was recorded after 21 days of storage (7.29). The decrease in quality score of texture was mainly due to the absorption of moisture that has negative effect on texture. Similar observations were also supported by Sahni [17] in carrot blended cookies and Khalaf [16] in red carrot powder blended cake.

### **3.5 Taste**

The experimental values of taste scores under different treatments and storage were presented in Table 4. Maximum mean taste score (8.12) was observed in T<sub>3</sub> (80:20::RWF:CF), whereas the minimum (6.83) was scored in T<sub>5</sub> (100:0::WF:CF). The mean taste scores showed a declining trend with the progress in storage period from 7.80 to 7.21. The results are consistent with those of Baltacioğlu et al. [19] and Khalaf [16] who studied the different attributes of carrot blended cookies and carrot powder blended cake, respectively. Taste was found to be affected probably due to lipolytic changes, hastened by increase in the moisture content of cake [20]. The decline in taste scores during storage might also be due to oxidation of fats (free and unsaturated fatty acids) leading to rancidity. The results are in accordance with the observations of Sahni [17], Agrawal et al. [21], Soni [22] and Sule et al. [23].

### **3.6 Flavour**

The data pertaining to flavour scores of carrot blended cake was presented in Table 4. Among treatments highest mean flavour score was recorded in treatment T<sub>3</sub> (8.05), whereas lowest mean flavour score was recorded in (T<sub>5</sub>) 100% wheat flour cake (6.71). Maximum mean flavor score was observed during initial days of storage (7.75) and minimum was observed after 21 days (7.12) of storage. These results are concurred with the findings of Sahni [17] and Ghodke and Kale [14]. This gradual decline in flavour scores might be due to certain enzymatic, physiological or biochemical changes. Mahmood et al. [24] reported that the decrease in flavour score as a function of storage might be due to oxidation of fats and maillard reaction. Similar findings were observed by Salehi et al. [25] while preparing sponge cake with air dried carrot and Pragyia [26] while developing various functional food using carrot [27].

## **4. CONCLUSION**

Cake can be prepared by using 80 per cent refined flour and 20 per cent carrot flour as the optimum proportion to produce the acceptable cake. The use of carrot flour improved the nutritional quality of cake with respect to phosphorus and iron content. On the basis of sensory evaluation (colour, texture, taste and flavour), treatment T<sub>3</sub> (80:20:: refined wheat flour: carrot flour) was found to be the best treatment.

### **DISCLAIMER (ARTIFICIAL INTELLIGENCE)**

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

### **COMPETING INTERESTS**

Authors have declared that no

## **REFERENCES**

1. Thakur N, Jamwal R S. Evaluation of European carrot (*Daucuscarota* L.) genotypes in mid hill conditions of Himachal Pradesh. *Himachal J of Agricultural Res.* 2012;38(1):80-84.
2. Meister MSK, Kava R. Sugar substitutes and your health; 2006.  
Available: <http://acsh.org>. 2006.
3. Rafiq N, Gupta N, Sood M, Bandral J D and Gupta V. Influence of sugar and jaggery syrup on quality and microbial characteristics of osmo-dehydrated carrot cubes during storage. *Plant Archives.* 2023;23(2):305-309.
4. Chopra SMA, Misra SG, Gupta R. Overweight, obesity and related non-communicable diseases in Asian Indian girls and women. *European J Clinica INutri.* 2013;67(7):688-696.
5. Singh P, Gupta N, Bhat A, Sood M, Bandral J D, Sharma S, Dutta U, Rahman R. Studies on mineral and microbial composition of blended pearl millet cake during storage. *Agri. Mech. Asia, Africa and Latin America.* 2022;53(2): 5893-5900.
6. AOAC. Official methods of analysis. 18th Edition, Association of Official Analytical chemists, Gaithersburg; 2007.
7. Amerine MA, Pangborn RM, Roessler EB, Anson ML, Mrak EM, Chichester CO Principles of sensory evaluation of food. *The Sense of Taste.* Academic Press INC. Elsevier Inc., USA. P, Cambridge, Massachusetts. 1965;28-144.
8. Gomez KA, Gomez AA. Statistical procedures for agricultural research. John Wiley and Sons; 1984

9. Pandey P, Grover K, Dhillon TS, Chawla N and Kaur A. Development and quality evaluation of polyphenols enriched black carrot (*Daucuscarota* L.) powder incorporated bread. *Heliyon*, 2024;10(3): 25109.
10. Singh P. Utilization of pearl millet for development of cake. M.Sc. Thesis, Sher-e-Kashmir University of Agricultural Sciences and Technology, Jammu, India; 2021.
11. Olatunde SJ, Ajayi OM., Ogunlakin G Ajala AS. Nutritional and sensory properties of cake made from blends of pigeon pea, sweet potato and wheat flours. *Food Res*, 2019;3(5):456-462.
12. Pandey L. Nutritional evaluation of wheat (*Triticumaestivum*) Varieties and development of value added products using wheat-sorghum-soybean flour blends. M.Sc. Thesis, CCS Haryana Agricultural University, Hisar, India; 2015.
13. ICMSF. Microorganisms in food 2. Sampling for microbiological analysis: Principles and specific applications. Second edition, Blackwell Scientific Publications, UK. 2009;1-131.
14. Ghodke SV, Kale AA. Studies on the development of carrot-brahmi biscuits. *International J of Sci. Eng. Management*, 2017;2(11):47-51.
15. Mittal S. Evaluation of ripe pumpkin flours and its utilization in bakery products. M.Sc. Thesis, DrYS Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, India; 2018
16. Khalaf AA. Effect of replacing wheat flour with red carrot powder on the physicochemical and sensory properties of cake. *International J Research in Eng. Innovation*. 2024;8(1):41-46.
17. Sahni P. Studies on the preparation and evaluation of fibre rich cookies utilizing fruit and vegetable pomace powder. M.Sc. Thesis, Vasant Rao Naik Marathwada Krishi Vidyapeeth, Parbhani, India; 2015.
18. Shende AB. Physico-chemical properties of pearl millet and its utilization in sponge cake. Ph.D. Thesis, Mahatma Phule Krishi Vidyapeeth, Rahuri, India; 2016
19. Baltacıoğlu H, Baltacıoğlu C, Tangüler H. Effect of waste fermented carrot powder addition on quality of biscuits. *Turkish J of Agri - Food Sci. Tech*. 2019;7(9):1237-1244.
20. Butt MS, Arshad MU, Alam MS, Nadeem MT. Bioavailability and storage stability of vitamin A fortificant (retinyl acetate) in fortified cookies. *Food Res. International*. 2007;40(10):1212-1219..
21. Agrawal RS, Shirale DO, Syed HM. Effect of storage on quality evaluation of cookies supplemented with germinated fenugreek seed powder. *International J of Advanced Biol Res*. 2017;7(3):630-633.
22. Soni P. Oat based apple pomace enriched bakery products. M.Sc. Thesis, Dr YSParmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, India; 2019.
  
23. Sule S, Oneh AJ, Agba I M. Effect of carrot powder incorporation on the quality of pasta. *MOJ Food Processing and Tech*. 2019;7(3):99-103.
24. Mahmood S, Butt MS, Anjum FM, Nawaz H. Baking and storage stability of retinyl acetate (vitamin A) fortified cookies. *Pakistan J Nutrition*. 2008;7(4):586-589.
25. Salehi F, Kashaninejad, M, Akbari E, Sobhani S M, Asadi F. Potential of sponge cake making using infrared-hot air dried carrot. *J of Texture Studies*. 2015;47(1): 34-39.
26. Pragya. Utilization of Black Carrot (*Daucuscarota l.*) for Development of Functional Foods. Ph.D. Thesis, Punjab Agricultural University, Ludhiana, Punjab, India; 2018.
27. Sehgal AS, Kawatra A. 2008. Development and nutritional evaluation of pearl millet rich banana cake. *J of Dairying, Foods and Home Sci*. 2008;27(2):138-141.