

Processing and health benefits of underutilized foxtail and barnyard millet for development of functional bakery products

Abstract

The bread sector in India holds the distinction of being the largest food industry in terms of yearly turnover, amounting to almost 3000 crores. Formerly considered a dietary choice limited to individuals with health ailments, bakery products have evolved into essential nutrition for a significant portion of the population. The majority of countries exhibit a notable increase in per capita consumption of bread goods. It is anticipated that there will be a sustained increase in the demand for bakery goods in the future. The anticipated growth rate of 9.8% appears to be quite modest when considering the existing market potential of bakery products among populations with lower and middle incomes. This suggests that there is significant opportunity for the promotion and consumption of bakery products as a means of enhancing dietary enrichment. Bakery products are among the most affordable processed food items available for consumption within the country. Over 70 % of the overall production has historically been contributed by the industry, which is still mainly from the unorganized sector.

Keywords: Bakery products, Barnyard, Foxtail, Functional, Health benefits, Millets, Underutilized

1. Introduction

India is the world's second largest producer of food grains after China and can stand first with the food and agriculture sector (Kalyaniet *al.*, 2014). According to FAOSTAT, India produced roughly 13.21 Megatons of millet overall in 2021 with an estimated harvested area of 97648 square kilometres. Millets are small seeded annual grains with different varieties, such as Pearl millet (*Pennisetumglaucum*), Finger millet (*Eleusinecoracana*), Kodo millet (*Paspalumsetaceum*), Proso millet (*Penicummillaceum*), Foxtail millet (*Setariaitalica*), Little millet (*Panicumsumatrense*), Sorghum (*Sorghum bicolor*), Oats (*Avena sativa*) and Barley (*Hordeumvulgare*) (Sunil *et al.*, 2016).

Millets are vintage foods known to humans and are unique due to their short growing season. Millets are originated in Asia and Africa, and were domesticated by the local populations. Important millet crop grown in India includes sorghum (great millet), bajra (pearl millet), ragi (finger millet) and small millets such as kora (foxtail millet), little millet, kodo millet, proso millet and barnyard millet. These were often referred to as coarse cereals but realizing the nutrient richness of the grains, they are now gazetted as "Nutricereals" by the Government of India (MILLETS- the Future Super Food for India, June 2022).

Moreover, Millets are rich source of Phytochemicals, micronutrients and antioxidants such as, phenolic acids and glycosylated flavonoids (Singh *et al.*, 2012). They are more nutritious and are non-glutinous, non-acid forming and easy to digest. They are good source of energy, protein, fatty acids, vitamins, minerals, dietary fibre and polyphenols (Reddy *et al.*, 2019). In addition to nutritive value, several potential health benefits such as prevention of cancer and cardiovascular diseases, reducing tumour incidence, lowering blood pressure, risk of heart disease, cholesterol and rate of fat absorption, delaying gastric emptying and supplying gastrointestinal bulk have been reported for millets (Truswell, 2002; Gupta *et al.*, 2012).

The demand for healthier bakery items has led to adjustments in ingredients, focusing on enhancing fibre, controlling calories, adding omega and essential fatty acids, and reducing saturated and trans fats. Natural sweeteners are favoured over artificial ones, and low-calorie options are explored, resulting in gluten-free baked goods. Millets, ancient grains cultivated in diverse climates, have been a staple for millennia in Asia and Africa, with China being an early grower around 2700 B.C., predating rice as a staple cereal.

Foxtail millet is remarkably similar grain to paddy rice. Foxtail millet was anciently domesticated in China, immensely aiding the rise of Chinese civilization and continuing to be a staple grain in dry and semi-arid areas (MILLETS- the Future Super Food for India, June 2022). Like most millets, foxtail millet is super source of crude fibre, aids in digestion, and encourages bowel movement, generating a laxative effect which promotes a healthy digestive tract. Additionally, foxtail millet has several health advantages, including the prevention of cancer and the reduction of blood sugar and cholesterol (Sharma & Niranjana, 2018).

Barnyard millet (*Echinochloa spp.*) is emerging as one of the most significant minor millet crops in Asia (Renganathan *et al.*, 2020). Amid various distinct, wild and cultivated species of *Echinochloa*, the two mainly grown and popular species are, *Echinochloa esculenta* (Japanese based barnyard millet) and *Echinochloa frumentacea* (Indian based barnyard millet). Moreover, *Echinochloa* genus constitutes of 20-35 annual and perennial species spread all over and have the potential to grow in any given climatic and agricultural condition. It has always been the staple cereal for the areas having unsuited climatic and soil conditions for cultivation of rice (Sood *et al.*, 2015).

Traditional barnyard millet, *Echinochloa species*, grows in warm and temperate locations including India, China, Japan, and Korea. Indian based Barnyard millet is grown up to 2300 from the mean sea level (MSL) predominantly in kharif season in the elevation of Uttarakhand and Tamil Nadu States (Gomasheet *et al.*, 2017).

Muffins, a popular bakery item, are not quick bread and do not contain yeast. Unlike cupcakes, muffins may have a glazed surface. They are typically consumed in one sitting and fit comfortably in an adult hand, often enjoyed as breakfast or snacks. Common ingredients include wheat flour, sugar, milk, butter, salt, baking soda, and baking powder. Whole grains, traditionally grown in India, have varied productivity, and new varieties have shown potential medicinal and functional benefits.

Recently the word 'nutraceuticals' has gained more importance in the field of nutrition. Nutraceutical is the food which can be used as preventing drug or food supplement. Millets are the underutilized food having nutraceutical value.

The dietary value of bakery products needs enhancement due to growing consumer awareness about nutrition and ideal health. There's an increasing demand for healthier bakery options as people become more conscious of food-related health issues like obesity, diabetes, and heart problems. This article focuses on the chemical composition, health benefits, and functional bakery products developed by incorporating foxtail and barnyard millet flour.

2. Millets

2.1 Foxtail millet

The names for foxtail millet (*Setaria italica*) include Warf, Enormous, Italian, German, Chinese, and Siberian. The small seeds of the foxtail millet have a diameter of 2 mm. After being threshed off, they are linked to a thin paper shell that is readily removed. The seed came in a wide range of hues, from pale yellow to orange, red, brown, and black. There are 2 g in one foxtail millet thousand kernel. Like maize, foxtail millet has a poor amino acid score and is deficient in lysine. In certain varieties, the quantities of leucine and methionine might exceed 100%. Barnyard, proso, and foxtail starches are more readily absorbed than maize starch. Ash is more prevalent than cereal grains. Dehulling resulted in a notable decrease in the amount of ash.

Starch, protein, vitamins, and minerals make up the majority of the nutrients in foxtail millet. Coarse grains of foxtail millet have 79 % digestible protein and undigestible part contains fibre and antinutritional compounds. High fibre contents induce bowel movement, increasing laxative effect to improve health of digestive system. Therefore, foxtail is a crucial element for preparing noodles, soup, brewed alcoholic beverages, cereal porridges, and pancakes, in addition to its functional and nutritional qualities. Also, it has anti carcinogenic, hypoglycaemic and hypolipidemic effects.

The value-added product obtained by foxtail millet processing significantly contributory economic development of many Asian and African countries. The foxtail and other millet have potential to care malnutrition problems (Sharma *et al.*, 2018).

2.2. Functional properties of foxtail millet

The wide variety of health-promoting functional compounds makes the foxtail millet unique among the cereals. The compounds present in foxtail millet improve the nutritional and organoleptic attributes like as aroma, flavour, and appearance.

Yang *et al.* (2013) studied that the foxtail millet contains 11.85% crude protein, 2.83 - 4.47 % of crude fat, 65.59-74.12 g/100 g total starch and 0.25-4.31 g/100 g amino acid. Over 80% of millet fat is unsaturated. Linoleic, oleic, palmitic, stearic, and linolenic acids were the primary fatty acids. According to Zhang *et al.* (2015), cultivar variety correlated with fatty acid content in different places. Thus, varietal breeding could increase foxtail millet's fatty acid composition.

2.3. Composition of foxtail millet

Foxtail millet contained 12.30 g/100g protein, 60.90 g/100g carbohydrates, 2.80 mg/100g iron, 250 mg/100g potassium, 15 mg/100g folic acid and 66.50 mg/100g linoleic acid (Table 1) (Hasan *et al.*, 2019).

Foxtail millet has a number of advantageous nutritional qualities. The thick seed coat of foxtail grains contributes to their high fibre content (8 g/100 g), as well as their high protein (12.30 g/100 g), fat (4.30 g/100 g), and mineral content (calcium 31 mg/100 g, phosphorus 290 mg/100 g, and iron content of around 2.80 mg/100 g). Foxtail grains are coarse and stand out for these qualities. They also have a significant quantity of antinutritional components such phytates and polyphenols, which prevent the absorption of nutrients (Garwadhiremath, 2011).

2.4. Barnyard millet

The family Poaceae, genus *Echinochloa*, and subfamily Panicoideae are where barnyard millet is classified. The warm and temperate regions of the planet are home to a large number of the 250 annual and perennial species belonging to the genus *Echinochloa* (Bajwaet *al.*, 2015). Most of the species from genus *Echinochloa* grow as weed in major crop field (Kraehmeret *al.*, 2016).

2.5. Composition of barnyard millet

When compared to kodo and foxtail millet, barnyard millet has a lower degree of hardness. There are two layers of epidermis on the seed coat of barnyard millet. The aleurone layer cutinized cell wall had 57-66 % carbohydrates, 6.4-12.2 % fibre, 5-8.5 % protein, 3.5-

4.6 % fat, 2.5-4.0 % ash. The shape of simple starch granules is spherical to polygonal with large 1.2-10 μ m diameter as compared to other millet (Kumari and Thayumanavan., 1998). Different genotypes have grain pericarps that range in colour from straw white to light grey and dark grey (Renganathan *et al.*, 2020).

Singh *et al.* (2022) studied the composition of barnyard millet grain. Data revealed that, barnyard millet contained 6-13 g/100g protein, 55-65.50 g/100g carbohydrates, 1.15-19.50 mg/100g iron, 280-340 mg/100g phosphate, 11-27.10 mg/100g calcium and ash 3.8-4.50 mg/100 g ash (Table 2).

2.6. Nutraceuticals properties of barnyard millet

The nutritive food choice for anaemic women and patients with lifestyle disease is barnyard millet. The barnyard millet had two times more polyphenol and carotenoids than finger millet with several health functions (Panwar *et al.*, 2016). The ethnomedical properties include wound healing ability, biliousness, anti-inflammatory, antibacterial, anti-carcinogenic, and easing constipation-related issues (Kim *et al.*, 2011). So, the barnyard millet is most secured and good choice for making nutritional food products.

2.7. Nutritional significance of barnyard millet

Singh *et al.* (2022) studied the amino acid profile of barnyard millet grain. Data revealed that, barnyard millet had very low lysine content ranging from 106 mg to 136 mg/g. The barnyard millet contained 175-219 mg/g cystine, 288-372 mg/g isoleucine, 725-762 mg/g leucine, 35-263 mg/g threonine, 131-133 mg/g methionine, 119-415 mg/g tyrosine, 388-415 mg/g valine (Table 3).

Barnyard millet may help type II diabetes. Millet has 10.50 % protein, 68.80 % carbs, 3.60 % fat, and 398 kcal/100 g. Dietary fibre, both soluble (4.20 %) and insoluble (8.40 %), was abundant (12.60 %). Dehulled (50.0) and heat-treated (41.70) grains had low glycemic indexes (Ugare *et al.*, 2014).

Foxtail and barnyard millet are equally nutritious as wheat and rice. They outperform rice and wheat in fibre and minerals. Millets have some potential for therapeutic nutrition, although this potential has not yet been fully realised. The development of millet-based foods that are suitable for diabetics has not been the subject of many studies (Pathak *et al.*, 2000).

3. Uses of non-wheat flours in bakery products

Donelson *et al.* (1988) replaced the flour starch with starch tailing or water-soluble fractions and gluten fraction. The researcher investigated the impact of including certain substances on the baking characteristics of cake and cookies. The data revealed that loss of volume of cake was observed with incorporation of starch tailing of water solution fraction for cake making.

While non-significant changes were observed in baking properties with incorporating of gluten fraction for cake making. The baking quality of cookies improved significantly with replacement of gluten or starch tailing fraction with flour starch.

Gaines *et al.* (1985) evaluated the effects of cake flour protein alteration on white layer thickness and softness. The air classification method used to change the protein content of flour by addition of gluten lever from 7 to 16 %. Result revealed that composition of flour did not significantly effect on yield and textural properties i.e. softness of cake. The increased flour protein content significantly effects on height and tenderness of angle cake.

Ammar *et al.* (2009) prepared the balady bread with replacement of taro flour of levels of 5, 10, 15 and 20% in wheat flour. Additionally, researchers looked at the farinograph, extensograph, sensory, and chemical characteristics of bread made with taro flour. The findings of the farinograph analysis indicated that the substitution of taro flour showed an increment in water absorption and dough weakening. However, it also led to a decrease in the time required for dough mixing and its overall stability. The energy required for dough markings, resistance to extension of dough and number of cells decreased with increased concentration of taro flour while extensibility of dough increased in case of extensograph. The bread with taro flours up to 10 % had similar sensory score with control for all attributes. The elevated concentration of tarp flour in bread leads to a reduction in crude protein and crude fat, while simultaneously resulting in an increment in ash, total carbs, and fiber content.

Sanfuleet *al.* (2010) evaluated the effect of addition of soy flour making bread on nutritional value, sensory score, composition of bread. The findings indicate that an elevated concentration of soy flour is associated with an increment of the moisture, ash, and protein levels in bread. The protein level of bread had a notable enhancement, resulting in a substantial improvement in its nutritional worth. The sensory evaluation for soy bread showed that bread with 30% of soybean flour liked more as compared to control while sensory acceptance decreases beyond 30% level of incorporation.

By including pea flour, a protein-rich and fiber-rich ingredient, baked products can be enhanced in terms of their nutritional value. Kamaljitet *al.* (2010) examined pea flour's functional properties for this purpose. The milled dried pea flour was added in bread and cookies at a 5 and 10 % level. The baking quality of bread and cookies were studied. Water absorption capacity increased with reduction in stickiness of dough. Increased level of pea flour causes collapse in volume and increase in weight of bread was observed. Pea flour lowered cookie spread ratio. Bread with 5 % pea flour had similar of sensory score with

control. The inclusion of 5 % pea flour improves the baking quality of cookies and bread without affecting on organoleptic quality.

4. Uses of millets in bakery products

The quality assessment of bakery goods made with pearl millet was researched by Singh *et al.* (2006). They used wheat and pearl millet flour to make sponge cake, eggless cake, and rusk. The millet flour replaced with wheat flour at level of 25, 50 and 75 %. The product prepared from wheat flour replaced with 25 and 50 % of pearl millet flour had overall acceptability between 7.40 and 8.20. The bakery product with a 50% substitution of pearl millet flour with wheat flour was acceptable by the majority of sensory panel assessors. The moisture content, protein content, fat content, ash content, crude fiber content, and carbohydrate content of baked fresh pearl millet items exhibited a range of 3.80-30.20 %, 11.20-17.20 %, 13.50-36 %, 1.1-1.8%, 0.4-0.78 %, and 51.0-67.20 %, respectively.

Anuet *et al.* (2007) prepared the pearl millet-based sponge cake for the survey of dietary assessment. The proportions of blanched pearl millet and green gram meal added to controls with 100 % refined wheat meal were 40:50:10 for Type I and 60:30:10 for Type II. Both varieties of sponge cakes were adored by the panellists. The type I and type II cakes' chemical composition were superior to that of the control cake.

Nazni and Karuna (2016) developed the rusk and muffin incorporated with barnyard millet bran (BMB). In comparison to calcium hydroxide-treated bran, the amount of dietary fibre, protein, and ash was stabilised in hot water-treated bran. The two products with BMB substitution that earned the highest overall acceptability from the panel were the rusk at ratio (85:15) and muffin at ratio (75:25). Increased replacement of BMB at slightly higher levels caused increase in hardness of rusk and muffin. The bran was first filtered and combined with a 20 % solution of 1 % calcium hydroxide before being dried at 50°C and used to make rusk and muffins. The bran was then steeped in boiling water (125°C) for 15 minutes.

Grigelmoet *et al.* (1999) included peach DF at 2, 3, 4, 5, or 10 % to make high-fruit, dietary-fibre muffins. Peach DF (dietary-fibre) holds water well, thus adding it to muffins enhanced moisture content. DF enhanced muffins' hardness, chewiness, and gumminess but not their weight (302 g), height (4.50 cm), springiness (0.880 cm), or cohesiveness (0.460). The little dark color came from peach DF. Consumer panellists rated muffins with peach DF levels of 2, 3, 4, and 5% as nice as the control on the hedonic scale.

Khatri *et al.* (2020) developed muffins with Palmyra sprout flour. In comparison to whole wheat flours (0.52 %) crude fibre content, palmyra sprout flour (SF) had (5.08 %). Since palmyra sprout flour had a lower water activity (0.55) than wheat flour, it was used to

extend muffin shelf life. The muffin that contained 50 % palmyra sprout flour was deemed to be the best. Additionally, palmyra SF muffins had better proximal composition than refined wheat muffins. The chosen muffin, which contained 50 % palmyra sprout flour, had a high nutritional value.

Millets are more nutritious than cereals, yet traditional consumers and low-income people still eat them. Food technologists and engineers have developed mechanised methods for numerous food products due to customer health awareness and benefits. Millets and products were used to make traditional and RTE (ready to eat) cuisines (Jaybhaye *et al.*, 2014).

Microscopy, rheology, and quality of muffin batter were investigated by Rajiv *et al.* (2011) in relation to substituting 0, 20, 40, 60, 80, and 100 % finger millet flour (FMF), emulsifiers, and hydrocolloids for wheat flour. Increased FMF replacement decreased amylograph peak, breakdown, and setback viscosity. Microscopy of muffin batter with FMF exceeding 60 % in blend showed fewer air cells, indicating inadequate air incorporation during mixing. Muffin batter viscosity, volume, density, and score decreased as FMF increased from 0 % to 100 %, however crumb hardness increased. Cake quality suffered above 60 % FMF. Polysorbate-60, hydroxypropyl-methylcellulose, and 60 % FMF enhanced muffin batter viscosity, volume, grain, and texture.

Hernandez *et al.* (2017) examined muffin texture and baking quality with modified cross-linked cassava starch (CCS) as a fat replacer. Based on wheat flour weight, CCS replaced fat at 8, 12, and 16 %. CCS and wheat flour were tested for thermal characteristics, and wheat flour-CCS mixes for pasting and farinography. CCS flour lowered dough stability and development time. CCS gelatinized and retrograded faster than wheat flour. Wheat flour-CCS mixes had lower setback and final viscosities. Weight loss, specific volume, crumb moisture content, and colour score of wheat flour-CCS muffins were non-significant. T8 muffins with 8 % CCS fat replacer had the closest texture to control muffins without CCS. Thus, this muffin and the control muffins were chosen for a 7-day shelf life research. On 2 and 4 days of storage, the T8 muffins had a higher crumb hardness and were equally acceptable. Despite being firmer during storage, consumers liked T8 muffins.

Gornaset *et al.* (2016) tested muffins with 50 g/kg pomace from strawberries, black currants, raspberries, and sour cherries at 140, 180, and 220 °C in conventional and halogen ovens to increase polyphenol stability. Anthocyanins (36-97 % lost) were more unstable than flavanol glycosides (0-21 %). Thermal degradation of ellagitannins and ellagic acid glycosides increased free ellagic acid in strawberry or raspberry muffins. High-temperature

short-baking preserved polyphenols. Enriching bakery items with fruit pomace maximises nutritional value. 50 g/kg apple pomace improved organoleptic quality and nutritional value of muffins compared to controls.

Rupasingheet *et al.* (2008) examined muffin dietary fibre and phenolics after adding apple skin powder (ASP). Apple processing waste contains nutritional fibre and phenolics. 41% total dietary fiber and 52 mg Trolox equivalents g⁻¹ dry weight of oxygen radical absorption capacity (ORAC) are found in blanched, dehydrated, and crushed ASP. ASP incorporation increased muffins dietary fibre, phenolic, and antioxidant content. Bakery recovered 61 %, 57 %, 53 %, 44 %, and 20 % of quercetin glycosides, catechins, chlorogenic acid, phloridzin, and cyanidingalactoside.

5. Baking characteristics of millets-based bakery products

Almaskiet *et al.* (2017) examined the sensory evaluations and in vitro starch digestion of millet muffins high in polyphenols. In lab trials, starch digestion and lowering sugar were measured in millet-based muffins. All three millet muffins released less sugar than a wheat muffin. Kodo millet grain muffins were less appealing than control and finger millet flour muffins. Kodo and finger millet muffin acceptance varied significantly.

Emmanuel *et al.* (2013) investigated millet sponge cake's nutritional and sensory qualities. The formulation with an overall acceptability scores greater than 6.0 was utilised to optimise levels. Cake carbohydrate, fat, and ash improved significantly. Ash increased 2.08±0.36% and fat 33.41±3.32% compared to 1.53% and 30.96% in the control sample. Carbohydrate content decreased 71.41±5.38% compared to control sample of 77.43%.

Kultheet *et al.* (2017) developed and tested cookie physical and textural features using pearl millet flour (PMF). For cookies, Shanti, Dhanshakti, and Pioneer 84M64 pearl millet flour was replaced at 0, 10, 20, 30, 40, and 50% with wheat maida. PMF increased cookie thickness but decreased diameter, spread ratio, and spread factor. PMF boosted a* and lowered L* and b* in cookies. PMF enhanced cookie hardness, breaking strength, and cutting strength. Based on cookie texture and appearance, PMF of Dhanshakti was better than Shanti and Pioneer 86M64.

Shiny *et al.* (2014) produced a high-fibre millet-based biscuit and studied its nutritional and sensory properties. Consumers now prioritise health benefits when eating properly. The high-fibre biscuit recipe used millets, whole grains, flax seed, and spices after several experiments. The newly created high-fiber biscuit was well-accepted during sensory examination, scoring 4.5 out of 5. The new high-fibre cookie contains the most fibre (19.21 g/100 g), the fewest calories (424.6 Kcal/100 g), and the least fat (13.10 g/100 g).

Jyotsna *et al.* (2016) examined millet-based gluten-free muffin rheology, texture, quality, and immunochemical validation. The project aimed to produce gluten-free (GF) wheat flour muffins. WP:FM blends were 100:0, 95:5, 90:10, and 85:15. Peak, hot, and cold paste viscosities decreased from 0 to 15% mix WP concentration. The 90:10 muffin blend had the highest volume and quality ratings. Thus, the 90:10 blend was optimised. Additives improved muffin texture profiles by lowering hardness and increasing springiness. FM: WP 90:10, distilled glycerol monostearate, and hydroxypropyl methylcellulose enhanced muffin quality.

Jadhav *et al.* (2021) examined the physico-sensory and textural properties of composite millet palm jaggery (CMPJ) muffins produced by replacing all-purpose flour (APFS) with CM (0:100, 50:50, 30:70) and cane sugar. Physico-sensory and texture profile analysis (TPA) was performed on CMPJ and cane sugar muffins. Millet flour and wheat flour mixed 70:30 produced a muffin with good sponginess. Palm jaggery muffins had more moisture ($21.84 \pm 0.01\%$) than sugar muffins ($19.58 \pm 0.01\%$). Due to palm jaggery's brown colour, CMPJ muffins had more redness (9.18 ± 0.07 , 12.12 ± 0.22) than APFS muffins (6.56 ± 0.15 , 10.61 ± 0.15). Jaggery muffins showed higher water activity (aw) and lower pH and sensory score than sugar muffins. TPA revealed CMPJ (6270 ± 7.2 g) was tougher than APFS (4729 ± 4.7). CMPJ muffins were safe for 12 days without preservative after microbial investigation (Total plate count, Yeast and mould count). CMPJ muffins without preservative could replace APFS without impacting quality.

A promising bio-compound-rich fruit waste, raspberry and cranberry pomace powder, was tested in muffins by Bajerskaet *et al.* (2016). Quality was tested with raspberry and cranberry pomaces in American-style muffins baked under various conditions. Baking didn't affect control muffin texture or microstructure. Baking obtained 156, 53, 48, 43, and 22% ellagic acid, flavanols, tocopherols, tocotrienols, and anthocyanins. More than flavanols, tocopherols, and anthocyanins, ellagic acid and tocotrienols held up at lower temperatures. Improved samples result from muffins baked at 180°C for 20 min with optimal microstructure, texture, and phytochemical retention.

Mrabet *et al.* (2016) evaluated date fruit (*Phoenix Dactylifera*) fibre concentration (DFC) muffins for quality and antioxidants. Due to poor commercial quality, Tunisian date varieties were underutilised. Steam pre-treatment yields DFC from these fruits. Bakery items used DFCs as antioxidant dietary fibre. 2.5 and 5% flour replacement muffins were made with DFCs from 165 and 180°C treatments. DFC-doughs yielded muffins with a lower volume than the control. Despite lesser cohesion and springiness, DFC-165 muffins were

softest. Samples shared proximal composition. DFC-muffins scored well in sensory evaluation and had more antioxidant capacity than the control. DFC-165 improves baked foods, however its effects on fat rancidity and staling delays need testing.

6. Shelf life study of millet-based bakery products

To increase muffin quality and shelf life, Bhise and Kaur (2015) added psyllium husk, oat, and barley fibre. After adding 5, 10, 15, and 20% oat, psyllium, and barley fibres, muffins were stored at room and refrigerated conditions. With increasing fibre addition, muffin weight increased significantly. Sensory scores were higher for muffins with 10% oat, 10% psyllium, and 15% barley fibres. Under refrigeration, product flavour did not change when free fatty acid concentration (% oleic acid) rose. Control muffins stored under ambient conditions grew faster and showed higher growth than fibre-incorporated muffins. The highest acceptability scores were for muffins made with 10% psyllium fibre (8.37), 15% barley fibre (8.34), and 10% oat fibre (7.95). After fibre integration, muffins lasted 28 days at ambient conditions and 35 days in LLDPE packed refrigerated conditions, however sensory qualities changed after 35 days. Fibres improved muffin shelf life from 21 to 35 days.

Kaur *et al.* (2018) examined wheat flour and wheat atta germinated and ungerminated flaxseed muffins. This study baked muffins with germinated and ungerminated flaxseed flour. Wheat atta muffins and flour were made with 5, 10, 15, 20, 25, and 30% flaxseed meal. Flaxseed increased muffin volume and texture. Organoleptic and textural evaluations favour 15% ungerminated and 10% germinated flaxseed meal. To evaluate the best shelf life, muffins were packed in LLDPE and stored in room and refrigerated conditions for one month. Muffins lasted 15 days at $30\pm 1^{\circ}\text{C}$ and 1 month at $4-6^{\circ}\text{C}$.

7. Conclusion

Millets like foxtail and barnyard millet are valuable bread ingredients due to their nutritional content and functional characteristics. Foxtail millet had 12.3g/100g protein, 60.9g carbs, 2.8 mg/100g iron, 250mg/100g potassium, 15mg/100g folic acid, 66.5 mg/100g linoleic acid, vitamins, and minerals, and may have anti-carcinogenic, hypoglycaemic, and hypolipidemic properties. However, barnyard millet's high polyphenol and carotenoid content helps treat anaemia and lifestyle disorders. Barnyard millet has 6-13g protein, 55-65.5g carbs, 1.15-19.5mg iron, 280-340mg phosphate, 11-27.1mg calcium, and 3.8-4.5mg ash. Foxtail and barnyard millet flours can replace wheat flour in baking recipes to make healthier products.

Millet flours increase protein, fibre, and antioxidant content in baked goods, as well as sensory qualities and customer acceptability.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

References

- Almaski, A., Coe, S., Lightowler, H., & Thondre, S. (2017). Sensory evaluation of polyphenol-rich millet-based muffins and their effect on in vitro starch digestion. *Proceedings of The Nutrition Society*, 76, 227.
- American Association of Cereal Chemists. Approved Methods Committee. (2000). *Approved methods of the American Association of Cereal Chemists*. AACC.
- Ammar, M. S., Hegazy, A. E., & Bedeir, S. H. (2009). Using of taro flour as partial substitute of wheat flour in bread making. *World Journal of Dairy and Food Sciences*, 4(2), 94-99.
- Anu, Sehgal, S., & Kawatra, A. (2007). Use of pearl millet and green gram flours in biscuits and their sensory and nutritional quality. *Journal of Food Science and Technology*, 44 (5), 536-538.
- Bajerska, Mildner-Szkudlarz, S., J., Gornas, P., Segliņa, D., Pilarska, A., & Jesionowski, T. (2016). Physical and bioactive properties of muffins enriched with raspberry and

- cranberry pomace powder: A promising application of fruit by-products rich in bio compounds. *Plant Foods for Human Nutrition*, 71, 165-173.
- Bajwa, A. A., Jabran, K., Shahid, M., Ali, H. H., & Chauhan, B. S. (2015). Eco-biology and management of *Echinochloa crus-galli*. *Crop Protection*, 75, 151-162.
- Bhise, S & Kaur A (2015). Fortifying muffins with psyllium husk fibre, oat fibre and barley fibre to improve quality and shelf life. *Carpathian Journal of Food Science & Technology*, 7 (2) 5-16.
- Donelson, J. R. (1988). The contribution of high-protein fractions from cake and cookie flours to baking performance. *Cereal Chemistry*, 65 (5), 389-391.
- Emmanuel, K., & Sackle, A. (2013). Nutritional and sensory analysis of millet-based sponge cake. *International Journal of Nutrition and Food Science*, 2 (6), 287-293.
- Gaines, C. S., & Donelson, J. R. (1985). Influence of certain flour quality parameters and post milling treatments on size of angel food and high ratio white layer cakes. *Cereal Chemistry*, 62 (1), 60-63.
- Garwadhiremath, A. (2011). Development of foxtail millet-based breakfast muffin. *Master of Home Science Thesis*: University of Agricultural Sciences, Dharwad.
- Gomashe SS. (2017). Barnyard millet: present status and future thrust areas. *Millet Sorghum Biology and Genetic Improvement*. (28) 184-98.
- Gornas, P., Juhnevica-Radenkova, K., Radenkova, V., Misina, I., Pugajeva, I., Soliven, A., & Segliņa, D. (2016). The impact of different baking conditions on the stability of the extractable polyphenols in muffins enriched by strawberry, sour cherry, raspberry or black currant pomace. *LWT-Food Science and Technology*, 65, 946-953.
- Grigelmo., Miguel, N., Carreras-Boladeras, E., & Martin-Belloso, O. (1999). Development of high-fruit-dietary-fibre muffins. *European Food Research and Technology*, 210, 123-128.
- Gupta, N., Srivastava, A. K., & Pandey, V. N. (2012). Biodiversity and nutraceutical quality of some Indian millets. *Proceedings of the National Academy of Sciences, India Section B: Biological Science*, 82 (2), 265-273.
- Hasan, M., Maheshwari, C., Garg, N. K., & Kumar, M. (2019). Millets: Nutri-cereals. *Biotech Express*, 69 (6), 18-20.
- Hernandez, Rodriguez-Sandoval, E., Prasca-Sierra, I., & V. (2017). Effect of modified cassava starch as a fat replacer on the texture and quality characteristics of muffins. *Journal of Food Measurement and Characterization*, 11, 1630-1639.

- Jadhav, S., Kavinya, V., Nirmal, R. V., Shameem, H. M., & Ramalakshmi, K. (2021). Physico-sensory and Textural Properties of Composite Millet Palm Jaggery Muffins. *Journal of Natural Remedies*, 37-43.
- Jaybhaye, R. V., Pardeshi, I. L., Vengaiah, P. C., & Srivastav, P. P. (2014). Processing and technology for millet-based food products: a review. *Journal of Ready to Eat Food*, 1 (2), 32-48.
- Jyotsna, R., Soumya, C., Swati, S., & Prabhasankar, P. (2016). Rheology, texture, quality characteristics and immunochemical validation of millet-based gluten free muffins. *Journal of Food Measurement and Characterization*, 10, 762-772.
- Kalyani, B., & Manjula, K. (2014). Food irradiation- technology and application. *International Journal of Current Microbiology and Applied Sciences*, 3 (4), 549-555.
- Kamaljit, K., Baljeet, S., & Kaur, A. (2010). Preparation of bakery products by incorporating pea flour as a functional ingredient. *American Journal of Food Technology*, 5 (2), 130-135.
- Kaur, A., Kaur, R., & Bhise, S. (2018). Baking and sensory quality of germinated and ungerminated flaxseed muffins prepared from wheat flour and wheat atta. *Journal of the Saudi Society of Agricultural Sciences*, 19 (1), 109-120.
- Khatri, R. M., Siddiqui, S., Nagamaniammai, G., & Athmaselvi, K. A. (2020). Development of Muffin Using Palmyra (*Borassus flabellifer*) Sprout Flour. *International Journal of Nutrition, Pharmacology, Neurological Diseases*, 10 (1), 14-20.
- Kim, J. Y., Chang, J. K., Park, B. R., & Han, S. I. (2011). Physicochemical and antioxidative properties of selected barnyard millet (*Echinochloa utilis*) species in Korea. *Journal of Food Science and Biotechnology*, 20, 461-469.
- Kraehmer, H., Jabran, K., Mennan, H., & Chauhan, B. S. (2016). Global distribution of rice weeds. *Crop Protection*, 80, 73-86.
- Krishna Kumari, S., & Thayumanavan, B. (1998). Characterization of starches of proso, foxtail, barnyard, kodo and little millets. *Plant Foods Human Nutrition*, 53 (1), 47- 56.
- Kulthe, A. A., Thorat, S. S., & Lande, S. B. (2017). Evaluation of physical and textural properties of cookies prepared from pearl millet flour. *International Journal of Current Microbiology and Applied Sciences*, 6 (4), 692-701.
- Larmond E (1970) Method of sensory evaluation of food. Canadian Department of Agriculture Publications, 1284.

- Mrabet, A., Rodriguez-Gutierrez, G., Rodriguez-Arcos, R., Guillen-Bejarano, R., Ferchichi, A., Sindic, M., & Jimenez-Araujo, A. (2016). Quality characteristics and antioxidant properties of muffins enriched with date fruit (*Phoenix dactylifera L.*) fibre concentrates. *Journal of Food Quality*, 39 (4), 237-244.
- Nazni, P., & Karuna, T. D. (2016). Development and quality evaluation of barnyard millet bran incorporated rusk and muffin. *Journal of Food and Industrial Microbiology*, 2 (116), 2.
- Panwar, P., Dubey, A., & Verma, A. (2016). Evaluation of nutraceutical and antinutritional properties in barnyard and finger millet varieties grown in himalayan region. *Journal of Food Science and Technology*, 53, 2779-2787.
- Pathak, Sarita Srivastava, Sema Grover, P. (2000). Development of food products based on millets, legumes and fenugreek seeds and their suitability in the diabetic diet. *International Journal of Food Sciences and Nutrition*, 51 (5), 409-414.
- Rajiv, J., Soumya, C., Indrani, D., & Venkateswara Rao, G. (2011). Effect of replacement of wheat flour with finger millet flour (*Eleusinecorcana*) on the batter microscopy, rheology and quality characteristics of muffins. *Journal of Texture Studies*, 42 (6), 478-489.
- Rao, B. R., Nagasampige, M. H., & Ravikiran, M. (2011). Evaluation of nutraceutical properties of selected small millets. *Journal of Pharmacy and Bio allied Sciences*, 3 (2), 277.
- Reddy, M., Shivkumara, C. S., & Aneesha (2019). Physico-chemical properties of different millets and relationship with cooking quality of millets. *International Journal of Multidisciplinary Research and Development*, 6(4), 97-101.
- Renganathan, V. G., Vanniarajan, C., Karthikeyan, A., & Ramalingam, J. (2020). Barnyard millet for food and nutritional security: current status and future research direction. *Frontiers in Genetics*, 11, 500.
- Renganathan, V. G., Vanniarajan, C., Nirmalakumari, A., Raveendran, M., & Thiyareshwari, S. (2017). Cluster analyses for qualitative and quantitative traits in barnyard millet (*Echinochloafrumentacea*) germplasm. *Bioscan*, 12, 1927- 1931.
- Rupasinghe, H. V., Wang, L., Huber, G. M., & Pitts, N. L. (2008). Effect of baking on dietary fibre and phenolics of muffins incorporated with apple skin powder. *Food Chemistry*, 107 (3), 1217-1224.

- Sanful, R. E., &Darko, S. (2010). Utilization of soybean flour in the production of bread. *Pakistan Journal of Nutrition*, 9 (8), 815-818.
- Sehgal, S., &Kawatra, A. (2007). Use of pearl millet and green gram flours in biscuits and their sensory and nutritional quality. *Journal of Food Science and Technology Mysore*, 536-538.
- Sharma, N., &Niranjan, K. (2018). Foxtail millet: Properties, processing, health benefits, and uses. *Food Reviews International*, 34 (4), 329-363.
- Sharma, N., &Niranjan, K. (2018). Foxtail millet: properties, processing, health benefits and uses. *Food Reviews International*, 34 (4),1-37.
- Shiny-Lizia, M., & John, S. (2014). Sensory and nutritional properties of millet based high fibre biscuit. *International Journal of Science and Research*, 3, 1824-1827.
- Singh, A., Bharath, M., Kotiyal, A., Rana, L., &Rajpal, D. (2022). Barnyard millet: the underutilized nutraceutical minor millet crop. *The Pharma Innovation Journal*, 11 (6), 115-128.
- Singh, K. P., Mishra, A.,& Mishra, H. N. (2012). Fuzzy analysis of sensory attributes of bread prepared from millet-based composite flours. *Journal of FoodScience and Technology*, 48, 276-282.
- Singh, R., G., & Chauhan, G. S. (2000). Development of soy-fortified biscuits and shelf-life studies. *Journal of Food Science and Technology*, 37 (3), 300-303.
- Singh, S., &Goyal, M. (2006). Quality evaluation of pearl millet-based convenience baked products. Sorghum Improvement Conference of North America, USA; *International Crops Research Institute for the Semi-Arid Tropics (ICRISAT)*.
- Sood S, Khulbe RK, Kumar A, Agrawal PK, Upadhyaya HD. (2015). Barnyard millet global core collection evaluation in the sub montane *Himalayan* region of India using multivariate analysis. *The Crop Journal*. 3 (6) 517-25.
- Sunil, C. K., Venkatachalapathy, N., Shanmuga, S. S., Akash, P., &Loganathan, M. (2016). Engineering properties of foxtail millet (*Setariaitalica*). *International Journal of Environmental Science and Technology*, 5 (2), 632-637.
- Truswell, A. S. (2002). Cereal grain and coronary heart disease. *European Journal of Clinical Nutrition*, 56(1), 1-14.

- Ugare, R., Chimmad, B., Naik, R., Bharati, P., &Itagi, S. (2014). Glycaemic index and significance of barnyard millet (*Echinochloafrumentaceae*) in type II diabetics. *Journal of Food Science and Technology*, 51, 392-395.
- Yang, X. S., Wang, L. L., Zhou, X. R., Shuang, S. M., Zhu, Z. H., Li, N., ... & Dong, C. (2013). Determination of protein, fat, starch, and amino acids in foxtail millet [*Setariaitalica (L.) Beauv.*] by Fourier transform near-infrared reflectance spectroscopy. *Food Science and Biotechnology*, 22 (6), 1495-1500.
- Zhang, A., Liu, X., Wang, G., Wang, H., Liu, J., Zhao, W., & Zhang, Y. (2015). Crude fat content and fatty acid profile and their correlations in foxtail millet. *Cereal Chemistry*, 92 (5), 455-459.

Table 1. Composition of foxtail millet (values/100 g)

Parameter	Composition (per 100g)
Energy	331 (kcal)
Carbohydrates	60.9g
Crude fibre	8g
Fat	4.3g
Protein	12.3g
Thiamine	0.59mg
Niacin	3.2mg
Riboflavin	0.11mg
Vitamin A	32mg
Folic acid	15mg

Vitamin Bs	0.82mg
Vitamin E	31mg
Iron	2.8mg
Zinc	2.4mg
Magnesium	81mg
Sodium	4.6mg
Potassium	250mg
Copper	1.4mg
Manganese	0.6mg
Palmitic	6.4mg
Stearic	6.3mg
Oleic	13mg
Linoleic	66.5mg

Hasan *et al.* (2019)

Table 2. Composition of barnyard millet (values/100 g)

Nutrient	Composition (per 100g)
Protein	6 - 13 g
Carbohydrates	55 - 65.5 g
Fat	2 - 4 g
Crude fibre	9.5 - 14 g
Thiamine	0.30 mg
Riboflavin	0.09 mg
Calcium	11 - 27.1 mg
Iron	1.15 – 19.5 mg
Ash	3.8 -4.5 mg
Phosphate	280 - 340 mg

Zinc	2.6 – 4.75 mg
Manganese	1.33 – 3.13mg
Magnesium	82 mg

Singh *et al.* (2022)

Table 3. Amino acid composition of barnyard millet

Amino acid	Composition (mg / g)
Cystine	175 - 210 mg
Isoleucine	288 - 372 mg
Leucine	725 - 762 mg
Lysine	106 - 136 mg
Methionine	131 - 133 mg
Phenylalanine	175 - 210 mg
Threonine	35 – 263 mg
Tryptophan	63 mg
Tyrosine	119 - 150 mg
Valine	388 - 415 mg

Singh *et al.* (2022)




Figure 1. Longitudinal section of a foxtail millet grain



