

## Efficacy of neem oil and selected plant extracts against anthracnose of soybean caused by *Colletotrichum truncatum* (Schw.) Andrus and Moore

### ABSTRACT

Soybean [*Glycine max* (L.) Merrill.] is one of the most important an Asiatic oil seed crop. Soybean comes under the leguminous crop is having symbiotic association with *Rhizobium*. It is a significant commercial crop that yields oil and is beneficial for both oil extraction and the production of numerous items that are used in human and animal nutrition. The present study was investigated on the effect of Neem oil, Eucalyptus leaf extract, Parthenium leaf extract, Lantana leaf extract, Darek leaf extract, Ficus leaf extract and Mancozeb were compared and tested against anthracnose leaf spot [*Colletotrichum truncatum* (Schw.) Andrus and Moore] of soybean under field conditions at the research plot of the Central Research Field, Department of Plant Pathology, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during the *Kharif* season of 2023. Among the treatments, T<sub>3</sub> – Eucalyptus leaf extract exhibited the minimum disease intensity (%) 50 DAS (10.66 %), 70 DAS (11.50 %) and 90 DAS (12.60) Significantly increased in T<sub>3</sub> – Eucalyptus leaf extract @ 10%, followed by T<sub>1</sub> – Parthenium leaf extract @ 10% as compared to other treatments including (Treated check)T<sub>6</sub>-Mancozeb @ 0.2 and (Untreated check) T<sub>0</sub> – Control.

**Key words:** Soybean, *Colletotrichum truncatum* (Schw.), Darek leaf extract, Eucalyptus leaf extract, Ficus leaf extract, Lantana leaf extract, Mancozeb, Neem oil and Parthenium leaf extract.

**Commented [S1]:** Removed an

**Commented [S2]:** crops

**Commented [S3]:** removed on

**Commented [S4]:** must be italic

**Commented [S5]:** must be italic

**Commented [S6]:** must be italic

**Commented [S7]:** must be italic

**Commented [S8]:** must be italic

**Commented [S9]:** of

**Commented [S10]:** ,

**Commented [S11]:** Add spce

**Commented [S12]:** Add six keywords and arrange alphabetically

## INTRODUCTION

Soybean [*Glycine max* (L.) Merrill.] is one of the most important an Asiatic oil seed crop. Soybean comes under the leguminous crop is having symbiotic association with Rhizobium. It is a significant commercial crop that yields oil and is beneficial for both oil extraction and the production of numerous items that are used in human and animal nutrition. As a result, it has spread throughout the entire world, including Europe. Originating in the temperate northern plains of China circa 1100 BC. Although the USA is a major producer of soybeans and their products, India depends heavily on its economy. Soybean a legume crop and is the second largest after groundnut oilseed in India. It is growing in diverse agro-climatic conditions. It is an important rainfed agricultural oil seed crop in the tropical and subtropical regions.

Soybean ranks first among the oilseeds in the world and contributes for nearly 25% of the world's total oil and fats production. Currently, soybean is cultivated over an area of 11.00 million ha and the production is 11.53 million tonnes registering a productivity of 10.45 q/ha (Online Agriculture Statistics, 2023). The USA leads in terms of area and production of soybean, while India ranks fourth in area and fifth in production in the world. USA, Argentina, Brazil, China and India are the major producers of soybean accounting for 90% of world production.

Major states in the nation that grow soybeans are Madhya Pradesh, Maharashtra, Karnataka, and Andhra Pradesh. Profitable soybean production depends critically on the health of the bean plants. The most prevalent species found on soybeans is *Colletotrichum truncatum*, which can affect soybeans at any point throughout their growth, but especially from bloom to pod fullness (Chavan and Suryawanshi, 2016).

Nutritionally soybean is very good source of protein, near about 42 % protein, 20 % oil and 31.3 % carbohydrates content occur in soybean seed. The protein quality of Soybean is equal to the meat, fish and egg and it is very much nutritious produces a lot of products as like soya milk, soya cake, soya peat, cattle feed etc. As well as it is used for the making bread, biscuits, cake,

Commented [S13]: Removed

Commented [S14]: Add reference

Commented [S15]: Add reference

Commented [S16]: is

Commented [S17]: removed the

Commented [S18]:

Commented [S19]: Add reference

Commented [S20]: add space

Commented [S21]: tones

Commented [S22]: Add reference

Commented [S23]: soybeans

Commented [S24]: Add reference

Commented [S25]: soybeans

Commented [S26]: not bould

Commented [S27]: carbohydrate

Commented [S28]: Soybeans

Commented [S29]: Add comma

Commented [S30]: Add reference

flour, etc. It is also called as poor man's meat because highest content of protein and other valuable essential components such as Vitamin C, A, E, B1, B2, B3, B5, B6, B12 and Folic acid etc.

Commented [S31]: Add reference

Soybean plant is highly able to fix the atmospheric nitrogen in to the soil nitrogen and it is the key that influence plant nutrients in crop production. Nitrogen requirement of the plant is very much high but chemically supplying of the nitrogen is very costly, but biologically it is fixed by different bio-inoculants and it is beneficially and eco-friendly for the crops. It will increase root nodulation, branch number, dry matter pod number as well as crop yield. Not only root nodules but also leaves residual effect for crop succeeding equal to the 35 to 40 kg nitrogen per hectare.

Commented [S32]: are

Commented [S33]: The

Commented [S34]: removed

Commented [S35]: Add reference

Commented [S36]: Add reference

A variety of pathogens, bacteria, viruses, nematodes, mycoplasma, including fungi, attack soybean. So far, 29 fungal, 6 bacterial, 18 viral, 6 nematodes and 3 mycoplasma diseases have been recorded on this crop (Sinclair and Dhingra, 1975; Sharma et al., 2016). Soybean cultivation is often subjected to several biotic stresses of which diseases like Anthracnose caused by (*Colletotrichum truncatum*), Fusarium wilt (*Fusarium solani*), Alternaria blight (*Alternaria tenuis*) are important. Among them, anthracnose/ pod blight of soybean caused by *Colletotrichum truncatum*., rust, which is caused by *Phakopsora pachyrhizi*, Rhizoctonia aerial blight, target leaf spot, which is caused by *Rhizoctonia solani*, Myrothecium leaf spot, which is caused by *Myrothecium roridum*, charcoal rot, which is caused by *Macrophomina phaseolina*, *Sclerotinia Xanthomonas axonopodis* pv. *Glycines* and *Xanthomonas campestris* pv. *Glycines* causes purple seed stain; Soybean mosaic virus (SMV); Soybean chlorotic mottle virus (SBCMV) causes soybean mosaic; and Bean Yellow Mosaic virus (BMW) causes bean yellow mosaic. In India it was considered the most serious disease (Khare and Chako, 1983). The disease causes reduction of seed germination, seed quality and yield losses occur in warm (20-25°C) and humid region of sub-tropics and tropics. In India, soybean anthracnose caused 16-25% of yield loss (Nataraj et al., 2020).

Commented [S37]: The soybean

Commented [S38]: to

Commented [S39]: Removed-

Commented [S40]: Add reference

Commented [S41]: disease

Commented [S42]: water-soaked

Commented [S43]: and shrivel

Commented [S44]: Add reference

Soybean crop is susceptible for anthracnose at all stages of development. Pre- and post-emergence damping - off occurs when infected seeds are planted. On emerging seedlings, dark brown, sunken lesions develop on the cotyledons. These lesions can extend along the stem when conditions favor diseases development. Causing one or both cotyledons to become water soaked, shrivel. Early season infection of pods or pedicels can result in fewer and smaller seed or no seed development, at advanced stages of diseases development, near soybean maturity, black fungal

Commented [S45]: disease

fruiting bodies called **acervulli** that produce minute black spines are abundant and randomly distributed on infected tissue (Sikora *et al.*, 2014).

Commented [S46]: acervuline

## MATERIAL AND METHODS

The details of the materials used and methods followed for various **experiment** are described here in the following paragraphs. Leaves and pods exhibiting typical symptoms of anthracnose disease of soybean were collected separately from the field-grown soybean seeds of JS-335 variety plants from the field experiment **was** carried out at the Central Research Field, Department of Plant Pathology, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj during *Kharif* season 2023.

Commented [S47]: experiments

Commented [S48]: was removed

Commented [S49]: and

### Isolation of fungal organism

Commented [S50]: Isolation of the fungal organism

Diseased samples **collected** from different areas during the season and isolation of pathogen was carried out in the laboratory. Firstly, **collected** diseased samples washed thoroughly under the tap water and then cut into small pieces 2-4 mm in size with the help of a sterilized blade in such a way that the sample contained a 50 per cent healthy portion as well as a 50 **per cent** diseased **portion**. The surface of the pieces was sterilized by using 1 **per cent** Sodium hypochlorite solution for 30 seconds to 1 minute, then finally **wash** well with the three changes of sterilized distilled water and to remove excess water then pieces **was** placed on blotter paper. With the help of a sterilized inoculating needle place the sample pieces on petri plates containing potato dextrose agar medium under the aseptic conditions in the laminar airflow chamber (**Plate 3**). Five pieces on PDA media on each plate. Inoculated Petri plates **kept** in an incubator at  $25^{\circ}\pm 2^{\circ}\text{C}$  and **examine** at frequent intervals to check the growth of the target fungal pathogen (Desai and Prasad, 1955).

Commented [S51]: were

Commented [S52]: the

Commented [S53]: percent

Commented [S54]: add refernce

Commented [S55]: percent

Commented [S56]: washed

Commented [S57]: were

Commented [S58]: are

Commented [S59]: examined

### Evaluation of treatments under field conditions

The efficacy of fungicide and **plant-extracts** against *Colletotrichum truncatum* was carried out in field conditions.

Commented [S60]: plant extracts

### Preparation of plant-extracts

Hundred grams of fresh healthy plant parts (leaves) collected from field was washed with distilled water and air dried and crushed in 100 ml of distilled water (w/v). The crushed product was filtered through double layer, muslin cloth and further filtrated through Whatsmann No. 1 filter paper using funnel and volumetric flasks (100 ml cap.). The prepared solution was 100 per cent concentration, which was further diluted to required concentrations of 10 per cent (Shekhawat and Prasada, 1971) (Plate 1). All the treatments were given as foliar sprays. Plant-extracts were sprayed @ 1 and 10 ml/litre of water, mancozeb @ 2 ml/ litre of water.

Commented [S61]: 100

Commented [S62]: were

Commented [S63]: a

Commented [S64]: of muslin

Commented [S65]: Whatman

Commented [S66]: a

Commented [S67]: the required

Commented [S68]: liter

Commented [S69]: liter

Plate 1 Preparation of plant-extracts for treatments

Commented [S70]: write A AND B




### Disease intensity (%)

The Percent disease intensity of 5 randomly selected plants was calculated at 50, 70 and 90 DAS. Disease intensity (%) formula is given by Wheeler (1969). It was calculated by using the following formula:

$$\text{Disease intensity (\%)} = \frac{\text{Sum of all disease ratings}}{\text{Total number of ratings} \times \text{Maximum disease grade}} \times 100$$

List 1. Rating Scale for Anthracnose of Soybean [*Glycine max* (L.) Merrill] (Singh et al., 1990)

Scale	Description
0	No of lesions / discoloration
1	1% area covered with lesions/ spots / discoloration
3	1. 1–10% area covered with lesions/ spots / discoloration
5	10. 1—25% area covered with lesions / spots / discoloration
7	25. 1—50% area covered with lesions / spots / discoloration
9	> 50% area covered with lesions / spots / discoloration



**Plate 2 Disease rating scale**

**List 2. Experimental Details:**

Name of the selected crop	Soybean
Variety	JS-335
Spacing	15 cm
Row to Row	30 cm
Plant to plant	30 cm x 15 cm
Experimental design	RBD
Number of treatments	7
Number of replications	3
Total number of plots	21

Plot size	2 × 1 = 2 m <sup>2</sup>
Size of bunds	0.2 m
Main irrigation channel	1m
Width of sub irrigation channel	0.5 m
Total net cultivated area	42 m <sup>2</sup>
Date of sowing	15 Sept.2023

Commented [S71]: sub-irrigation

### List 3. Treatment Details:

The treatment details are given below under the following headings:

S.No.	Treatments	Treatment details	
		Seed treatment	Foliar spray
1	T <sub>0</sub>	Control (untreated)	Water spray
2	T <sub>1</sub>	Mancozeb (S.T)	Neem oil @1% + Parthenium leaf extract @10%
3	T <sub>2</sub>	Mancozeb (S.T)	Neem oil @1% + Lantana leaf extract @10%
4	T <sub>3</sub>	Mancozeb (S.T)	Neem oil @1% + Eucalyptus leaf extract @10%
5	T <sub>4</sub>	Mancozeb (S.T)	Neem oil @1% + Darek leaf extract @10%
6	T <sub>5</sub>	Mancozeb (S.T)	Neem oil @1% + Ficus leaf extract @10%
7	T <sub>6</sub>	Mancozeb 0.2% (F.S)	

## RESULTS AND DISCUSSION

### Disease intensity (%) of Anthracnose leaf spot on soybean at 50 DAS

The data presented in the table 1 and depicted in figure 1 reveals that disease intensity (%) of soybean significantly decreased in treatment T<sub>3</sub> – Neem oil + Eucalyptus leaf extract (10.66 %) followed by T<sub>1</sub> – Neem oil + Parthenium leaf extract (11.50 %), T<sub>2</sub> – Neem oil + Lantana leaf extract (12.00 %), T<sub>4</sub> – Neem oil + Darek leaf extract (13.00 %) and T<sub>5</sub> - Neem oil + Ficus leaf extract (14.63 %) as compared to (Treated check) T<sub>6</sub> – Mancozeb (8.73 %) and (Untreated check)

T<sub>0</sub> – Control (15.50 %). Comparing the treatments with CD value (0.21), all the treatments (T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, T<sub>4</sub>, T<sub>5</sub>, T<sub>6</sub>) were found significant over (Untreated check) T<sub>0</sub> – Control.

#### Disease intensity (%) of Anthracnose leaf spot on soybean at 50, 70 and 90 DAS

**Table 1 Effect of treatments on disease intensity (%) of Anthracnose leaf spot on soybean at 50, 70 and 90 DAS**

Commented [S72]: ADD COMMA

Treatments	Treatment details		Disease intensity (%)		
	Seed treatment	Foliar spray	Before spray of botanicals	10 days after 1 <sup>st</sup> spray of botanicals	10 days after 2 <sup>nd</sup> spray of botanicals
			50 DAS	70 DAS	90 DAS
T <sub>0</sub>	Control (untreated)	Water spray	15.50 <sup>a</sup>	17.06 <sup>a</sup>	19.86 <sup>a</sup>
T <sub>1</sub>	Mancozeb (S.T)	Neem oil @1% + Parthenium leaf extract @10%	11.50 <sup>e</sup>	12.10 <sup>e</sup>	13.23 <sup>e</sup>
T <sub>2</sub>	Mancozeb (S.T)	Neem oil @1% + Lantana leaf extract @10%	12.00 <sup>d</sup>	13.10 <sup>d</sup>	14.66 <sup>d</sup>
T <sub>3</sub>	Mancozeb (S.T)	Neem oil @1% + Eucalyptus leaf extract @10%	10.66 <sup>f</sup>	11.50 <sup>f</sup>	12.60 <sup>f</sup>
T <sub>4</sub>	Mancozeb (S.T)	Neem oil @1% + Darek leaf extract @10%	13.00 <sup>c</sup>	14.10 <sup>c</sup>	15.13 <sup>c</sup>
T <sub>5</sub>	Mancozeb (S.T)	Neem oil @1% + Ficus leaf extract @10%	14.63 <sup>b</sup>	15.46 <sup>b</sup>	19.76 <sup>a</sup>
T <sub>6</sub>	Mancozeb 0.2% (F.S)		8.73 <sup>g</sup>	9.23 <sup>g</sup>	10.20 <sup>g</sup>
	<b>SE(m) ± 1</b>		<b>0.07</b>	<b>0.98</b>	<b>0.99</b>
	<b>CD at 5%</b>		<b>0.21</b>	<b>0.23</b>	<b>0.25</b>


#### Disease intensity (%) of Anthracnose leaf spot on soybean at 70 DAS

The data presented in the table 1 and depicted in figure 1 reveals that disease intensity (%) of soybean significantly decreased in treatment T<sub>3</sub> – Neem oil + Eucalyptus leaf extract (11.50 %)

followed by T<sub>1</sub> – Neem oil + Parthenium leaf extract (12.10 %), T<sub>2</sub> – Neem oil + Lantana leaf extract (13.10 %), T<sub>4</sub> – Neem oil + Darek leaf extract (14.10 %) and T<sub>5</sub> - Neem oil + Ficus leaf extract (15.46 %) as compared to (Treated check) T<sub>6</sub> – Mancozeb (9.23 %) and (Untreated check) T<sub>0</sub> – Control (17.06 %).

### Disease intensity (%) of Anthracnose leaf spot on soybean at 90 DAS

The data presented in the table 1 and depicted in figure 1 reveals that disease intensity (%) of soybean significantly decreased in treatment T<sub>3</sub> – Neem oil + Eucalyptus leaf extract (12.60 %) followed by T<sub>1</sub> – Neem oil + Parthenium leaf extract (13.23 %), T<sub>2</sub> – Neem oil + Lantana leaf extract (14.66 %), T<sub>4</sub> – Neem oil + Darek leaf extract (15.13 %) and T<sub>5</sub> - Neem oil + Ficus leaf extract (19.76 %) as compared to (Treated check) T<sub>6</sub> – Mancozeb (10.20 %) and (Untreated check) T<sub>0</sub> – Control ( 19.86 %). Comparing the treatments with CD value (0.25), all the treatments (T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, T<sub>4</sub>, T<sub>5</sub>, T<sub>6</sub>) were found significant over (Untreated check) T<sub>0</sub> – Control. Among the treatments (T<sub>6</sub>, T<sub>0</sub>) were found non-significant to each other. (T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, T<sub>4</sub> and T<sub>5</sub>) were found to be significant over all the other treatments.



**Figure 1 Effect of treatments on disease intensity (%) of Anthracnose leaf spot on soybean at 50, 70 and 90 DAS**

Commented [S73]: Must be italic

**Konar and Kushari (1995)** using Eucalyptus extract on different crop plants. **Prasad and Subhashini (1994)** reported that the allelochemicals greatly inhibited porphyrin synthesis. The study results are consistent with the comparable conclusions made by **Patni et al. (2005)**. The Eucalyptus spray had a substantially lower minimum number of spots (30.73) than all other treatments, except the Mancozeb spray. (Table 2). Eucalyptus spray also showed minimum number of spots/ 10cm<sup>2</sup> leaf area (4.16), which was significantly lower than all other treatments but at par with Mancozeb spray (4.25). Eucalyptus spray showed significantly minimum size (5.99mm) of leaf spot in comparison to other treatments except mancozeb spray. Minimum diseases index was observed in Eucalyptus spray (29.05%) followed by Ashok spray (29.31%) and mancozeb spray (29.41%), which were significantly different to all other treatments but at par with each other. Significantly minimum number of spots on pod was observed in Eucalyptus spray (8.26) followed by mancozeb spray (8.60). phyto-extracts of Eucalyptus and neem significantly inhibits the pathogen could lead to better health of the plants which in turn helps in producing minimum disease intensity and leading to a healthier plant.

Commented [S74]: Follow one style

Commented [S75]: italic

Commented [S76]: a

Commented [S77]: Must be italic

Commented [S78]: from

Commented [S79]: he

Commented [S80]: sub-irrigation

Commented [S81]: and

Commented [S82]: Add conclusion

Commented [S83]: Add conflict of interest

## REFERENCES

**Anonymous (2023)**. Online Agriculture Statics. [Http/www.faostal.org](http://www.faostal.org).

**Chavan, S.S. and Suryawanshi, A.P. (2016)**. Bioefficiency of botanicals against *Colletotrichum truncatum*, causing pod blight of soybean. *Journal of Agroecology and Natural Resource Management*, (3):172-174.

**Nataraj, V, Shivakumar, M., Kumawat, G., Gupta, S., Rajput, LS., Kumar, S., Sharma, A.N. and Bhatia, V.S. (2020)**. Genetic inheritance and identification of germplasm sources for anthracnose resistance in soybean [*Glycine max* (L.) Merr.]. *Genetic Resources and Crop Evolution*, 67: 1449-1456.

**Sikora, E. J., Allen, T. W., Wise, K. A., Bergstrom, G., Bradley, C. A., Bond, J. and Zidek, J. (2014).** A coordinated effort to manage soybean rust in North America: A success story in soybean disease monitoring. *Plant of Disease*, 98(7): 864-875.

**Desai, M. V. and Prasad, N. (1955).** A new *Colletotrichum* from India. *Indian Phytopath.*8(1): 52-7.

**Shekhawat, P. S. and Prasada, R. L. (1971).** Antifungal activities of some plant extracts: Inhibition of spore germination. *Indian Phytopathology*, 24: 800-802.

**Wheeler, B. E. J. (1969).** An Introduction to plant disease. John wiley sons limited London. 301.

**Singh, D. P. (1990).** Relative resistance of soybean cultivars to pod blight caused by *Colletotrichum truncatum*. *International Journal of Phytopathology*,11(27): 439.

**Konar, J. and Kushari, D. P. (1995).** Effects of *Eucalyptus globulus* leachates on the growth and diosgenin content of *Costus speciosus*. *Allelopathy Journal*, 2: 215-218.

**Prasad, M. N. V. and Subhashini, P. (1994).** Mimosine-inhibited seed germination, seedling growth and enzymes of *Oryza sativa* L. *Journal of Chemical Ecological*, 20: 1689-1696.

**Patni, C. S., Kolte, S. J. and Awasthi, R. P. (2005).** Efficacy of botanicals against *Alternaria* blight (*Alternaria brassicae*) of mustard Department of Plant Pathology, G.B. Pant University of Agriculture and Technology, Pantnagar. *Indian Phytopathology*, 58 (4): 426-430.

**Commented [S84]:** Please follow journal style

UNDER PEER REVIEW

