

Effect of Different Potting Media on Sprouting, Survival and Growth Performance of Bougainvillea (*Bougainvillea* spp.) Cuttings.

ABSTRACT

The present investigation was carried out during academic year 2023-24 at College of Horticulture, Dapoli, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli, Maharashtra, India. The experiment was laid down in Randomized Block Design (RBD) replicated thrice with eight treatments i.e. T₁ Soil, T₂ Soil + Sand + FYM (1:1:1), T₃ Soil + Sand + Rice husk (1:1:1), T₄ Soil + Sand + Cocopeat (1:1:1), T₅ Soil + Sand + Vermicompost (1:1:1), T₆ Soil + Sand + FYM + Rice husk (1:1:1:1), T₇ Soil + Sand + FYM + Cocopeat (1:1:1:1), T₈ Soil + Sand + FYM + Vermicompost (1:1:1:1). The maximum number of sprouts (3.10), sprouting percentage (95.00), survival percentage (88.33), plant height (56.76 cm), relative growth rate (0.0068 cm/cm/day) were recorded in treatment T₈ i.e. Soil + Sand + FYM + Vermicompost (1:1:1:1).

Keywords: Potting media, bougainvillea, sprouting, survival, relative growth rate.

INTRODUCTION

Bougainvillea is a vibrant and iconic flowering shrubby vine native to south America, popularly cultivated in tropical and subtropical regions worldwide. It imparts beauty to gardens. Belonging to the family Nyctaginaceae. Bougainvillea is a stunning addition to any garden or landscape, with its radiant colours and adaptability making it a popular choice among gardeners and horticulturists. It flourishes well everywhere and even in heavily polluted areas of large cities and industrial towns and that's why it called as 'Glory of the Garden' [1].

Use of different types of rooting media having different characteristics is important for quality rooting in cuttings. Growing media is an essential material for the production of high quality plants grown in containers Problems of nursery growing plant which relate to the growing media are often due to physical characteristics of the media. Soil and FYM have

tendency to become compacted when used in nursery bags. This compaction is often accompanied by reduction in water holding capacity, drainage, aeration, water infiltration rate and even root penetration occur frequently with this compaction. While, plants grown in growing media composed of mix of cocopeat, vermicompost, farm yard manure, sand etc. will exhibit higher growth rates and improved health indicators (such as leaf colour, root development, and resistance to disease) compared to plants grown in traditionally media.

MATERIALS AND METHODOLOGY

The experiment was conducted during the year 2023-24 at College of Horticulture, Dapoli, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli, Dist. Ratnagiri, Maharashtra. The experiment was conducted in the Randomized Block Design with eight treatments and three replications. The planting material i.e. of bougainvillea hardwood cuttings of 15 cm length of pencil size thickness were selected prior to planting.

Eight different growing media were used: T₁ Soil, T₂ Soil + Sand + FYM (1:1:1), T₃ Soil + Sand + Rice husk (1:1:1), T₄ Soil + Sand + Cocopeat (1:1:1), T₅ Soil + Sand + Vermicompost (1:1:1), T₆ Soil + Sand + FYM + Rice husk (1:1:1:1), T₇ Soil + Sand + FYM + Cocopeat (1:1:1:1), T₈ Soil + Sand + FYM + Vermicompost (1:1:1:1). Potting mixtures were prepared with different proportions of media on volume basis and trichoderma was added at the rate 10g per 100 kg. Then as per the treatments, the mixture filled in polythene bags of size 9" x 11". A slanting cut was given at the base of the cuttings and each cutting had about three to four buds. A transverse cut was given at top of each cutting. For keradix powder treatment, the basal end of the cutting was dipped in water and later dipped in keradix powder taken in a beaker. Then the treated cuttings were planted in polybags (9" x 11") containing media as per treatments.

Number of sprouts and sprouting percentage recorded at 60 days after cutting. Survival percentage recorded at 180 days after cutting. Plant height (cm), relative growth rate (cm/cm/day) were recorded at 30 days interval up to 180 days after cutting. The data obtained in the present investigation were statistically analysed by the method suggested by Panse and Sukhatme [2].

RESULT AND DISCUSSION

Number of sprouts:

All treatment indicate significant difference with respect to number of sprouts (Table 1). The highest number of sprouts (3.10) found in treatment T₈ (soil + sand + FYM + vermicompost) which was at par with treatment T₇ (2.80) and T₆ (2.73). While lower number of sprouts recorded in treatment T₁ (2.00). During investigation the maximum number of sprouts observed in soil + sand + FYM + vermicompost (1:1:1:1) combination which might be due to potting media and environmental factors had significant influence on sprouts developed in cuttings and presence of growth promoting factors in vermicompost and FYM which helped in better utilization of stored carbohydrates, nitrogen and other factors. Similar findings was reported by Kapre *et al.* [3] for cape jasmine in media soil + FYM + rice husk , Minj *et al.* [4] for bougainvillea in soil + sand + cocopeat media.

Sprouting percentage:

The statistically highest per cent sprouting (Table 1) was found in treatment T₈ (95.00) (soil + sand + FYM + vermicompost) which was at par with T₆ (93.33), T₇ (91.67), T₄ (90.00) and T₅ (88.33). While lowest per cent sprouting observed in treatment T₁ (83.33). Potting media and the weather condition during experimental period was important for increasing cell the activity for formation of root. Maximum sprouting percentage observed in media containing soil + sand + FYM + vermicompost might be due to FYM and vermicompost contain bioactive principles and balanced composition of nutrients for sprouting and root initiation of cutting. All these factors contributed for sprouting [5].

Similar findings obtained by Kamruzzaman and Quadir [6] in sand for bougainvillea, Kumar *et al.* [7] in soil + sand + vermicompost media in pomegranate.

Survival percentage:

The highest per cent survival of plants (Table 1) in treatment T₈ (88.33) (soil + sand + FYM + vermicompost) which was at par with T₆ (86.67), T₇ (85.00), T₄ (81.67) and T₅ (83.33). While lowest per cent sprouting observed in treatment T₁ (75.00). In present investigation media combination i.e. soil + sand + FYM + vermicompost (1:1:1:1) might have provide physical conditions and sufficient nutrients to cuttings particularly for better metabolic and physiological activities along with better development of root system. Maximum number of roots with higher length and thickness absorb more nutrients and water from the soil and resulted in maximum percentage of survival [8]. The results are in conformity with the findings of Rahman *et al.* [9], Mehmood *et al.* [10], Singh *et al.* [11].




Fig. 1: Effect of different potting media on sprouting and survival percentage of bougainvillea (*Bougainvillea* spp.) cuttings.

Plant height

The maximum plant height (Table 2) was found in treatment T₈ (56.76 cm) (soil + sand + FYM + vermicompost) and followed by treatments T₆ (51.43 cm) and T₇ (50.94 cm) which was found superior over rest of the treatments. The lowest plant height was recorded in treatment T₁ (40.60 cm). Plant height is the most important morphological character of plant. In present investigation, it was found that different media had significant effect on height of cuttings. The media containing FYM and vermicompost produced the highest plant height. It might be due to vermicompost and FYM play important role to improve physical and chemical properties of rooting media by improving its water holding capacity, aeration, organic matter mineralization and solubilisation and availability of micronutrient permitted gaseous exchange between roots and atmosphere which might have increased the growth of cuttings producing more length of shoots. Similar results were also recorded by Seifeldin and Samah [12] in (silty soil 75 per cent + compost 25 per cent) for bougainvillea, Minj *et al.* [4] (soil + sand + cocopeat) for bougainvillea.

Relative growth rate on height basis (cm/cm/day):

The highest relative growth rate on height basis (Table 2) recorded in treatment T₈ (0.0068 cm/cm/day) (soil + sand + FYM + vermicompost) and lowest relative growth rate on height basis observed in treatment T₃ (0.0035 cm/cm/day) (soil + sand + FYM). The maximum relative growth rate on height basis observed in media containing FYM and vermicompost. The combination provides excellent structures, porosity and nutrients in available form such as nitrate nitrogen and soluble phosphorus it might be main reasons for excellent plant growth. The results are in conformity with the findings of Bendre *et al.* [13] in soil + rice husk + compost (2:1:1) in bush pepper, Khedkar *et al.* [14] in soil + FYM (3:1) in cashew graft.

Table 1. **Effect of different potting media on sprouting parameters and survival percentage of bougainvillea (*Bougainvillea* spp.) cuttings.**

Treatments	Number of sprouts (60 DAC)	Sprouting percentage (60 DAC)	Survival percentage (180 DAC)
T ₁	2.00	83.33	75.00
T ₂	2.27	86.67	80.00
T ₃	2.06	85.00	76.67
T ₄	2.06	90.00	81.67
T ₅	2.37	88.33	83.33
T ₆	2.73	93.33	86.67
T ₇	2.80	91.67	85.00
T ₈	3.10	95.00	88.33
Mean	2.42	89.17	82.08
Range	2.00- 3.1	83.33-95.00	75.00-88.33
'F' test	SIG	SIG	SIG
S.Em. ±	0.19	2.41	2.28
C.D. at 5 %	0.56	7.31	6.92

Table 2. Effect of different potting media on plant height (cm) and relative growth rate (cm/cm/day) of bougainvillea (*Bougainvillea* spp.) cuttings at 180 days after cutting.

Treatments	Plant height (cm)	Relative growth rate (cm/cm/day)
T ₁	40.60	0.0046
T ₂	45.13	0.0044
T ₃	41.09	0.0035
T ₄	46.55	0.0041
T ₅	47.97	0.0055
T ₆	51.43	0.0053
T ₇	50.94	0.0053
T ₈	56.76	0.0068
Mean	47.56	0.0049
Range	40.60-56.76	-
'F' test	SIG	-
S.Em. ±	1.11	-
C.D. at 5 %	3.37	-

CONCLUSION

From present investigation it can be concluded that among different treatments, T₈ (soil + sand + FYM + vermicompost) (1:1:1:1) treatment recorded the maximum number of sprouts, sprouting percentage, survival percentage, plant height and relative growth rate. Soil + sand + FYM + vermicompost media combination gives best results for better growth performance of bougainvillea.



REFERENCES

1. Sindhu, S. S. and Swaroop, K. (2015). Bougainvillea repository at IARI. *Indian Bougainvillea Annual*. **26**(3): 27-29.
2. Panse, V. G. and Sukhatme, P. V. (1985). Statistical Methods for Agricultural Workers. *ICAR Rev. Ed.* 97-156.
3. Kapre, G. C., Thorat, S. B., Dalvi, N. V., Khobragade, N. H., Malshe K. V. and Swarda Parab. (2022). Performance of cape jasmine (*Gardenia jasminoids* Ellis) to different propagation media on survival by stem cuttings. *The Pharma Innovation J.*, **11** (12): 3885-3887.
4. Minj, S. K., Gupta, P., Sahu, R. K. and Khilari, M. (2023). Standardization of media for propagation of bougainvillea (*Bougainvillea comm.* Ex Juss). *The Pharma Innovation J.*, **12**(5): 927-930.
5. Bachman, G. and Metzger, J. (2008). Growth of bedding plants in commercial potting substrate amended with vermicompost. *Bioresources Technology*, **99**(8): 3155-3161.
6. Kamruzzaman, M., and Quadir, M. A. (1998). Rooting performance of Bougainvillea as influenced by different types and methods of stem cuttings. *Thailand J. Agri. sci.*, **31** (1): 98-103.
7. Kumar, S., Prakash, S., Malik, S. and Dhayani, B. P. (2021). Effect of different rooting media treatments on rooting and growth of stem cuttings in pomegranate (*Punica granatum* L.). *The Pharma Innovation J.*, **10**(3): 475-479.
8. Gupta, V. N., Banerjee, B. K. and Datta, S. K. (2002). Effect of auxin on rooting and sprouting behaviour of stem cuttings of bougainvillea under mist. *Haryana J. Hort. Sci.* **31**: 42-44.
9. Rahman, N., Hussain, L., Awan, A. A. (1999). Effect of different media on propagation of bi-color bougainvillea cuttings. *Pakistan J. Biologi. Sci.*, **2** (3): 877-878.
10. Mehmood, S., Ayub, Q., Khan, S. M., Arif, N., Khan, M. J., Mehmood, A., Shahzad, Q., Haq, N., Tanoli, M.T.Z. and Ayub, M.U., (2020). Responses of fig cuttings (*Ficus carica*) to different sowing dates and potting media under agro-climatic conditions of Haripur. *RADS J. Biol. Res. Appl. Sci.*, **11**(2) :1-8.

11. Singh, K. K., Singh, S.P., Bairwa, B. and Chauhan, J.S. (2020). Production technology of Bougainvillea spp. *Biotica Research Today*. **2**(6): 472-474.
12. Seifeldin, A. M. and Samah , O. M. M. (2022). Effect of different potting media on growth and flowering of bougainvillea sp. *Nile J. Agri. Sci.*, **7**(1): 1-11.
13. Bendre, P. D. (2019). Effect of different potting media and plant growth regulator on survival and growth of bush pepper (*Piper nigrum* L.) *M.Sc.(Hort.)* thesis submitted to Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth Dapoli, Maharashtra.
14. Khedkar, S. P., Bhingarde, R. T., Chavan, V. B., Chatse,D. B., Khandekar, R. G. and Salvi, B. R. (2023). Effect of potting media on survival and growth performance of rebagged of cashew grafts cv. Vengurla 4. *The Pharma Innovation J.*, **12**(6): 428-432.