

## Original Research Article

# Effect of magnetized irrigation water and seeds on the production of OKRA (*Hibiscus esculentus*)

**Abstract:** Very recently special magnetic devices are developed and used to magnetize irrigation water and plant seeds. Such magnetic treatments were reported to play a great role in increasing the germination rate of seeds, reducing soil salinity and consequently resulting in increasing yield. The objectives of the study to improving crop productivity by using magnetic technologies. A split plot experimental design was used for growing two varieties of okra (*Hibiscus esculentus*). The treatments included: Non magnetized water + None magnetized seeds (NMW+NMS), Non magnetized water + Magnetized seeds (NMW+MS), magnetized water + Non magnetized seeds (MW+NMS), Magnetized water + Magnetized seeds (MW+MS). Each treatment was replicated three times. The parameters considered included: number of leaves per plant, leaf length, plant height, plants density and yield. There was a significant difference ( $p < 0.05$ ) between treatments, the combination of (MW) and (MS) gave higher of: Number of leaves per plant, plant density, and plant height and leaf length. There was a significant difference ( $p < 0.05$ ) in yield between treatments (MW) record increasing in yield by 50% as compared to (NMW), (MW) + (MS) record increasing in yield by 24% as compared to (NMW), (MS) record increasing in yield by 18% as compared to (NMS) when both were irrigated by (NMW), The combination of (MW) and (MS) gave higher yields for both okra (25.5 and 35,5 Kg/ha)

It was concluded that magnetizing water may lead to better crop establishment and further improvement could be achieved by magnetizing the seeds.

**Key word:** *magnetic devices, magnetic technologies, magnetizing water, crop productivity, okra.*

## INTRODUCTION

Irrigation has often been defined as the method of applying water to the soil to supplement that from rainfall for maximizing production per unit area. Irrigated agriculture is the largest consumer of water in the world. In areas with dry climate, irrigation water amount to 50-85% of total water use (Hamdy, 2001). The growing scarcity and misuse of the available water resources, particularly in arid and semi-arid regions, constitute challenges to water demands for various utilities.

Sudan is envisaged to face water shortage very soon. Utilization of water from the river Nile and tributaries is abided by international agreements with neighboring countries. Rainfalls are unpredictable and evaporation losses are high under such arid conditions.

Agriculture in Sudan is either irrigated or rain fed. Irrigated agriculture depends mainly on water from the river Nile and its tributaries. The utilization of the Nile water is controlled by the 1959 Nile water agreement with Egypt (CFA 2010).

The use of magnetic devices for magnetizing irrigation water started in the 1960s, which there has been paid increased attention as an efficient and sustainable utilization method (Kney and Parsons, 2006, Mostafazadeh et al., 2008) Magnetization causes physical and chemical changes of natural water characteristics, resulting in an improvement of its infiltration rate and an increase in its dissolving properties. These changes result in an increased ability of soil to get rid of salts which will consequently result in a better assimilation of nutrients and fertilizers in plants during their growing cycles. Magnetic water is obtained by passing water through permanent magnets or through the electromagnets installed in or on a feed pipeline (Higashitani et al. 1993). Plants need mineral salts and trace elements from the soil for healthy growth and maximum yield. The mechanism of magnetization of water and proposed a theory based on the molecular structure of water Deng and Pang, (2007). Minerals and nutrients must be well balanced in the soil at an appropriate pH level in order to be fully utilized by the plants, (Busch et al, 1997), magnetized technology has great beneficial for soil improvement and plant growth (Ahmed and Bassem, 2013, Higashitani et al., 1993).

Watering plants with magnetized water dissolves more nutrients because it lowers the surface tension of water. This lets more minerals be suspended in concentration. This buffers the pH and causes more minerals and nutrients to pass through the cell walls of plant roots. Magnetized water penetrates the soil faster and deeper, allowing roots to penetrate and grow larger. In addition, the changes caused by the magnetic influence depend on many factors, such as strength of the magnetic field, direction of applied magnetized field, duration of magnetic exposure, flow rate of the solution, additives present in the system, and the pH (Higashitani et al. 1993; Marcus and Rashin 1994; Baker and Judd 1996; Parsons et al. 1997; Gabrielli et al. 2001; Chibowski et al. 2005). Magnetized water dissolves more nutrients within the root zone and become readily available to be up taken by the roots and hence stimulate plant growth and increase water-use efficiency (WUE) and crop yields (Tan and Wang, 2017), These may be the reasons why plant growth rates are increased. Crop yields are higher in a shorter period of time, and with much less needs for water, fertilizers and pesticides. This is the reason why magnetized water is recommended for irrigation. These results in an increased top quality, crop production coupled with savings in cost and reduction in environmental hazards. It has been found that magnetized water treatment can promote seed germination, crop growth, increase yield, and improve fruit quality (Wang et al., 2018a, Wang et al., 2018b). Haq et al. (2016), Gholami and Sharafi (2010) reported that the Magnetic seed treatment enhanced seed germination, seedling growth, fresh and dry weight, activities of some enzymes and seed yield in wheat. On the other hand, Iqbal et al. (2012) obtained that the magnetic field exhibited increased germination of pea seeds.

This study has been conducted with a view to evaluating the effect of magnetizing irrigation water and /or seeds on crop production under Sudan condition, okra (*Hibiscus esculentus*) have been taken as an indicator plant.

## **Material and methods**

The experiment was conducted in the Demonstration Farm of the faculty of Agriculture, University of Khartoum, Shambat which lies at latitude 15° 40' N and longitude 32° 32' E.

The study was carried out during 2020 summer season. The experimental period extended over April – July. The climate of the area is tropical semi-arid, characterized by low relative humidity with mean daily maximum and minimum temperatures of about 36° C and 21° C respectively. The annual rainfall is about 158 mm mainly during July, August and September (Sudan Meteorological Department 1951-1980). During the experimental period of April – June, no rainfall was recorded.

The soil in the experimental field is heavy clay with percentages ranging between 65% in the top 15 cm and 55% in the 100-140 cm depth. The soil reaction is moderately alkaline with a pH ranging from 7 to 8 (Saeed, 1968). The infiltration rate is low and has been estimated to be about 20 mm/hr in the first two hours and 5mm/hr after 10 hours (Ferguson, 1986).

An area of 170 m<sup>2</sup> was divided into two sections and planted with two varieties of okra (Madni and Kosti). One section was irrigated by magnetized water. The treatments were arranged in a split plot design, and each treatment was replicated three times giving a total of 12 plots. The seeds were magnetized either on the dry and when they were soaked in water

#### **Magnetization processes:**

A magnetic funnel and a container were used to soak seeds in, the required amount of water was poured to pass through the magnetic funnel into the container; the seeds were then passed through the magnetic funnel into the container with magnetized water. The seeds were left in the magnetized water for about 30 minutes and then the water was poured out of the container and the seeds were passed through the magnetic funnel again, and were ready for sowing according to Aladjadjiyan, (2007).

Magnetization of the dry seeds was done simply by passing the seeds dry through a magnetic funnel.

#### **Land preparation:**

The experimental field was ploughed with a standard integral mounted disc plough at a depth of about 0.25 m. The land was then left for two weeks then leveled with a general-purpose blade. Ridging was done at a spacing of 0.7 m with a general-purpose ridge.

#### **Planting:**

Okra seeds were sown on 28th April 2020 at a seed rate of about 8Kg per feddan. Two to three seeds were planted manually in rows on the eastern side of the ridge at a spacing of 30 cm between plants.

#### **Irrigation system:**

The experiment was cured out under furrows irradiation system, water source from domestic water supply system with centrifugal pump (2hp).

#### **Data collection:**

Plant growth parameters measured were as follows:

a) Number of leaves per plant:

Nine plants were taken at random from each three ridges and the mean number of leaves per plant was recorded.

b) Density of plant:

Number of plants for each three ridges was counted then the mean density was recorded as number of plants per unit area.

c) Plant height:

Eight weeks from sowing, plants height was measured by a wooden ruler from the base of the stem to the tip of youngest leaf. Nine plants were taken at random from each three ridges, their mean

Height was recorded.

d) Length of leaves:

Nine plants were randomly selected from each three ridges, and the mean length of the leaves was recorded.

e) The yield:

A sensitive balance was used to weigh the end product.

#### **RESULTS AND DISCUSSION**

The experimental results are presented in the tabular form of tables (1& 2) they exhibit the following information:

##### **Number of leaves per plant:**

Table 1 shows that there was no significant difference ( $p < 0.05$ ) in the number of leaves per plant when either or both (MW) + (MS), but a significant difference is indicated when neither one of them was magnetized (NMW) + (NMS).

Both okra varieties (Kosti and Madani) gave higher values of number of leaves per plant when both water and seeds were magnetized (MW)+(MS), they are respectively 17.23 and 12.7 leaves /plant (table 2). these findings were agreed with Ziaf, *et al* (2022)

##### **Plant density:**

Table (1) shows a significant difference ( $p < 0.05$ ) when either or both water and seeds were magnetized (MW) + (MS) as compared to when neither one of them was magnetized. The

combination of magnetized water and magnetized seeds gave higher plant density per m<sup>2</sup> for both okra varieties; they are (14.9 plants /m) for Kosti and (13.87)

Plants/m<sup>2</sup>) for Madani (Table 2). these findings were agreed with Neo E. Nyakane *et al* (2019) and Ziaf, *et al* (2022)

### **Plant height:**

With respect to plant height there was no significant difference when either water or seeds were magnetized. Furthermore, it was found that both okra varieties (Kosti and Madani) gave higher values of plant height for the combination of magnetized water and magnetized seeds; they are (59cm) for Kosti and (57.1 cm) for Madani (Table 2).

### **Leaf length:**

A significant difference ( $p < 0.05$ ) in leaf length was obtained from magnetized water (8.05cm) as compared to (6.15cm) when non magnetized water was used (Table 1)

It was found that when magnetized water was used there was no significant difference ( $p < 0.05$ ) in the leaf length whether the seeds were magnetized or not magnetized (Table 1).

Referring to Table (1) it was noticeable that when non magnetized water was used there was a significant difference ( $p < 0.05$ ) in the leaf length of magnetized seeds (8.0 cm) as compared to non-magnetized seeds (6.15 cm).

The combination of magnetized water and magnetized seeds gave higher averages of leaf length for both okra varieties, they are (9.2 cm) for Kosti and (8.9cm) for Madani (Table 2).

### **Yield:**

There was a significant difference ( $p < 0.05$ ) in yield when okra was irrigated by magnetized water (14.8 Kg/ha) as compared to non-magnetized water (7.4 Kg/ha).

A significant difference ( $p < 0.05$ ) was also obtained with combinations of magnetized water and magnetized seeds (19.4Kg/ha) as compared to a combination of magnetized water and non-magnetized seeds (14.8 Kg/ha) (Table 1).

Table (1) also shows a significant difference in the yield of magnetized seeds (9.0 Kg/ha) as compared to non-magnetized seeds (7.4 Kg/ha) when both were irrigated by non-magnetized water. The combination of magnetized water and magnetized seeds gave higher yields for both okra varieties, namely (25.5 Kg/ha) for Kosti and (35.5 Kg/ha) for Madani (Table 2), these results agree with Sabah L. Alwan, *et al* (2013) and Neo E. Nyakane *et al* (2019) who reported that, using proper combination of magnetic field intensity and time exposure is a fundamental key to consider in order to enhance the crop productivity and development

## CONCLUSION AND RECOMMENDATIONS

From the results of this study the following conclusion and recommendations can be drawn:

- Magnetizing irrigation water can lead to better crop production and further improvement may be attained by magnetizing the seeds.
- The major problem associated with magnetization technology may be the high initial cost for the device and its requirement for special conveyance accessories.
- Since the technology of magnetization has been newly introduced in Sudan and proves to have good potentialities in agricultural production, further research studies are highly recommended in this area.

**Table 1 Analysis of the effects of magnetized and non-magnetized water and seeds on some agronomic parameters of okra crop.**

Parameters	Magnetized water		Non magnetized water		
	Mag. seed	Non mag. seeds	Mag. seeds	Non mag.seeds	LSD
Number of leaves/plants	a 12.12	a 12.12	a 14.97	a 7.85	a 2.88
Plant density (plants/m <sup>2</sup> )	a 12.8	ab 11.45	ab 11.45	b 8.0	3.70
Plant height (cm)	a 48.45	a 45.77	a 43.33	a 39.55	12.07
Leaf length (cm)	a 9.05	a 8.05	a 8.0	b 6.15	1.49
Yield (Kg/ha)	a 19.4	b 14.8	c 9.0	d 7.4	1.38

**Table 2 Mean comparison test:**

Magnetized water					Non magnetized water					
Kosti		Madani			Kosti		Madani			LSD
Mag. Seeds	Non mag. seed	Mag. Seeds	Non mag. seed	Mag. Seeds	Non mag. seed	Mag. Seeds	Non mag. seed			
a 17.23	bc 12.43	bc 12.7	de 9.9	b 14.33	ef 8.9	cd 11.8	f 6.8	2.14		
a 14.9	b 8.0	a 13.87	b 9.1	a 13.6	b 8.0	a 12.0	b 8.0	2.75		
a 59	a 37.9	ab 57.1	cd 39.8	ab 53.63	bc 47.9	de 31.2	e 27.67	8.97		
a 9.2	bc 7.8	Ab 8.9	C 7.1	Ab 8.9	C 9.6	Ab 8.3	d 5.4	1.11		
b 25.5	e 4.3	a 35.5	c 13.9	e 4.2	f 2.5	d 10.6	ef 3.4	1.02		

## References:

- Agreement on the Nile River Basin Cooperative Framework (CFA). 2010. <https://nilebasin.org/images/docs/CFA.pdf>.
- Aladjadjiyan A., 2007. Study of the Influence of Magnetic Field on Some Biological Characteristics of Zea mais. *J. of Central Europ. Agric.* 3(2). 89-94.
- Baker, J. S., and Judd, S. J. (1996). "Magnetic amelioration of scale formation." *Water Res.*, 30(2), 247–260.
- Busch, A.E. & Suessbrich, H. (1997). Role of the IsK protein in the IminK channel complex. *Trends Pharmacol. Sci.*, 18, 26– 29.
- Chibowski, E., Szcześ, A., and Hołysz, L. (2005). "Influence of sodium dodecyl sulfate and static magnetic field on the properties of freshly precipitated calcium carbonate." *Langmuir*, 21(18), 8114–8122.
- Deng, Bo and Pang Xiao Feng, 2007. Variations of optic properties of water under action of static magnetic field, *Chinese Science Bulletin*, Vol.52, No.23, pp. 3179-3182.
- Ferguson, R.I. (1986) River loads underestimated by rating curves. *Water Resources Research*, 22, 74–76. Gifford, G.F. (1976) Applicability of some infiltration formulae to rangeland infiltrometer data. *Journal of Hydrology*, 28,
- Gabrielli, C., Jaouhari, R., Maurin, G., and Keddam, M. (2001). "Magnetic water treatment for scale prevention." *Water Res.*, 35(13), 3249–3259.
- Gholami, A. and Sharafi, S. (2010). Effect of magnetic field on seed germination of two wheat cultivars. *World Academy of Science, Engineering Technology*, 62, 279-282.
- Hamdy A. 2001. Agricultural water demand management is a must for water saving an advanced short course on water saving and increasing water productivity challenges and options. Faculty of Agriculture, University of Jordon, Amman, pp. 618.1-618.30.
- Higashitani, K., Kage, A., Katamura, S., Imai, K., and Hatade, S. (1993). "Effects of a magnetic field on the formation of CaCO<sub>3</sub> particles." *J. Colloid Interface*
- Iqbal, M., Haq, Z.U., Jamil, Y. and Ahmad, M.R. (2012). Effect of pre-sowing magnetic treatment on properties of pea. *International Agrophysics*, 26, 25-31
- K. Higashitani et al. Effects of magnetic field on formation of CaCO<sub>3</sub> particles, *J. Coll. Int. Sci.* (1993).
- Kney A.D, Parsons S.A," A Spectrophotometer-based study of magnetic water treatment: Assessment of ionic vs. surface mechanisms", *Water Res.*, 40, (2006), pp.517.
- Marcus, Y., and Rashin, A. (1994). "A simple empirical model describing the thermodynamics of hydration of ions of widely varying charges, sizes, and shapes." *Biophys. Chem.*, 51(2–3), 111–127.

- Mostafazadeh-Fard, B., Heidarpour, M., Aghakhani, A., and Feizi, M. (2008). "Effects of leaching on soil desalinization for wheat crop in an arid region." *Plant, Soil Environ*, 54(1), 20–29.
- Neo E. Nyakane (2019), The Effects of Magnetic Fields on Plants Growth: A Comprehensive Review, *International Journal of Food Engineering* Vol. 5, No. 1, March 2019, doi: 10.18178/ijfe.5.1.79-87
- Parsons, S. A., Wang, B. L., Judd, S. J., and Stephenson, T. (1997). "Magnetic treatment of calcium carbonate scale-effect of pH control." *Water Res.*, 31(2), 339–342.
- Sabah L. Alwan Ahmed N.Hameed AL-Fakhr Al-din Hayder A. Ali AL-Shebly 2013, The possibility of manufacturing a device for magnetization of water and test its efficiency in seed germination and seedling growth of Pamia, *Kufa Journal of Agricultural Sciences / Volume (4) / Issue (2) 2012*, (175-179
- Saeed, A. M. (1968). Some Physical and Chemical Properties of Certain Soils in Shambat. M. Sc. Thesis. University of Khartoum, Sudan. *Sci.*, 156(1), 90–95.
- Tan S. WangQ.XuD. ZhangJ.ShanY.2017Evaluating effects of four controlling methods in bare strips on soil temperature, water, and salt accumulation under film-mulched drip irrigation. *Field Crops Research*214, 350–358
- Turker, M., C. Temirci, P. Battal and M.E. Erez, 2007. The effects of an artificial and static magnetic field on plant growth, chlorophyll and phytohormone levels in maize and sunflower plants. *Phyton Ann. Rev. Bot.* 46 (2), 271-284.
- Ul Haq, Z.; Iqbal, M.; Jamil, Y.; Anwar, H.; Younis, A.; Arif, M.; Fareed, M.Z.; Hussain, F. Magnetically treated water irrigation effect on turnip seed germination, seedling growth and enzymatic activities. *Inf. Process. Agric.* 2016, 3, 99–106
- Wang H.B. et al.Effect of magnetized water drip irrigation on soil salt and corn yield and qualityJ. *Soil.* (2018) b
- Y. Wang *et al.*Effect of magnetic field on the physical properties of waterJ. *Results Phys.*(2018) a
- Ziafi, M. Amjad, M. A. Ghani, I. Ahmad, M. Ayub, M. Sarwar, Q. Iqbal and M. A. Nawaz, Comparative Efficacy of Magnetic Field Seed Treatment and Priming in Improving Growth and Productivity of Okra 2022, *Journal of Animal & Plant Sciences*, 32(1): 2022, Page: 84-90 ISSN (print): 1018-7081; ISSN (online): 2309-8694 <https://doi.org/10.36899/JAPS.2022.1.0405>