

Original Research Article

Phytochemical analysis and effect of *Myristica fragrans* extract on *Escherichia coli* and *Staphylococcus aureus* strains isolated from Automated Teller Machine, Bikaner, Rajasthan

ABSTRACT

Aims: Plants produce compounds known as secondary plant products, as they are not required for the survival of plants. Earlier it was thought that these compounds were only the waste product of metabolism in plants but later it was found that they have important functions. These phytochemicals possess the ability to combat pathogenic bacteria and lethal diseases. These phytochemicals even don't have the same side effects as modern medications. So the objective was to analyze the phytochemicals and antimicrobial activity of a traditional herb *Myristica fragrans* (nutmeg).

Study design: Quantitative analysis of raw seed powder of *M. fragrans* was done for moisture content and solubility in water and 70% ethanol. Phytochemical analysis was conducted for both water and 70% ethanolic extract. Beside this the antimicrobial activity of both water and ethanolic extract was analyzed for gram-positive (*S. aureus*) and gram-negative (*E. coli*) bacteria isolated from ATM of Bikaner, Rajasthan.

Place and Duration of Study: The study was conducted in the laboratory of MN College and Research Institute, Bikaner, Rajasthan (<https://www.mncollegebkn.com/>) for the period of 2 months (May and June).

Methodology: Moisture content and solubility of *M. fragrans* was determined using the using the procedure given by B.C. O'Kelly and Elena (2012). The extraction of powdered seeds of *M. fragrans* was done following the procedure of Ozaki *et al.* (1989) procedure with slight modification. Alkaloid content of both the extracts was determined using the method given by Harbone (1973). Tannins and phenol contents were analyzed using the method given by Tomar and Shrivastava (2016). Beside this the saponin content was analyzed using two methods (TLC and foam method) given by Ginovyan *et al.* (2020) and Mendhekar *et al.* (2017) respectively. Further the antimicrobial activity of both the extracts for *S. aureus* and *E. coli* were determined using the method given by Ibrahim *et al.* (2013).

Results: Quantitative analysis of raw seed powder showed high moisture content and solubility in water and ethanol. The extraction process in 70% ethanol and water yielded extracts containing alkaloids, saponins, tannins, and phenols in the ethanolic extract, while the water extract contained alkaloids and saponins but lacked tannins and phenols. The ethanolic extract exhibited higher alkaloid content compared to the water extract, with percentages of 13.25% and 7.14% respectively. Notably, the ethanolic extract demonstrated effectiveness against both gram-positive (*Staphylococcus aureus*) and gram-negative (*Escherichia coli*) bacteria, which were of human origin (isolated from ATM).

Conclusion: Our findings suggest that phytochemicals found in seeds of nutmeg have antimicrobial activity and can effectively combat pathogenic bacteria.

Key words: Antimicrobial activity, E. coli, Myristica fragrans, phytochemicals, S. aureus.

INTRODUCTION

The use of natural products to relieve illness dates back thousands of years (Christophersen *et al.*, 1991). Since herbs have healing potential, botanical medicine stands as one of the oldest practised professions by mankind (Van Wyk, and Gerike, 2000). The utilization of traditional medicinal plants can be traced

back 4000 years to Mesopotamia, where seeds and plant extracts were employed for disease prevention and treatment, as documented in clay tablets (Dahham *et al.*, 2018). Approximately 25% of modern prescribed medications are derived from plant-based substances (Hamburger and Hostettmann., 1991). Some of these drugs are synthesized chemically or semi-synthetically, the rest are extracted and purified directly from plants (Farnsworth, 1990). Unlike current therapies which may have limited effectiveness and side effects (Al-Rawi *et al.*, 2024), plants and seeds contain valuable compounds (secondary metabolites and aromatic substances) for treating diseases, which can be used alone or with other medications to enhance efficacy and reduce adverse effects (Ibrahim *et al.*, 2011).

Myristica fragrans (nutmeg) is a tropical, evergreen dioeciously tree commonly found in Penang, Malaysia, India, Indonesia and Southeast of Asia (Orwa *et al.*, 2009). Nutmeg, the dried kernel of the seed of *Myristica fragrans*, is known for its pleasant aroma and warm taste, making it a popular spice in many countries. Hence it is widely used to flavour meats, vegetables, teas, baked goods and more (Al-Rawi *et al.*, 2011). The distinctive odour of nutmeg is attributed to the presence of essential oil-containing terpenes (α -pinene, p-cymene, sabinene, camphene, myrcene and γ -terpinene) terpene derivatives (terpinol, geraniol, and linalool) and phenylpropanes (myricetin, safrole, and elemicin) (Widelski *et al.*, 2017). Nutmeg is known for its various properties, including aphrodisiac, stomachic, carminative, tonic, nervous stimulant, aromatic, narcotic, astringent, hypolipidemic, antithrombotic, antifungal, antidysentric and anti-inflammatory effects (Francis *et al.*, 2014; Pashapoor *et al.*, 2019). Recent studies indicate that nutmeg can potentially mitigate damage caused by gamma radiation and improve memory in mice. Furthermore, nutmeg has been found to possess anti-inflammatory properties and exhibit insulin-like biological activity. In ancient Indian medicine, nutmeg is believed to have cardio tonic properties, leading to nutmeg aqueous extract (NMAET) for treating hepatotoxicity linked to ISO (isoproterenol)-induced oxidative stress. Further studies showed the antihyperglycemic and hyperlipidemia effects of NMAET (Kareem *et al.*, 2013).

In addition, *M. fragrans* exhibits significant antimicrobial activity against various bacterial pathogens (*Streptococcus mutans*, *S. sanguis*, *S. salivarius*, and *Lactococcus casei*) and antioxidant properties as it is rich in many bioactive compounds (Chung *et al.*, 2006; Matulyte *et al.*, 2020). Recently, the disc diffusion method was used to investigate the antimicrobial activity of nutmeg against Gram-

positive bacteria (*Staphylococcus aureus*, *Listeria monocytogenes*, *Bacillus cereus*) and Gram-negative bacteria (*Klebsiella pneumoniae*, *Escherichia coli*, *Proteus vulgaris*, *Pseudomonas aeruginosa*) (Nikolic *et al.*, 2021).

Besides this, it was also shown that terpene hydrocarbons (60-80% of the oil) are the main constituents of *M. fragrans* (pinene, sabinene, camphene, p-cymene, terpinene, phellandrene, myrcene, and limonene), followed by aromatic ethers (myristicin, elemicin, safrole, elemicin, and eugenol derivatives) representing 15% to 20% of the oil, and oxygenated terpenes (geraniol, linalool, and terpineol), which make up approximately 5% to 15% of the *M. fragrans* structure (Muchtaridi *et al.*, 2010; Rashidian *et al.*, 2022). Additionally, phytochemical studies of *M. fragrans* seeds have also shown that it is rich in alkaloids, glycosides, flavonoids tannins, saponins and steroids (Cruz *et al.*, 2024). So the resourcefulness of nutmeg makes it a significant ingredient in a variety of fields, including culinary arts, healthcare, and pharmaceuticals (Ramteke *et al.*, 2024).

The objective of this study was to investigate the phytochemical constituents and antimicrobial activity of *M. fragrans* against gram-negative (*Escherichia coli*) and gram-positive (*Staphylococcus aureus*) bacteria isolated from ATM (Automated Teller Machine) of Bikaner, Rajasthan, India.

MATERIAL AND METHODS

Collection and Preparation of *M. fragrans* Seeds Extract

Dried seeds of nutmeg collected from a local market in Bikaner, Rajasthan were washed and the extract was prepared according to Ozaki, *et al* procedure with slight modification (Ozaki *et al.*, 1989). To extract bioactive compounds from the seeds of nutmeg, 14 gms of the crude powder of seeds was refluxed with 1000 ml of water and 70% ethanol, respectively in the soxhlet apparatus for 8 hours. To eliminate residual solids the solution was filtered through a filter paper (Whatman GradeNo.18) and evaporated to dryness under vacuum at 40⁰ C. Subsequently, the dried extract was weighed and the stock solution was prepared by dissolving one gram of dried extract in 10 ml of 70% ethanol and water respectively.

Determination of Moisture Content and Solubility

The moisture content was determined using the procedure given by B.C. O'Kelly (Kelly *et al.*, 2016) using the following formula.

$$\text{Moisture content (\%)} = \frac{W_1 - W_2}{\text{Weight of sample}} \times 100$$

The solubility was determined using the method given by Elena, 2012.

Phytochemical Analysis

The nutmeg extracts were screened for the presence of alkaloids, tannins, phenols and saponins using the standard procedures.

Alkaloid determination

Five grams of the sample was weighed into a 250ml beaker followed by the addition of 200ml of 10% acetic acid in ethanol. The mixture was allowed to stand for 4 minutes before being filtered. The resulting extract was then concentrated on a water bath to one-quarter of the original volume. Precipitate was formed by adding concentrated ammonium hydroxide dropwise to the extract. The solution was then allowed to settle, and the precipitate was collected, washed with dilute ammonium hydroxide, and filtered. The residue obtained was alkaloid, which was subsequently dried and weighed (Harbone, 1973).

$$\% \text{Alkaloid} = \frac{W_3 - W_2}{W_1} \times 100\%$$

Where

W_1 = initial weight of sample

W_2 = weight of extract

W_3 = final weight of the residue

Test for Tannins

The presence of tannins was assessed using the procedure given by Tomar and Shrivastava, 2016. Specifically, 2 ml of leaf extract was combined with a few drops of 5% ferric chloride. The development of brown colour indicates the presence of tannins.

Test for Phenols

The presence of phenols was determined using the procedure given by Tomar and Shrivastava, 2016. Approximately 1 ml of extract was mixed with 2 ml of distilled water and a few drops of 10% of aqueous FeCl₃. The presence of phenol was indicated by the formation of blue or green colour.

Test for saponin

Extracts were spotted onto a layer of silica gel-activated glass plates. The plate's bottom was immersed in the solvent, and once the solvent reached the top, the plate was taken out and the solvent was evaporated. The solvent system utilized for separation was butanol: water: acetic acid (12:2:1). Following the application of a visualisation agent (iodine vapour), brown spots of saponin was identified in the chromatogram upon examination under UV light (Ginovyana *et al.*, 2020).

Furthermore, tests for saponins were conducted using the **Foam test method**. Approximately 1 ml of the extract was diluted with 20 ml of distilled water and vigorously shaken in a test tube. The presence of foam formation in the upper part of the test tube indicates the presence of saponins (Mendhekar *et al.*, 2017).

Antimicrobial Activity of *M. fragrans*

The agar well diffusion method was used for the antimicrobial activity of nutmeg extract using the method given by Ibrahim *et al.* (Ibrahim *et al.*, 2013) using Gram-negative (*E. coli*) and Gram-positive (*S. aureus*) bacteria. The microbes used in this study were previously isolated in the Department of Microbiology, MN College and Research Institute, Bikaner. The bacteria were grown in Muller Hilton agar for 18 hours at 37°C. Afterwards, a loop full of culture was added to 5 ml fresh Muller Hilton broth and adjusted to 0.5 McFarland by using a saline solution. A sterile swab was used for further seeding these bacteria on Muller Hilton agar. After drying the agar, a sterile cork borer was used to cut 4mm diameter wells. The wells were

then filled with 0.1 ml of nutmeg extract and allowed to diffuse for 2 hours at room temperature. After incubating plates at 37^o C for 24 hours, the diameters of inhibition zones (mm) were measured.

Statistical Analysis

The data collected from laboratory experiments was analyzed using descriptive statistical methods, specifically calculating simple means to determine the average value of the data points and percentages to represent the distribution of values within the dataset. This analysis helps to summarize and interpret the data clearly and concisely, providing valuable insights into the results of the experiments.

RESULTS

Moisture Content and Solubility of *M. fragrans* Seeds

Quantitative analysis of *M. fragrans* (nutmeg) extract showed that the moisture content (Loss on Drying) in powdered raw material of nutmeg was 95% and solubility of raw seeds in water was 40% and ethanol was 70%.

Phytochemical Compounds of *M. fragrans* Seeds Extract

The phytochemical analysis of water and 70% ethanolic extract of *M. fragrans* seeds were given in table 1. It was observed that ethanolic extract contains alkaloids, saponins, tannins and phenols while water extract only contains alkaloids and saponins but lacks tannins and phenols. Besides this ethanolic extract has a high percentage of alkaloids in comparison to water extract, which is 13.25 and 7.14% respectively (fig 1).

Antimicrobial activity

The results of the antimicrobial activity of different extracts against chosen microorganisms are shown in table 2. In the case of *Escherichia coli*, the maximum zone of inhibition was by 70% ethanol extract of nutmeg followed by extract in water and then raw material in ethanol. Raw material in water didn't show any activity against *E. coli*. Whereas, in the case of *staphylococcus aureus*, the maximum zone of inhibition was by 70% ethanol extract of nutmeg followed by extract in water, raw material in 70% ethanol and raw material in water (fig 2).

DISCUSSION

Medicinal plants have great significance due to the synthesis of a wide range of medicinally important chemical compounds. These compounds shield us from various pathogenic diseases (Pradhan and Huidrom, 2022). *Myristica fragrans* is one such plant that synthesizes a wide range of chemicals as described in the introduction part of this manuscript.

In the present study, phytochemical analysis was done specifically for alkaloids, phenols, saponins and tannins. Other researchers screened these components as they are the main components of medicinal plants having antimicrobial properties (Chandra, 2013). The results of the present study revealed that the ethanolic extract of *M. fragrans* contains alkaloids, saponins, Phenol and tannins while water extract only contains alkaloids and saponins but lacks tannins and phenols. Similar findings have also been reported by other workers (Gayathri and Anuradha, 2015). Various bioactive compounds found in the extract exhibit important biological properties. For instance, alkaloids demonstrate cytotoxicity and are utilized in allopathic systems (Tackie and Schiff, 1993). Steroidal compounds are particularly significant in pharmacy due to their association with compounds like sex hormones, commonly used in the production of female contraceptive pills (Akhtar *et al.*, 2017). Tannins inhibit the growth of fungi, yeast, bacteria, and viruses, acting as antioxidants comparable to phenols (Darout *et al.*, 2000). Phenols and flavonoids possess antioxidant, anti-allergic and antibacterial properties, while saponins exhibit anti-inflammatory, antiviral, and plant defence activities (Kumar, 2019). Henceforth, the presence of all these phyto-constituents contributes to the treatment of severe human infections (Olivia *et al.*, 2021).

The ethanolic extract of *M. fragrans* showed the highest effectivity against both gram-positive (*S. aureus*) and gram-negative (*E. coli*) bacteria while the water extract of nutmeg showed less activity for both bacteria. Similar findings were attained by other researchers (Mahady *et al.* in 2005; Shafiei *et al.*, 2012; Thanoon and Kamal, 2013). The essential oil of *M. fragrans* exhibits significant inhibitory effects against a variety of bacteria, including bacteria that cause food poisoning, plant and animal pathogens, and spoilage (Jose *et al.*, 2024). The essential oil of *M. fragrans* has the potential to treat several hazardous diseases (Algaffar *et al.*, 2024). The present study revealed the antimicrobial activity of *M. fragrans* against *S. aureus* (Al-Qahtani *et al.*, 2022) and *E. coli* (Romeiro *et al.*, 2024). Hence, the results

of our study also strengthened the previous studies that the antimicrobial potential of the extracts may be due to the presence of rich phytochemical compositions.

CONCLUSION

This study evaluated the antimicrobial activity of *M. fragrans* extracts against selected microorganisms isolated from local ATMs, demonstrating a wide range of antimicrobial sensitivity. The bioactive compounds extracted from *M. fragrans* using water and 70% ethanol as solvents may account for these results. The findings suggest that this spice holds promise for new formulation development. However, further investigation is required to elucidate the precise mechanism underlying the antimicrobial activity of the extract, potentially enhancing its pharmacological applications. Based on the current results, *M. fragrans* shows potential as a natural antimicrobial agent against infections caused by *E. coli* and *S. aureus* of human origin. Further research is needed to identify the principle active component and the mechanism underlying the antimicrobial activity of the seed extracts of *M. fragrans*.

CONSENT

Not applicable

ETHICAL APPROVAL

Not applicable

REFERENCES

1. Abd Algaffar SO, Satyal P, Ashmawy NS, Verbon A, van de Sande WW and Khalid SA. In Vitro and In Vivo Wide-Spectrum Dual Antimycetomal Activity of Eight Essential Oils Coupled with Chemical Composition and Metabolomic Profiling. *Microbiology Research*. 2024; 15(3):1280-1297.
2. Akhtar N, Ali SS, Ahmed S, Jan S, Khan MA and Khan MS. Phytochemical Analysis of *Lepidium didymum*. *Pakistan Journal of Weed Science Research*. 2017; 23(2).

3. Al-Qahtani WH, Dinakarkumar Y, Arokiyaraj S, Saravanakumar V, Rajabathar JR, Arjun K and Appaturi JN. Phyto-chemical and biological activity of *Myristica fragrans*, an ayurvedic medicinal plant in Southern India and its ingredient analysis. *Saudi Journal of Biological Sciences*. 2022; 29(5):3815-3821.
4. Al-Rawi SS, Ibrahim AH, Ab Rahman NNN, Nama MMB, Majid AMA and Ab Kadir MO. The effect of supercritical fluid extraction parameters on the nutmeg oil extraction and its cytotoxic and antiangiogenic properties. *Procedia Food Science*. 2011; 1:1946-1952.
5. Al-Rawi SS, Ibrahim AH, Ahmed HJ and Khudhur ZO. Therapeutic, and pharmacological prospects of nutmeg seed: A comprehensive review for novel drug potential insights. *Saudi Pharmaceutical Journal*. 2024; 102067.
6. Chandra M. Antimicrobial activity of medicinal plants against human pathogenic bacteria. *International Journal of Biotechnology and Bioengineering Research*. 2013; 4 (7):653 - 658.
7. Christophersen C, Larsen C and Dimayuga RE. Traditional Medicine-A potential resource exploitation of natural products. The H.C. OrstedInstitute,Copenhagen, Denmark. 1991; 8-12.
8. Chung JY, Choo JH, Lee MH, Hwang JK. Anticariogenic Activity of Macelignan Isolated from *Myristica Fragrans* (Nutmeg) against *Streptococcus Mutans*. *Phytomedicine*. 2006; 13:261–266.
9. Cruz A, Sánchez-Hernández E, Teixeira A, Oliveira R, Cunha A and Martín-Ramos P. Phytoconstituents and Ergosterol Biosynthesis-Targeting Antimicrobial Activity of Nutmeg (*Myristica fragrans* Houtt.) against Phytopathogens. *Molecules*. 2024; 29(2):471.
10. Dahham SS, Al-Rawi SS, Ibrahim AH, Majid ASA, Majid AMSA. Antioxidant, anticancer, apoptosis properties and chemical composition of black truffle *Terfezia clavaryi*. *Saudi J. Biol. Sci*. 2018; 25 (8):1524–1534.
11. Darout IA, Christy AA, Skaug N, Egeberg PK. Identification and quantification of some potentially antimicrobial anionic components in Miswak extract. *Indian Journal of Pharmacology*. 2000; 32(1):11- 14.
12. Elena DL. Pharmacognostic methods for analysis of herbal drugs, According to European Pharmacopoeia. *Promising Pharmaceuticals*. 2012; 38-62.

13. Farnsworth NR and Buigel AS. Problems and Prospects of Discovering New Drugs from Higher Plants by Pharmacological Screening. Springer Verlag, Berlin, Germany. 1977.
14. Francis, KS, Suresh, E and Nair MS. Chemical constituents from *Myristica fragrans* fruit. Natural Product Research. 2014; 28(20):1664-8.
15. Gayathri R and Anuradha V. Phytochemical screening and total phenolic content of aqueous and acetone extracts of seed, butter, mace of nutmeg (*Myristica fragrans* Houtt). Int J Pharm Sci Rev Res. 2015; 33(1): 236-239.
16. Ginovyan M, Ayvazyan A, Nikoyan A, Tumanyan L and Trchounian A. Phytochemical screening and detection of antibacterial components from crude extracts of some Armenian herbs using TLC-bioautographic technique. Current Microbiology. 2020; 77:1223-1232.
17. Hamburger M and Hostettmann K. Bioactivity in plants. The link between phytochemistry and medicine. Phytochemistry. 1991; 30(12): 3864-3874.
18. Harbone JB. Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis. Chapman and Hall, London. 1973; 267-270.
19. Ibrahim KM, Naem RK, Abd-Sahib AS. Antibacterial activity of nutmeg (*Myristica fragrans*) seed extracts against some pathogenic bacteria. J Al-Nahrain Univ. 2013; 16(2):188-92.
20. Ibrahim TA, Opawale BO and Oyinloye JMA. 1. Antibacterial Activity of Herbal Extracts Against Multi Drug Resistant Strains of Bacteria from Clinical Origin by 1 TA Ibrahim, 2 BO Opawale and 3 JMA Oyinloye. Life sciences leaflets 15. 2011; 490-to.
21. Jose JS, Raj H and Thomas TA. *Myristica Fragrans*: A Potent Antibacterial Agent against Food Borne Pathogens. Asian Journal of Biological and Life Sciences. 2024; 13(1):133.
22. Kareem MA, Gadhamsetty SK, Shaik AH, Prasad EM and Kodidhela LD. Protective effect of nutmeg aqueous extract against experimentally-induced hepatotoxicity and oxidative stress in rats. Journal of Ayurveda and integrative medicine. 2013; 4(4):216.
23. Kelly B. C. O'. Oven-drying characteristics of soils of different origins. Dry. Technol. 2005; 23(5):1141–1149, 2005.
24. Kumar SA. Natural phytochemicals: potential anti - HCV targets in silico approach. Journal of Applied Pharmaceutical Science. 2019; (8):94 – 100

25. Mahady GB, Pendland SL, Stoia A, Hamill FA, Fabricant D, Dietz BM, Chadwick LR. In Vitro susceptibility of *Helicobacter pylori* to botanical extracts used traditionally for the treatment of gastrointestinal disorders. *Phytother. Res.* 2005; 19:988–991.
26. Matulyte I, Jekabsone A, Jankauskaite L, Zavistanaviciute P, Sakiene V, Bartkiene E, Ruzauskas M, Kopustinskiene DM, Santini A, Bernatoniene J. The Essential Oil and Hydrolats from *Myristica Fragrans* Seeds with Magnesium Aluminometasilicate as Excipient: Antioxidant, Antibacterial, and Anti-Inflammatory Activity. *Foods.* 2020; 9:37.
27. Mendhekar SY, Padekar SG, Date SH, Jadhav SL and Gaikwad DD. Pharmacognostic, Phytochemical, Physicochemical And Detail Microscopical Evaluation Of *Capparis Zeylanica* Linn. Leaves. 2017.
28. Mughtaridi SA, Apriyantono A, Mustarichie R. Identification of compounds in the essential oil of nutmeg seeds (*Myristica fragrans* Houtt.) that inhibit locomotor activity in mice, *Int J Mol Sci.* 2010; 1111:4771– 81.
29. Nikolic V, Nikolic L, Dinic A, Gajic I, Urosevic M, Stanojevic L, Stanojevic J, Danilovic B. Chemical Composition, Antioxidant and Antimicrobial Activity of Nutmeg (*Myristica Fragrans* Houtt.) Seed Essential Oil. *J. Essent. Oil Bear. Plants.* 2021; 24:218–227.
30. Olivia NU, Goodness UC and Obinna OM. Phytochemical profiling and GC - MS analysis of aqueous methanol fraction of *Hibiscus asper* leaves. *Futur J Pharm Sci.* 2021; 7:59. <https://doi.org/10.1186/s43094-021-00208-4>.
31. Orwa C, Mutua A, Kindt R, Jamnadass R and Simons A. Agroforestree Database: a tree reference and selection guide. *Agroforestree Database: a tree reference and selection guide.* 2009; 4.
32. Ozaki Y, Sekita S, Soedigdo S and Harada M. Antiinflammatory effect of *Graptophyllum pictum* (L.) Griff. *Chemical and pharmaceutical bulletin.* 1989; 37(10):2799-2802.
33. Pashapoor A, Mashhadrafie S and Mortazavi P. Ameliorative effect of *Myristica fragrans* (nutmeg) extract on oxidative status and histology of pancreas in alloxan induced diabetic rats. *Folia morphologica.* 2020; 79(1):113-119.

34. Pradhan S and Huidrom P. Antimicrobial Properties and Phytochemical Analysis of Some Medicinal Plants: A Review. *International Journal of Science and Research (IJSR)*. 2022; 508-511.
35. Ramteke AM, Gourand PB and Malode SM. Role of isolated compounds from spices and herbs in natural medicinal care or therapy-A review. In *AIP Conference Proceedings*. 2024; 2974(1).
36. Rashidian G, Shahin K, Elshopakey GE, Mahboub HH, Fahim A, Elabd H and Faggio C. The dietary effects of nutmeg (*Myristica fragrans*) extract on growth, hematological parameters, immunity, antioxidant status, and disease resistance of common carp (*Cyprinus carpio*) against *Aeromonas hydrophila*. *Journal of Marine Science and Engineering*. 2022; 10(3):325.
37. Romeiro PFR, Coelho AS, de Almeida JM, Alonso VPP, Prata AS, Crippa BL and Silva NCC. Unraveling the antimicrobial activity of nutmeg and turmeric essential oils against *Staphylococcus aureus*, *Listeria monocytogenes*, *Escherichia coli*, and *Salmonella spp.* *Food Science and Technology*. 2024; 44.
38. Shafiei Z, Shuhairi NN, Md Fazly Shah Yap N, Harry Sibungkil CA, Latip J. Antibacterial activity of *Myristica fragrans* against oral pathogens. *Evid.-Based Complement. Altern. Med.* 2012; 825362.
39. Tackie AN, Schiff PL. Cryptospirolepine, a unique spiro-noncyclic alkaloid isolated from *Cryptolepis sanguinolenta*. *Journal of Natural Products*. 1993; 56:653-5.
40. Thanoon SG and Al-Refai Kamal A. Antibacterial effect and healing potential of nutmegoil for chemically induced oral ulcerations in rabbits. *Zanco Journal of Medical Sciences*. 2013; 17(2): 393-399.
41. Tomar RS and Shrivastava V. Phytochemical profiling and efficacy evaluation of antibacterial potential of *Myristica fragrans* seed extracts against selected microorganisms. *Asian J Pharm Clin Res*. 2016; 9(6):1-3.
42. Widelski J and Kukula-Koch WA. Psychoactive drugs. In *Pharmacognosy*. 2017; 363-374).
43. Wyk BV and Gericke N. *People's plants: A guide to useful plants of Southern Africa*. 2000; 351.

Table-1: The Result of Phytochemical Analysis of *MyristicaFragrans* Seed Extract

Phytochemicals	70% ethanolic extract of <i>M. fragrans</i> seed	Water extract of <i>M. fragrans</i> seed
Alkaloid	+(13.25%)	+(7.14%)
Tannin	+	-
Saponin	+	+
Phenol	+	-

Table-2: Antimicrobial analysis of *M. Fragrans* raw material and extract in mm.

S.no	Bacteria	Effectivity of raw material in water	Effectivity of raw material in 70% ethanol	Effectivity of extract in water	Effectivity of extract in 70% ethanol
1	<i>Escherichia coli</i>	nil	6	9	11
2	<i>Staphylococcus aureus</i>	6	7	10	11




Figure-1: Showing percentage of Alkaloid in *M. fragrans*(seed)raw material, extract in water and extract in 70% ethanol




Figure-2: Antimicrobial analysis of raw material and extracts of *M. fragrans*