

## Original Research Paper

### A Comparative Evaluation of the Antisickling, Elemental and Proximate Analysis of the Leaves of Three *Khaya* species found in Nigeria

#### ABSTRACT

Sickle cell disease (SCD) is traditionally treated in Nigeria by the Traditional Medical Practitioners (TMPs) with medicinal plants and other plant products. One of the herbs commonly used singly or in combination with other plant materials is *Khaya*, which is widely distributed in Africa, with three species existing in Nigeria. These are *Khaya senegalensis* A. Juss (KS), *Khaya grandifoliola* (WELW) CDC (KG) and *Khaya ivorensis* A. Chev. (KI). In this study we evaluated the anti-sickling activities of the leaf extracts of the three *Khaya* species leaves with a view to authenticating the ethnomedicinal claims, and assess the proximate as well as the elemental constituents that may contribute to the properties.

Dried leaves of the *Khaya* species were extracted by cold maceration and Soxhlet methods using water, ethanol and petroleum spirit as extractives. Filtered extracts were concentrated *in vacuo* at 40°C and dried in a lyophilizer. Inhibitory and reversal models were employed to evaluate their antisickling effects on Hb SS red blood cells following standard procedures. Proximate and elemental analysis were determined using the simple analytical procedure as described by the Association of Official Analytical Chemistry (AOAC) of 1987, 13<sup>th</sup> Edition Washington DC P1094 and Dry Ashing method of AOAC 2006 respectfully.

Anti-sickling results showed that the Soxhlet ethanol extracts of the *Khaya* species demonstrated significant ( $p \leq 0.05$ ) ability to reverse and inhibit the sickling of red blood cells *in-vitro* with KI having the highest inhibitory and reversal of 60.04±1.77% and 74.97±2.23% respectively at 4 mg/ml concentration. The proximate analysis revealed the richness of the *Khaya* leaves in

minerals, proteins and carbohydrates. *Khaya* leaves contained calcium, magnesium, sodium, phosphorus, and other essential elements within the limits of dietary requirement.

*Khaya* species could therefore be recommended for further development into herbal remedy in the management of sickle cell disease.

**Keywords:** *Khaya sp*, Red Blood Cells, Sickle Cell, Anti-sickling, Physicochemical.

## 1.0 INTRODUCTION

Sickle cell diseases (SCD) are a group of disorders that have in common the propensity of the red blood cells to become deformed when oxygen in the blood is lowered, causing occlusion of blood vessels by misshapen cells, anemia, and various associated clinical consequences, including death [1]. Sickle cell disease is caused by the substitution of glutamic acid with valine at the sixth position of the beta-globin chain of hemoglobin S (HbS). This mutation causes the hemoglobin to form abnormal or long and rod-like shape within the red blood cell, because the mutated genes have low solubility in the bloodstream. Under hypoxic conditions, deoxy-HbS molecules polymerise, forming rigid, sickled cells, which in turn causes the deformation of the normal disc biconcave red blood cells [2]. SCD is a hereditary disease passed down through families. Inheriting one gene for hemoglobin S, together with a normal gene, results in the formation of red cells that contain approximately 40 per cent of the abnormal hemoglobin and 60 per cent of the normal hemoglobin, an essentially harmless state that is designated as sickle cell trait [3]. However, if the gene inherited together with the sickle gene is not normal, then the sickle cell disease may develop. The most common hemoglobin that interacts with sickle hemoglobin is hemoglobin C, and the  $\beta$ -thalassemia (beta-thalassemia) mutation also interacts with the sickle gene by restricting the formation of normal hemoglobin. Sickle gene, and genes that interact with it, are common in a number of different populations, but the highest gene

frequencies are observed in Africa [4]. The gene is also found in Southern Europe, the Middle East, and India. Among African Americans, approximately 7.8 per cent are carriers of the sickle mutation, while 2.3 per cent have hemoglobin C trait and 0.8 per cent have  $\beta$ -thalassemia trait [4].

Signs and symptoms of sickle cell anemia usually commence after an infant is 4 months old and may include anemia which is a chronic shortage of red blood cells [5], [6]. SCD is often accompanied with crisis such as Vaso-occlusive crisis, splenic sequestration crisis, aplastic, hemolytic crisis and death [7]. Episodes of pain are one of the major symptoms of sickle cell anemia which develops when sickle-shaped red blood cells block blood flow through tiny blood vessels to the chest, abdomen and joints [7]. These severe pains have been claiming life of children, youths and adults across the world especially among the African population.

After centuries of research by both orthodox and trado-medical researchers, there has been no other cure either traditionally or scientifically aside bone marrow transplantation (BMT). Current therapeutic options remain limited to hydroxyurea, although more recently, L-Glutamine (Endari), Voxelotor (Oxbryta) and Crizanlizumab (Adakyeo) were approved by the US Food and Drug Administration (FDA) [8, 9, 10], however, the drugs are not available to the low-income countries where SCD is prevalent. Hence, researches to discover more therapeutic options are still ongoing.

Scientists have been carrying out research into ethno-medicine to confirm the potency of medicinal plants used by the traditional medical practitioners in the treatment and management of various diseases including SCD, standardizing and formulating them into useful dosage forms. One of such medicinal plants used to manage sickle cell individuals in Southern Nigeria is *Khaya* Species commonly referred to as Oganwo by the Yoruba speaking people of Nigeria, and

the trade name is African Mahogany. *Khaya* belongs to family Meliaceae, native to tropical Africa and found along the Western Africa sub- region. In spite of the plants being morphologically different they are all commonly called African mahogany and are used to treat different types of diseases. Ethno-medicinally, the different parts of *Khaya* are employed in the treatment of headaches, convulsion, cough, whooping cough, stomach ache, fever, threatened abortion, as lotion for rheumatism, dermatomycosis, and malaria fever in Nigeria. When mixed with black pepper it is used to treat diarrhea and dysentery; bark decoction is used as a laxative, as an aphrodisiac, and used for treating back pains, syphilis, jaundice, and mental illness [11]. Although Elenga *et al.*, [12] reported the anti-sickling activity of *K. senegalensis* stem bark, we designed this research to determine and compare the anti-sickling properties of the leaf extracts of the three species (*Khaya ivorensis* A.Chev., *K. grandifoliola* C.Dc. and *K. senegalensis* A. Juss.) found in Nigeria, evaluate the proximate and elemental constituents with a view to confirm the ethnomedicinal use and quality of the plant drug components.

## 2.0 MATERIALS AND METHODS

### 2.1 Plants Collection and Processing

Fresh leaves of *Khaya senegalensis* (KS), *Khaya grandifoliola* (KG) and *Khaya ivorensis*(KI) (Fig. 1) were individually collected between the month of December and January, with the geographical and altitudinal coordinates of the location recorded using the GPS device. *K. senegalensis* was collected from the Botanical Garden (7°31'20.42"N, 4°31'20.93"E) of Obafemi Awolowo University, Ile-Ife, Osun State; *K. grandifoliola* from Abata Egba village (7°19'11.18"N, 4°36'04.89"E), Osun State and *K. ivorensis* A from Oke Ado (Okomu Forest Reserves) (6°20'00"N, 5°16'00"E), in Ovia Local Government, Okada, Edo State, Nigeria. The plants were identified and authenticated by the curator at the IFE herbarium located in the

Department of Botany, Obafemi Awolowo University, Ile-Ife and the voucher specimens deposited and assigned numbers as follows: 16289 IFE (KS), 16348 IFE (KG) and 16343 IFE (KI).

The leaves were first air dried at room temperature and afterwards activated in the oven at 40°C before grinding into powder using Christy machine. The powdered leaves were stored in well-sealed amber colored bottles until ready for use. All experiments were carried out at Drug Research and Production Unit, Faculty of Pharmacy, Obafemi Awolowo University, Ile-Ife, Nigeria.



**Figure 1: Pictures of the three *Khaya* Spp found in Nigeria. (A) *Khaya senegalensis*, (B) *Khaya grandifoliola*, and (C) *Khaya ivorensis***

## **2.2 Extraction Procedures**


### **2.2.1 Cold Extraction**

Powdered plant materials 250 g each of KS, KG and KI were macerated separately in water, ethanol and petroleum spirit for 72 h with constant shaking using electrically operated

mechanical shaker, to ensure exhaustive extraction. The extracts were filtered and concentrated *in vacuo* at 40°C or freeze dried to achieve complete dryness.

### 2.2.2 Hot Extraction (Soxhlet)

Powdered plant materials 250 g each of KS, KG and KI were exhaustively extracted, using a Soxhlet extractor with water, ethanol and petroleum spirit separately. The extracts were filtered, concentrated *in vacuo* at 40°C and then lyophilized to achieve complete dryness.



**Figure 2: Flow Diagram of Plant Preparation and Extraction**

## 2.3 Anti-sickling Assay

### 2.3.1 Collection of Blood Samples

Whole blood sample (5 ml) from sickle cell anemia patients in steady state between the ages of 12 and 23 years collected into an EDTA (Ethylene Diamino Tetra-acetic Acid) bottle, by vein puncture was used. This exercise was carried out by the hematologists from confirmed sickle cell patients attending the regular hematology out-patient clinic at Obafemi Awolowo University Teaching Hospital Complex (OAUTHC) after securing Ethical Clearance from the OAUTHC Ethical Committee.

### 2.3.2 Inhibitory and Reversal Anti-sickling Test

The anti-sickling assays were carried out using modified methods of Egunjobi *et al.*, [13]. Vanillic acid solution, 4 mg/ml, was used in place of the extract as positive control while 0.2 ml of phosphate buffered saline (pH 7.0) was used as negative control. All experiments were carried out in triplicates. Using a light microscope, sickled and un-sickled cells were counted to calculate the percentage inhibitory or reversal activity [14].

## 2.4 Proximate Analysis and Elemental Analysis Procedures

### 2.4.1 Proximate Analysis

Simple analytical procedure for proximate analysis was used as described by the Official methods of Analysis of the Association of Official Analytical Chemistry [15], 13<sup>th</sup> edition Washington D.C. P1094. United States of America and Official methods of Analysis of the Association of Official Analytical Chemistry [16], revised edition Washington D.C. United States of America. The Moisture Content, Total Ash, Total Crude Fibre, Ether Extract (% Fat, % Oil or % Lipid) and Total Crude Protein were determined according to AOAC, 2006 methods.

### 2.4.2 Elemental Analysis Procedure

#### *Determination of digestion for the determination of Minerals (Dry Ashing Method).*

Samples of powdered *Khaya* leaves (2g) were placed in a ceramic crucible, heated in a muffle furnace of hot air oven at 600<sup>o</sup>C for 3 h and thereafter allowed to cool in a desiccator. 5 ml of 6M HCl was added into each container and allowed to stand for 30 mins for proper digestion. After digestion the contents were filtered with Whatman No 1 filter paper into a 50 ml conical flask. The filtrate was made up to mark (50 ml) with distilled water, shaken vigorously and the elemental content of each of the three species were read with the aid of Atomic Absorption Spectrophotometer (AAS) [16].

## 2.5 Statistical Analysis


All experiments were carried out in triplicate and result presented as mean  $\pm$  SEM. One-way ANOVA was used to detect significant differences and the mean significant value was set at  $p \leq 0.05$ .

## 3.0 RESULTS

### 3.1 Anti-Sickling Activities


The antisickling results showed concentration dependent inhibitory and reversal properties. The Soxhlet extracts of the three extraction solvents, water, ethanol and petroleum spirit, exhibited higher anti-sickling properties than the extracts of cold maceration. For the cold extracts, ethanol extracts of KI gave the highest inhibitory and reversal activities (Figures 3 and 4). Constituents of the petroleum spirit extracts are non-polar phytochemicals and not active in preventing nor reversing sickling of red blood cells.

For the Soxhlet extracts, ethanol extract of KI gave the highest inhibition of  $60.04 \pm 1.77\%$  followed by KG ( $49.71 \pm 1.14\%$ ) and KS with the least activity ( $44.88 \pm 1.95\%$ ) at 4 mg/ml concentration (Figure 5). The same trend was observed with the reversal activity. At 4 mg/ml concentration, KI reversed sickled Hb SS red blood cells ( $74.97 \pm 2.23\%$ ) followed by KG ( $59.96 \pm 3.10\%$ ) and lastly KS with  $51.62 \pm 1.83\%$  reversal activity (Figure 6). The study generally revealed that reversal activities were concentration dependent as activities increased with concentration.



**Figure 3: Inhibitory properties of the macerated aqueous, ethanol and petroleum spirit extracts of KS, KG and KI at various concentrations.**


**Key: KS: *Khaya senegalensis*; KG: *Khaya grandifoliola*; KI: *Khaya ivorensis***



**Figure 4: Reversal properties of the macerated aqueous, ethanol and petroleum spirit extracts of KS, KG and KI at various concentrations.**

**Key: KS: *Khaya senegalensis*; KG: *Khaya grandifoliola*; KI: *Khaya ivorensis***


UNL



**Figure 5: Inhibitory properties of the aqueous, ethanol and petroleum spirit Soxhlet extracts of KS, KG and KI at various concentrations.**

**Key: KS: *Khaya senegalensis*; KG: *Khaya grandifoliola*; KI: *Khaya ivorensis***

UNL



**Figure 6: Reversal properties of the aqueous, ethanol and petroleum spirit Soxhlet extracts of KS, KG and KI at various concentrations.**

**Key: KS: *Khaya senegalensis*; KG: *Khaya grandifoliola*; KI: *Khaya ivorensis***

### 3.2 Proximate Composition

The proximate and elemental analysis of KS, KG and KI are presented in tables 1 and 2. The leaves of the three plants are rich in carbohydrates and crude protein, and contain trace levels of nickel, cadmium, and lead.

**Table 1: Proximate Composition of the three *Khaya* species leaves.**

Sample Description	% Moisture	% Total Ash	% Crude Fibre	% Ether Extract	% Crude Protein	% Nitrogen Free Extract (NFE)	% Total Carbohydrate
KS	10.55	10.09	15.26	3.18	23.63	37.29	52.55
KG	10.35	7.72	24.89	3.02	20.56	33.46	58.35
KI	9.32	7.22	17.29	6.71	18.38	41.08	58.37

**Key: KS: *Khaya senegalensis*; KG: *Khaya grandifoliola*; KI: *Khaya ivorensis***

**Table 2: Elemental Analysis of the three *Khaya* species leaves**

Extracts	Cu (Copper)	Ni (Nickel)	Fe (Iron)	Zn (Zinc)	Mn (Manganese)	K (Potassium)	Cd (Cadmium)	Na (Sodium)	Cr (Chromium)	Ca (Calcium)	Mg (Magnesium)	Pb (Lead)
<b>KS</b>	0.237 ± 0.004	0.160 ± 0.007	12.68 ± 0.008	0.842 ± 0.002	1.005 ± 0.008	10.27 ± 0.091	0.033 ± 0.001	0.036 ± 0.001	-	8.384 ± 0.069	0.714 ± 0.002	0.108 ± 0.013
<b>KG</b>	0.991 ± 0.001	0.038 ± 0.007	9.139 ± 0.016	1.016 ± 0.005	1.267 ± 0.002	10.16 ± 0.312	0.022 ± 0.001	0.323 ± 0.002	0.416 ± 0.041	4.322 ± 0.019	0.824 ± 0.002	0.053 ± 0.007
<b>KI</b>	0.440 ± 0.001	0.063 ± 0.002	11.34 ± 0.020	0.713 ± 0.002	0.943 ± 0.002	9.051 ± 0.134	0.018 ± 0.001	0.059 ± 0.001	0.491 ± 0.010	5.422 ± 0.032	0.832 ± 0.000	0.045 ± 0.027

**Key: KS: *Khaya senegalensis*; KG: *Khaya grandifoliola*; KI: *Khaya ivorensis***

#### 4.0 Discussion

The inhibitory activities exhibited by the aqueous and ethanol extracts of the *Khaya* species at 4 mg/ml were significantly comparable to that of Vanillic acid ( $58.20 \pm 0.92$  %) while the reversal activities were significantly higher (Figures1-4).

The antisickling activities of KS, KG and KI were dose dependent as the highest activities were recorded at 4 mg/ml (Figures 1-4). This was observed across the three-plant species regardless of their mode and solvent of extraction. We employed water, ethanol and petroleum spirit as the extraction solvents to determine the class of putative compounds responsible for the activities, and also used cold and hot extraction methods to know whether active compounds are thermolabile or not.

Across all the tested plant extracts, the ethanol extracts gave the highest inhibitory properties followed by the aqueous and the non-polar petroleum spirit extracts. Although flammable, alcohol is recognized as non-toxic and has fewer handling risks [17]. Due to the organic nature of ethanol, it has a better extractive capability of plant metabolites, the putative compounds, which are often organic in nature. The putative active compounds responsible for inhibiting and reversing the sickling of red blood cells in the *Khaya* species were polar compounds. Compounds such as amino acids, flavonoids and phenols which have been reported to display antisickling properties were reported to be present in KG [18, 19]. These compounds were also not thermolabile as their activities were not hindered by the exertion of heat during extraction.

The general trend of the anti-sickling activities of the three plants was  $KI > KG > KS$  in all the tested concentrations, irrespective of hot or cold extraction methods.

The percentage moisture contents (MC) of KS, KG and KI were at acceptable levels as specified by the British pharmacopoeia which stated that the general requirement of MC in crude drug should not be more than 14% [20] and the value obtained in this study is within the acceptable range of 9.32%-10.55% (Table 1) implying a long storage period with less chances of microbial degradation of the drugs [21].

The physicochemical data would help in determining the richness and usefulness of the *Khaya* species in combating SCD and for the identification of the drug from its substitute or adulterants. The total ash contents of *Khaya* leaves ranged between 7.22%-10.09% (Table 2). In all, the highest % ash content was found in KS leaf 10.09%. The high ash content in *Khaya* species showed that it contained high mineral contents and may contain inorganic radicals like carbonates, phosphates, and silicates of sodium, potassium, calcium, magnesium [22] which may be the reason for their effect in inhibiting and reversal of sickled red blood cells (Table 1). Values are comparable to other leafy vegetables reported such as *Amaranthus hybridus*, and *Curcubita pepo* [23]. The quantitative evaluation is an important parameter in setting standard of crude drugs and the physical constant parameters could be useful in detecting any adulterant in the drug. The crude fibers of the *Khaya* species studied were within 15.26-24.89%, which is within the allowed values of 8.50 - 20.90% for some Nigerian vegetables [24]. Crude fiber analysis is an effective tool to determine the nutritional value of animal feed and some plant-based foods. The effects of dietary fiber on health outcomes include improving digestion, reducing cholesterol levels, weight management, and preventing chronic illnesses like heart disease and diabetes [25, 26]. The crude protein content ranged from 18.38 - 23.63%. KS leaves have the highest with 23.63%, KG with 20.56%, and KI with 18.38% protein content. The result showed that the crude proteins were generally high in the leaves of the three species studied (Table 1). The crude protein content is higher than what was reported for some other green leafy vegetables such as *Momordica balsamina* (11.29%), *Moringa oleifera* (20.72%) and *Leptadenia hastane* (19.10%) [27, 28, 29]. Plant foods that provide more than 12% of their calorific values from protein has been shown to be good sources of protein [30]. Besides this, the high protein content will help in replacing worn out cell/tissues of SCD patients thereby supporting their well-

being and growth. Although the crude protein content of KS was the highest it did not correlate to a higher anti-sickling property.

Qualitative and quantitative elemental analysis of KI, KG and KS are presented in Table 2. Calcium, iron, manganese, zinc and copper are of great importance and essence to our daily body due to their physiological and biological roles and functions [31]. Manganese (Mn) is a vital mineral needed for normal growth and skeletal formation. It helps to break down fats, carbohydrates and proteins, and serves as co-factor for enzymes [32]. Mn deficiency causes diabetes, nervous instability, disorder of bony cartilaginous growths in infant children and rheumatic arthritis in adults [33]; and since rheumatic arthritis is a symptom in SCD patient then Mn supplement in medicinal plants like *Khaya* can alleviate this. Zinc is an intercellular cation present in all body tissues and fluids next to Iron. It is the second abundant trace elements in humans. It is important for enzymatic functions. It takes part in synthesis of DNA, protein, insulin. It is also essential for normal functioning of the cells including protein synthesis, carbohydrates metabolism, cell growth and cell division [31, 32]. *Khaya* leaves contain significant amount of zinc implying its importance in replacement of worn-out tissues which is important to SCD patients. *Khaya* species can supply substantial portion of Mn, Zn and Cu requirement of adults in the tropics who cannot afford other sources of these minerals such as milk, cheese and eggs. Since SCD has effect on blood content and quality, the use of *Khaya* will go a long way in fighting the anemic state frequently presented by SCD patients in crisis. Calcium is another important mineral present in the leaves of the three *Khaya* species. Patients with SCD are always also presented with bone problems therefore calcium in the leaves will help in formation and maintenance of strong bones and teeth if developed into remedy. Sodium and potassium are also important and are present in the leaves of these species; they have important

intracellular and intercellular activities respectively. Sodium is involved in the regulation of plasma volume, acid-base balance, nerve and muscle contraction [34]. Heavy metals present in the leaves of *Khaya* species are cadmium, nickel, lead and chromium within permissible limit (Table 2) and implies the three *Khaya* are safe for consumption if developed into herbal remedy for management of SCD.

## 5.0 CONCLUSION

The three species of *Khaya* leaves showed a concentration and thermal dependent anti-sickling activities. The proximate analysis also reveals that *Khaya* leaves are very rich in nutrients like minerals, protein, carbohydrates in amounts required by the body and a moisture content that will allow storage if eventually developed into herbal remedy or drugs. In addition, the elemental analysis also revealed that the leaves of the three *Khaya* species studied supports healthy living because they contain elements within permissible limit for the body.

### **Ethical Approval:**

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

### **Consent**

As per international standards or university standards, patient(s) written consent and parental written consent has been collected and preserved by the author(s).

## Disclaimer (Artificial intelligence)

### Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during writing or editing of manuscripts.

### Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

## 6.0 REFERENCES

1. Platt OS, Brambilla DJ, Rosse WF. Mortality in sickle cell disease. Life expectancy and risk factors for early death. *N. Engl. J. Med.* 1994; 330 (23):1639–44.
2. Patric A, Daging H, Ingrid H, Denisa D. Pro-coagulant state resulting from high levels of soluble P-selectin in blood. *PNAS* 2000; 97 (25):13835–40.
3. Roach ES. "Sickle Cell Trait". *Archives of Neurology.* 2005; 62 (11):1781-2.
4. Wellems TE, Hayton K, Fairhurst RM. The impact of malaria parasitism: from corpuscles to communities. *J. Clin. Invest.* 2009; 119 (9): 2496–505.
5. Chrouser KL, Ajiboye OB, Oyetunji TA, Chang DC. "Priapism in the United States: the changing role of sickle cell disease". *Am. J. Surg.* 2011; 201(4):468–74.

6. Weatherall DJ, Clegg JB. Inherited haemoglobin disorders: an increasing global health problem. *Bull. World Health Organ.* 2001; 79 (8): 704–12.
7. Geller AK, O'connor MK. The sickle Cell Crisis: a dilemma in pain relief. *Mayo Chen Proc.* 2008; 359(21):2254-6.
8. Yenamandra A, Marjoncu D. Voxelotor: A Hemoglobin S Polymerization inhibitor for the treatment of sickle cell disease. *J Adv Pract Oncol.* 2020; 11(8):873-877.
9. Stevens DL, Hix M, Gildon BL. Crizanlizumab for the Prevention of Vaso-Occlusive Pain Crises in Sickle Cell Disease. *J Pharm Technol.* 2021;37(4):209-215.
10. Elenga N, Loko G, Etienne-Julan M, Al-Okka R, Adel AM, Yassin MA. Real-World data on efficacy of L-glutamine in preventing sickle cell disease-related complications in pediatric and adult patients. *Front Med (Lausanne).* 2022;9:931925.
11. World Agroforestry Centre (WAC). *Khaya senegalensis*. In: Agroforestry Database. <<http://www.worldagroforestry.org/Sites/TreeDBS/AFT/SpeciesInfo.cfm?SpID=1027>>[Accessed 14 September 2004].
12. Fall AB, Vanhaelen-Fastre R, Vanhaelen M, Lo I, Toppet M, Fester A, Fondu P. in vitro Anti-sickling Activity of a re-arranged Limonoid isolated from *Khaya senegalensis*. *Planta Med.* 1999; 65(3):209-12.
13. Egunyomi A, Moody JO, Eletu MO. Anti-sickling Activities of Two Ethno-medicinal Plant Recipes used for the Management of Sickle Cell Anemia in Ibadan, *Nigeria. Afr. J. Biotechnol.* 2009; 8(1): 020-025.
14. Cyril-Olutayo CM, Agbedahunsi JM. Effects of the Ethanolic Extracts of *Cniscodolus aconitifolius* (Mill.) I.M. Johnst. on Hbs Red Blood Cells In-vitro. *Nig. J. Nat. Prod. and Med.* 2015; Vol. 19, 116-122.

15. Official Methods of Analysis of the Association of Official Analytical Chemistry (AOAC). 13th edition Washington D.C. P1094. United States of America (1987).
16. Official Methods of Analysis of the Association of Official Analytical Chemistry (AOAC). Revised edition Washington D.C. United States of America (2006).
17. Rittner U, Jess DR. Phenolics and In-vitro degradability of protein and fibre in West Africa Browse. *Journal of science of food and agriculture*(1992); 58, (1):21-28.
18. Mishra S, Sonter S, Dwivedi MK, Singh PK. Anti sickling potential and chemical profiling of traditionally used *Woodfordia fruticosa* (L.) Kurz leaves, *Arabian Journal of Chemistry*, 2022;15:1, Doi:[10.1016/j.arabjc.2021.103539](https://doi.org/10.1016/j.arabjc.2021.103539)
19. Njayou FN, Amougou AM, Fouemene Tsayem R, Njikam Manjia J, Rudraiah S, Bradley B, Manautou JE, Fewou Moundipa P. Antioxidant fractions of *Khaya grandifoliola* C.DC. and *Entada africana* Guill. et Perr. induce nuclear translocation of Nrf2 in HC-04 cells. *Cell Stress Chaperones*. 2015;20(6):991-1000. doi: 10.1007/s12192-015-0628-6.
20. British Pharmacopoeia. Ash value, acid insoluble, water soluble, extractive and alcohol extractive, 11, appendix xii, His majesty's Stationary Office, London (1980); pp. 1276 – 1277.
21. Ajazuddin A, Saraf S. Evaluation of physicochemical and phytochemical properties of Safoof-E-Sana, a Unani polyherbal formulation. *Pharmacognosy Res.*2010; 2(5):318-22.
22. Oduntan A, Olaleye O, Akinwande B. Effect of Plant Maturity on the Proximate Composition of *Sesamum Radiatum* Schum Leaves. *Journal of Food Studies*. 2012;1(1):69-76
23. Iheanacho K, Ubebani AC. Nutritional composition of some leafy vegetables consumed in Imo state, *Nigeria. J. Appl. Sci. Environ Manage.* 2009; 13(3):35-38.

24. Isong E, Idiong U. Comparative studies on the nutritional and toxic composition of three varieties of *Lesianthera africana*. *Plant Foods Hum. Nutr.* 1997; 51:79–84.
25. Asaolu SS, Adefemi O, Oyakilome IG, Ajibulu KE, Asaolu MF. Proximate and Mineral composition of Nigerian leafy vegetables. *J. Food Res* 2012; 1(3):
26. Shaikh S, Yaqoob M, Kumar A, Mca S. Importance in dietary fiber in food: a review. *SSRN Electronic Journal.* 2019; 6:810-817.
27. Sena LP, Vanderjagt DJ, Rivera C, Tsin AT, Muhamadu I, Mahamadou O, Millson M, Pastuszyn A, Glew RH. Analysis of nutritional components of eight famine foods of the Republic of Niger. *Plant Foods Hum Nutr.* 1998; 52(1):17-30.
28. Lockett CT, Calvert CC, Grivetti, LE. Energy and micronutrient composition of dietary and medicinal wild plants consumed during drought. Study of rural Fulani, Northeastern Nigeria. *Int J Food Sci Nutr.* 2000; 51:195- 208
29. Hassan LG, Umar KJ. Nutritional Value of Balsam Apple (*Momordica balsamina* L.) Leaves. *Pak J. Nutr.* (2006); 5: 522-529.
30. Ali A. Proximate and mineral composition of the Marcubeh (*Asparagus officinalis*). *World j. dairy food sci.* 2009; 4(2):142-149.
31. Garcia-Rico L, Leyva-Perez J, Jara-Marini ME. Content and daily intake of copper, zinc, lead, cadmium and mercury from dietary supplements in Mexico. *Food Chem. Toxicol.* 2007; 45:1599-1605.
32. Gomez MR, Cerutti S, Olsina RA, Silvia MF, Martínez LD. Metal content monitoring in *Hypericum perforatum* pharmaceutical derivatives by atomic absorption and emission spectrometry. *J Pharm Biomed Anal.* 2004; 24:569-576.

33. Underwood EU. Trace metals in human and animal nutrition, 3rd ed., Academic Press, New York, USA. 1971; 253.

34. Akpanyung EO. Proximate and Mineral composition of bouillon cubes produced in Nigeria. *Pak J Nutr.* 2005; 4(5):327-329.

UNDER PEER REVIEW