Topography based soil profile characterization of KVK farm, Sakhigopal, Puri, Odisha located in the East and Southeastern Coastal Plain Agro-Climatic Zone of Odisha

Abstract

To identify soil-related crop production constraints and to provide the necessary information for soil management decisions, horizon-wise profile soil samples (from three profiles) at three topographic positions present at upland,medium landandlowland from KVK farm, Sakhigopal were collected and analysed for various physico-chemical, and nutrient parameters. The soil's physical and chemical properties andavailable nutrient status showed distinct variations among the pedons. The soil texture varied from sandy to sandy clay loam; bulk density and particle density values varied from 1.31 to 1.91 and 2.33 to 2.68 Mgm^{-3}, respectively. Soil pH ranged from 6.32 to 7.50 and the electrical conductivity was recorded below 1 dSm^{-1}. The organic carbon content ranged from 8.6 to 10.3 g kg^{-1}. Available nitrogen,phosphorus, potassium, and sulphur contents increased towards the lower topographic position, which maybe attributed to increased levels of organic carbon content in the lower topographic positionowing to high moisture content and higher cropping intensity. Proper land use planning and soil management practices can helpin optimizingcropproductivityas well as sustainingsoil health.

Keywords: topography, Sakhigopal, pedons, soil profile characterization, nutrients, physico-chemical soil properties

1. Introduction

Soil fertility and crop productivity are significantly impacted by the properties of the soil profile, which are influenced by several soil-forming processes (Digal et al. 2018, Sethy et al. 2019, Swain et al. 2019, Lokya et al. 2020, Singh et al. 2021,Pattanayak et al., 2023). Soil genesis has a significant impact on soil physico-chemical properties and available nutrient status as well (Dash et al. 2019a, 2019b). The physico-chemical properties and distribution of available macro nutrients provide a comprehensive idea about the fertility status of the studied area (Revathi et al., 2024). Of all the variables that contribute to soil formation, topography and parent material have a greater impact (Dash et al., 2019c). Given that soil and water are the primary natural resources needed for both crop production and

UNDER PEER REVIEW

human habitation, a thorough investigation that includes soil profile characterization and evaluation of soil nutrient statushave a greater importance (Dash et al., 2022a, 2022b).

Studying variations in soil properties along a toposequence notonly helpstodetecttopography relatedcropproductionlimits,butalsohelpsinselecting suitable crops, cropping methods, land use planning and soil and watermanagementactivitiesfordiverselandtypes (Dash et al., 2022c).Additionally, it will be helpful to identify soil fertility-related agricultural production constraints on the farm based on which necessary remedial measures can be recommended for maximizing crop yield. Moreover, any research on soil survey is a great asset for future research purposes. Since works on soil characterization on various landtypesofKVKfarm,Sakhigopalhavenotbeenpreviously carriedout, therefore,inthe present study,an attempthas been intended tocharacterize andclassify soils bystudying representative pedons at different topographic positions viz. upland, medium land and lowland ofKVKfarm, Sakhigopal.

2. Materials and Methods

2.1. Thestudy area: KVK, Sakhigopal

Sakhigopal, one of the oldest Coconut ResearchStation in our country was established in the year 1955 by the Government of Odisha. The study area is located in the East and South Eastern Coastal Plain AgroClimaticZoneofOdisha.Itislocatedatadistanceof20kilometersfromtheBayofBengal.It is situated between $19^0$48' N latitude and $85^0$52' E longitude. The study area has been divided into three physiographic units based on slopeand elevation such as gently sloping upland (pedon-1; elevation of 30 feet above Mean SeaLevel), very gently sloping medium land (pedon-2; elevation of 26 feet above Mean Sea Level)and nearly levellow land (pedon-3; elevation of 22 feetabove Mean SeaLevel). The climate of the study area is relatively warm and humid, with shortandmild winters. The mean annual rainfall is 1409 mm, mainly due to the southwestmonsoon. The maximum average summer temperature is 39^0C. The summer seasonstarts from March to mid-June, followed by rainy season from June to September.

Comment [Ma1]: Stations

Comment [Ma2]: By the rainy

2.2. Methodsof soilsurveyandsamplecollection:

The landforms of the researchstation were determined by exploring theareaand using a GPS device (Garmin MAKE; model: 76MAPCSx) to monitor and recordelevationdata aboveMeanSeaLevel (MSL)at several locations. In the research area, three typical soil

UNDER PEER REVIEW

profiles (pedons) were dug based on thetype of land viz., upland, medium land and low land comprising of 7, 8 and 7 horizonsrespectively.

Comment [Ma3]: remove

2.3. Soil analysis

The soil samples from three soil profiles collected from the study area were air driedand ground with a wooden hammer and passed through a 2 mm sieve for analysis of various physico chemical properties.The textural classes of soil were determined by Bouyoucous hydrometer method (Bouyoucos, 1962). Soil pH was determined in 1:2.5 soil water ratio by pH meter as described by Jackson (1973). As suggested by Jackson (1973), the electrical conductivity of soil saturated paste was determined in 1:2.5 soil-water suspension by a conductivity meter. The organic carbon content of soil was determined by wet digestion procedure of Walkley and Black (1934). Furthermore the soils were analyzed for bulk density (Klute 1986), particle density (Chopra and Kanwar 1986), water holding capacity (Piper 1950), cation exchange capacity (Chapman 1965), exchangeable acidity (Thomas 1996), exchangeable cations (Page et al. 1982), available nitrogen (Subbiah and Asija, 1956), phosphorus (Bray and Kurtz, 1945), potassium (Hanway and Heidel, 1952) and sulphur (Chesnin and Yien, 1950).

3. Results and Discussion

3.1. Physical properties of pedons

Physical characteristics of the soil properties of pedon 1, 2, and 3 have been presented in Table 2a, 2b, and 2c, respectively.

3.1.1. Particle size distribution

The soil texture varied from sandy to loamy sand for pedon1 and 2 and sandy to sandy clay loam for pedon 3. Thepercentageofsand,siltandclay contentrangedbetween 88.4 to 97.4, 0.2 to 4 and 4.6 to 9.6 percentage in pedon 1 (Table 1a); 88.4 to94.4, 0.2 to 4 and 4 to 7.6 percentage in pedon 2 (Table 1b) and 64.4to92.4,2to14 and 5.6 to 23.6 percentage in pedon 3 (Table 1c), respectively. The sand content increased gradually from upper horizons towards lower horizons. Large coarse fragments were found throughout the soil profile which might be attributed to soil erosion and absence of elluviation process. The clay content was found to be decreasing with depth in all the three pedons. Similar results were observed by Dash (2018).

Comment [Ma4]: eluviation

3.1.2. Bulk density

Pedon 1, pedon 2 and pedon 3 recorded the bulk density values in the range of 1.62 to 1.91 Mg m^{-3}, 1.53 to 1.90 Mg m^{-3} and 1.31 to 1.51 Mgm^{-3} respectively. Higher bulk density values were recorded in lower most horizons due to higher sand content, lower clay

UNDER PEER REVIEW

content, low organic carbon content and increased effect of compaction, whereas the bulk density was lower in surface horizons due to high organic carbon content. Similar findings were reported by Dash (2019a, 2022a) and Barla (2021).

3.1.3. Particle density

The particle density values of 2.44to2.68Mgm^{-3},2.36 to 2.67 Mg m^{-3} and 2.33to 2.58Mgm^{-3} were recorded in pedon 1, 2 and 3 respectively. The increasing order of particle density values from upper to lower horizons is might be attributed to lower organic carbon in the lower horizons than that of the upper horizons of the soil profiles. Acharya (2021) reported similar results.

Comment [Ma5]: horizons might

3.1.4. Total porosity

The total porosity values of pedon1, 2 and 3 varied from 26.4 to 34%, 28.9 to 36.3% and 41.5 to 45%, respectively. The total porosity in all the three pedons was found to be positively correlated with percentage clay. Dash (2019a) found similar observations.

3.1.5. Maximum water holding capacity

The maximum water holding capacity (WHC) values varying from 27.2 to 35.6%, 31.5 to 35.79% and 33.6 to 44.2% were recorded for pedon 1, 2 and 3, respectively. In all three pedons, WHC was seen to be decreasing with depth, which may be related to the varying clay percentages in the pedons.

3.2. Chemical characteristics of pedons

Chemical characteristics of the soil properties of pedon 1, 2, and 3 have been presented in Table 2a, 2b, and 2c, respectively.

3.2.1. Soil reaction

In pedon 1, the surface soil and lower horizon recorded pH values of 6.80 and 7.50 respectively. The slightly acidic surface soil of pedon 2 had a pH value of 6.34 and increased upto 6.55 with depth at lower horizon. The pH values of 6.32 and 6.68 were recorded in surface and lower most horizon of pedon 3, respectively. The pH values of all three pedons showed a consistent trend of increasing from upper to lower horizons due to leaching of basic cations from upper to lower horizons followed by heavy rainfall. Similar results were reported by Barla (2021) and Kumar et al. (2012).

Comment [Ma6]: in the surface

Comment [Ma7]: must be in iatlic

3.2.2. Electrical conductivity

UNDER PEER REVIEW

The electrical conductivity (EC) of all the three soil profiles remained below 1 dSm^{-1}, making them safe for all types of agricultural production without any salinity hazard. Such lower electrical conductivity could be attributed to leaching of soluble salts during intensive rainfall as prevalent in the study area.

3.2.3. Organic carbon

The organic carbon content of surface horizons of pedon 1, 2 and 3 were found to be 8.6, 9.1 and 10.3 g kg^{-1}, respectively and decreased with depth across the three soil profiles. Fresh accumulation and decomposition of crop residues in the surface horizons of all pedonscontributed higher organic carbon in the upper horizons. Similar findings were reported by Dorji et al. (2014), Kumar et al. (2012) and Khanday et al. (2018).

3.2.4. Exchangeable bases, cation exchange capacity (CEC), and base saturation

The concentration of exchangeable bases *viz* calcium, magnesium, sodium and potassium in different pedons was found to be in the order of Ca^{2+} >Mg^{2+} >Na^{+} >K^{+}. The CEC values of 13.3, 12.8and 13.6 cmol(p^{+})kg^{-1} were recorded in the surface horizons of pedon 1, 2 and 3, respectively, which gradually decreased with depth in all the threepedons. This could be due to decrease in clay percentage with soil depth.These results are in conformity with the findings of Pattanayak (2016). The surface horizons of pedon 1, 2 and 3 recorded base saturation percentage of 65.4, 68.7 and 75.6, respectively. Base saturation was observed to be increasing with soil depth, which could be attributed to increase in the basic cations with depth in all the three pedons.

Comment [Ma8]: to a decrease

Comment [Ma9]: percentages

3.2.5. Exchangeable Sodium percentage (ESP)

The ESP varied from 5.2 to 5.6, 3.5 to 5.8 and 4.4 to 5.9% in pedon 1, 2 and 3, respectively. The gradual increment of ESP with depth in all the three soil profiles could possibly be due to leaching of sodium ions from upper to lower horizons during intensive rainfall and the findings are in agreement with the findings of Mishra (2008).

3.3. Depth wise vertical distribution of available macro nutrients in different pedons

Thedepth-wise vertical distribution of available macronutrients *viz*. nitrogen, phosphorus, potassium and sulphur are presented in Table 3. In pedon 1, the depth wise available N, P, K and S content varied from 142 to 198, 12.7 to 22, 447 to 548 and 1.7 to 1.8 kg ha^{-1}. Similarly, the N, P, K and S content ranged from 137 to 218, 16 to 37.5, 469 to 561, 2.2 to 3.3 kg ha^{-}

UNDER PEER REVIEW

[1]inpedon 2 and 162 to 240, 14 to 37.7, 481 to 565 and 3.2 to 5.2 kg ha^{-1}inpedon 3, respectively. The concentration of available N, P, S was higher in surface horizons and the lowest concentration was in the bottom horizons. In all the three pedons, available N, P, S content decreased with increase in depth, whereas available K was found to be increasing with the soildepth. The increased concentrations of nitrogen, phosphorus and sulphur towards lower topographic positions could be attributed to higher organic carbon content. The presence of potassium bearing parent materials like Feldspar and Mica could be the reason for increased potassium availability with soil depth (Dash et al., 2019b; Kishore et al., 2020; Kishore et al., 2022).

4. Conclusion

In the present investigation, three representative pedons located at three distinct topographic positions viz., upland, medium land, and low land inside the premises of KVK farm Sakhigopal located in the Puri district of Odisha were studied. Depth wise vertical distribution of physico-chemical properties and available macro nutrient status were characterized. It was observed that the bulk density and particle density values were relatively lower in the upper horizons and higher in the lower horizons. In all three pedons, the pH of the soil progressively rose with increasing depth and the electrical conductivity values of less than 1 dSm^{-1} were recorded for each pedon. The organic carbon content was high in the surface layers of all pedons. The exchangeable acidity showed a decreasing trend with soil depth due to increase in other exchangeable cations saturating the exchange sites. In all the three pedons, calcium and magnesium constituted the majority of exchangeable cations. The cation exchange capacity values of the soils decreased vertically with increasing depth whereas, the percentage base saturation of soils increased steadily as depth decreased vertically. The decrease in the concentrations of available macronutrients such as nitrogen, phosphorus, and sulphur with soil depth can be explained by the declining organic matter content with depth. However, the available potassium increased with depth in all three pedons, possibly due to the nature of parent material that have been weathered and transported by rivers.

5. References

AcharyaBP.2021.PreparationofGPSandGISbasedsoilfertilitymapsandidentificationofsoilrelatedcropproductionconstraintsofKVKFarm,Jashipur,M.Sc.Thesis,DepartmentofSoilScienceandAgriculturalChemistry,Collegeof Agriculture,OUAT, Bhubaneswar.

UNDER PEER REVIEW

BarlaFX.2021.PreparationofGPSandGISbasedsoilfertilitymapsandidentification of Soil related Crop production constraints of KVK Farm, R.Udayagiri,M.Sc.Thesis,DepartmentofSoilScienceandAgriculturalChemistry,Colle geof Agriculture, OUAT,Bhubaneswar.

Bouyoucos GJ. 1962. Hydrometer method improved for making particle size analysisofsoils,*AgronomyJournal*, **54**: 464.

Bray RH and Kurtz Lt. 1945. Determination of total, organic and available forms insoils,SoilScience,**59**: 39-45.

ChapmanHD.1965.MethodsofSoilAnalysis,Part-II,AmericanSocietyofAgronomy.Inc. Winconsin, USA: 891-900.

ChesninLandYienCH.1950.Turbidimetricdeterminationofavailablesulphates,

ChopraSLandKanwarJS.1986.AnalyticalAgriculturalChemistry,KalyaniPublishers,New Delhi.

Dash PK. Chapter 22 - Remote sensing as a potential tool for advancing digital soil mapping, in: Dharumarajan, S., Kaliraj, S., Adhikari K, Lalitha M, Kumar N. (Eds.), Remote Sensing of Soils. Elsevier. 2024. 357–370.

Dash PK, Mishra A, Patro H, Nayak RK, Jena B, Satapathy M .2022a. Soil characterization along a toposequence located in North-Central Plateau Agro-climatic zone of Odisha. Ecology Environ Conserv, pp 471—476.

Dash PK, Mishra A, Patro H, Patra RK, Sahoo BC, Panda RK, Saren S .2022b. Morphological Characterization of Soil Profiles in a Toposequence Located in the North-Eastern Coastal Plains Agroclimatic Zone of Odisha. Environ Ecol40 : 1762—1772.

Dash PK, Mishra A, Saren S .2019a. Characterization and taxonomic classification of soils under a toposequence located in Eastern India. Environ Ecol 37 (4) : 1240—1249.

Dash PK, Mishra A, Saren S .2019b. Vertical distribution of available nutrients in an Eastern Indian Catena. Annals Pl Soil Res 21 : 320—325.

Dash PK, Mishra A, Saren S, Revathi B and Sethy SK. 2018. Preparation of GPS andGIS based soil fertility maps and identification of soil related crop productionconstraints of RRTTS and KVK farm, Dhenkanal located in the mid centraltablelandagroclimaticzoneofOdisha,India,*InternationalJournalofChemicalStudi es*,**6**: 934-943.

UNDER PEER REVIEW

Dash PK, Nayak B, Mishra A, Saren S .2019c. Geomorphological Characterization of Soils under a Toposequence Located in Eastern India. Environ Ecol, pp 37.

Dash PK, Panigrahi N, Mishra A.2022c. Identifying opportunities to improve digital soil mapping in India: A systematic review. Geoderma Regional. 28 : e00478. https://doi.org/ 10.1016/J.GEODRS. 2021.E00478 1349—1359.

Digal M, Saren S, Mishra A, Dash PK, Swain N, Acharya BP .2018. Soil fertility status of some villages in Phiringia Block of Kandhamal District under North-Eastern Ghat Agro-Climatic Zone of Odisha, India. J PharmacogPhytochem 7 (6) : 659—662.

Dorji T, Odeh IOA and Field DJ. 2014. Vertical distribution of soil organic carbondensity in relation to land use cover, altitude and slope aspect in the EasternHimalayas, *Land*, **3**: 1232-1250.

Hanway JJ and Heidel H. 1952. Soil analysis methods as used in Iowa State College,SoilTesting Laboratory,IowaStateCollege Bulletin,**57**:1-31.

Jackson ML. 1973. Soil Chemical Analysis, Prentice Hall of India. Private Limited,NewDelhi.

KhandayM,WaniJA,RamDandKumarS.2018.Depthwisedistributionofavailablenutrientsofsoilsofhorticulturegrowingareasof GanderbalDistrictofKashmirvalley,*JournalofPharmacognosyandPhytochemistry*,**7**(1):19-22.

Kishore M, Baskar A, Dash PK, Bagavathi AU, Mishra A, Saren S .2020. Distribution of different potassium fractions in the soil profiles of Karaikal, Puducherry. *Int J Chem Studies* 8(2): 01—05.

Kishore MA, Mishra A, Dash PK, Saren S, Das TK, Priyadarsini S, Sethy H, Sardar MS. 2022. Vertical distribution of available n utrients in Boudh Block of Boudh District Catena. *Int J Pl Soil Sci* 34: 1681—1686.

Klute. 1986. Methods of Soil Analysis, part I, American Society of Agronomy, SoilandSocietyof America,371-373.

Kumar R, Rawat KS and Yadav B. 2012. Vertical distribution of physico-chemicalpropertiesunderdifferenttoposequenceinsoilsofJharkhand,*JournalofAgriculturalPhysics*, **12**(1): 63-69.

Lokya T, Mishra A, Saren S, Dash PK .2020. GPS and GIS-based soil Fertility maps of KVK farm, Nayagarh located in the North-Eastern Ghat Agro-Climatic zone of Odisha. *Ind J PharmacogPhytochem*Sp 9 (4) : 599—605.

UNDER PEER REVIEW

MishraA,PradhanNK,NandaSKandJenaB.2008.Soiltestbasedfertilizerrecommendationfor targetedyield of Sesamum (*Sesamum indicum*) underrice-sesamum cropping system in an *Inceptisol* of Orissa, *Environment andEcology*,**26**(4A): 1756-1758

Page AL, Miller RH and Keeney DR. 1982. Methods of Soil Analysis, part-2 (Edn.),Monographno-9,AmericanSociety of Agronomy,Agronomy seriesASASSA,Publishers, Medision, Wisconsin, USA, 621-622.

Pattanayak T .2016. Preparation of GPS-based soil fertility maps and identification of soil-related crop production constraints for Dhenkanal District, Odisha (PhD thesis). Department of chemistry, institute of technical education and Research, Siksha 'O' Anusandhan University, Bhubaneswar.

Pattanayak T. 2016. Preparation of GPS based soil fertility maps and identification ofsoil related crop production constraints for Dhenkanal District, Odisha, Ph. DThesis,DepartmentofChemistry,InstituteofTechnicalEducationandResearch,Siksha "O"AnusandhanUniversity,Bhubaneswar.

PiperCS .1950. Soil andPlantanalysis, UniversityofAdelaide, pp-368.*Proceedingsof Soil ScienceSocietyof America*,**14**: 149-51.

RevathiB, Saren S, Dash, PK, Mishra A, Sethy SK, Patnaik T, Dey P. 2024. Optimizing Nutrient Recommendations for Bottle Gourd Crop by Formulation of Targeted Yield Equations in an Inceptisol of Odisha, India,*Asian Journal of Soil Science and Plant Nutrition*, 10 (2): 223-235.

Sethy SK, Mishra A, Dash PK, Sraen S, Dey P .2019. Geo-Infor-mation based soil fertility status of Deogarh district of Odisha. *Ind Int J Curr Microbiol Appl Sci* 8(12) : 255—262.

Singh RP, Mishra A, Dash PK, Saren S, Mallik H, Mishra BB .2021. Soil fertility assessment of some villages in Kankadahad block of Odisha. *Environ Ecol*39 (1A): 250—255.

Subbiah BV and Asija GL. 1956. A rapid procedure for the determination of availablenitrogenin soils,*Current Science*, **25**:259-25260.

Swain N, Mishra A, Saren S, Dash PK, Digal M, Mishra BB .2019. Soil fertility status of some villages in Khordha and Bhubaneswar block of Khordha district under Northeastern Ghat agro climatic zone of Odisha, India. *Int J Curr Microbiol Appl Sci* 1 (8) : 2319—7706.

Thomas GW. 1996. Soil pH and Soil Acidity. In Sparks D.L.ed., *Methods of SoilAnalysis*, Part 3, Chemical Methods Madison, WI, Soil Science Society ofAmerica,*American*

UNDER PEER REVIEW

Society of Agronomy, 475-490.

Walkley AJ and Black CA. 1934. Estimation of soil organic carbon by the chromicacidtitration method,*Soil Science*, **37**: 29-38.

UNDER PEER REVIEW

Table 1a Physical characteristics of Pedon 1 (Upland)

Horizon	Depth (cm)	Sand	Silt	Clay	Textural Class	Bulk density	Particle density	Porosity	WHC
		(%)				(Mg m^{-3})		(%)	
Ap	0-12	88.4	2	9.6	Loamy sand	1.62	2.44	34	35.6
C1	12-27	88.4	4	7.6	Loamy sand	1.63	2.45	33.5	32.4
C2	27-60	92.4	2	5.6	Sand	1.77	2.54	27.2	27.2
C3	60-96	94.4	0.2	5.4	Sand	1.90	2.65	28.4	28.9

Table 1b Physical characteristics of Pedon 2 (Medium land)

Horizon	Depth (cm)	Sand	Silt	Clay	Textural Class	Bulk density	Particle density	Porosity	WHC
		(%)				(Mg m^{-3})		(%)	
Ap	0-12	92.4	2	5.6	Sand	1.53	2.36	35.2	33.7
A2	12-27	94.4	0.2	5.4	Sand	1.56	2.44	36.1	32.8
C1	27-39	94.4	0.2	5.4	Sand	1.58	2.48	36.3	31.5
C2	39-79	88.4	4	7.6	Loamy sand	1.64	2.56	34.5	35.8
C3	79-115	92.4	2	5.6	Sand	1.84	2.65	29.6	34.2

Table 1c Physical characteristics of Pedon 3 (Lowland)

Horizon	Depth (cm)	Sand	Silt	Clay	Textural Class	Bulk density	Particle density	Porosity	WHC
		(%)				(Mg m^{-3})		(%)	
Ap	0-12	64.4	12	23.6	Sandy Clay loam	1.31	2.33	44.8	37.4
C1	12-30	66.4	15	21.6	Sandy Clay loam	1.33	2.37	44	43.5
C2	30-75	64.4	14	21.6	Sandy Clay loam	1.35	2.45	44.3	44.2
C3	75-120	92.4	2	5.6	Sand	1.42	2.53	44	34.8

UNDER PEER REVIEW

Table 2a Chemical characteristics of Pedon 1 (Upland)

Horizon	Depth (cm)	pHw (1:2.5)	ECw (1:2.5)	Organic carbon (g kg^{-1})	Exchangeable cations [cmol(p$^+$) kg^{-1}]					Total exchangeable acidity	Cation Exchange Capacity	Base saturation	ESP
					Ca	Mg	Na	K	Sum	[cmol(p$^+$) kg^{-1}]		(%)	
Ap	0-12	6.80	0.01	8.6	6.6	0.9	0.7	0.5	8.7	1.3	13.3	65.4	5.2
C1	12-27	6.84	0.02	8.1	6.4	0.8	0.7	0.6	8.5	1.1	12.8	66.5	5.4
C2	27-60	7.43	0.02	6.8	6.3	0.7	0.5	0.7	8.2	1.1	12.3	66.6	5.6
C3	60-96	7.46	0.03	5.2	6.5	0.8	0.7	0.5	8.3	0.8	12.3	67.5	5.6

Table 2b Chemical characteristics of Pedon 2 (Medium land)

Horizon	Depth (cm)	pHw (1:2.5)	ECw (1:2.5)	Organic carbon (g kg^{-1})	Exchangeable cations [cmol(p$^+$) kg^{-1}]					Total exchangeable acidity	Cation Exchange Capacity	Base saturation	ESP
					Ca	Mg	Na	K	Sum	[cmol(p$^+$) kg^{-1}]		(%)	
Ap	0-12	6.34	0.04	9.1	7.8	0.3	0.2	0.5	8.8	1.1	12.8	68.7	3.5
A2	12-27	6.37	0.05	8.8	7.9	0.2	0.2	0.5	8.8	1.1	12.7	69.2	4.0
C1	27-39	6.43	0.05	7.6	7.5	0.3	0.2	0.7	8.6	0.9	12.6	68.2	5.4
C2	39-79	6.47	0.06	6.5	7.9	0.4	0.6	0.7	9.8	0.8	12.7	77.1	5.5
C3	79-115	6.53	0.06	5.7	7.6	0.5	0.6	0.7	9.4	0.7	11.7	80.3	5.8

UNDER PEER REVIEW

Table 2c Chemical characteristics of Pedon 3 (Low land)

Horizon	Depth (cm)	pHw (1:2.5)	ECw (1:2.5)	Organic carbon (g kg^{-1})	Exchangeable cations [cmol(p^{+}) kg^{-1}]					Total exchangeable acidity	Cation Exchange Capacity	Base saturation	ESP
					Ca	Mg	Na	K	Sum	[cmol(p^{+}) kg^{-1}]		(%)	
Ap	0-12	6.32	0.07	14.8	8.8	0.7	0.6	1.1	11.2	1	14.8	75.6	4.4
C1	12-30	6.47	0.07	14.3	8.7	0.7	0.7	1.0	11.1	1	14.4	77	5.5
C2	30-75	6.56	0.08	14.4	8.8	0.7	0.7	0.9	11.1	0.7	14.3	77.6	5.8
C3	75-120	6.70	0.08	13.6	8.6	0.6	0.8	0.9	10.9	0.5	13.6	80	5.9

UNDER PEER REVIEW

Table 3. Depth wise distribution of available nutrients in representative pedons of the study

Horizon	Depth (cm)	N	P	K	S
		(kg ha^{-1})			
Pedon 1 (Upland)					
Ap	0-12	198	22	447	1.8
C1	12-27	175	20.8	536	1.8
C2	27-60	166	19.4	543	1.7
C3	60-96	142	12.7	548	1.7
Pedon 2 (Medium land)					
Ap	0-12	218	37.5	470	2.6
A2	12-27	195	28.2	469	2.8
C1	27-39	176	25.3	552	3.3
C2	39-79	158	21.4	556	2.4
C3	79-115	137	16	561	2.2
Pedon 3 (Lowland)					
Ap	0-12	240	37.7	481	4.8
C1	12-30	196	23	552	5.2
C2	30-75	181	19.7	554	4.2
C3	75-120	162	14	565	3.2

UNDER PEER REVIEW