

# QUALITY PARAMETERS OF TRAY DRIED JACKFRUIT SEED FLOUR

## ABSTARCT

India has a large jackfruit farming industry; however, after the edible part is eaten, the nutrient-rich seeds are thrown away as waste. The study was aimed at investigating the functional properties and sensorial attributes of jackfruit seed flour from various treatments. The results revealed a significant difference between the functional characteristics and sensory attributes of jackfruit seed flour samples processed by soaking the seeds in various soaking media. Based on the results, two seed flour samples (VS<sub>1</sub> and KS<sub>1</sub>) were selected as the best and evaluated for their proximate composition. The cultivar *Varikka* had more crude fiber (2.96%), ash (2.61%), protein (13.17 g/100g) and carbohydrate content (76.82 g/100g). Both the seed flour samples had negligible amount of fat (VS<sub>1</sub> – 1.72 and KS<sub>2</sub> – 1.68%), less fat allows for the prevention of rancidity issues and longer shelf life of the product. The tray drying process can alter the colour, aroma, and textural properties of the seed samples. This procedure makes it feasible to preserve jackfruit seeds by converting them into flour, extending their shelf life. The findings revealed that jackfruit seed flour can be employed as a key ingredient in a variety of nutrient-dense food products.

## Keywords

*Jackfruit seed, flour, soaking media, tray drying, functional properties.*

## 1 INTRODUCTION

The focus of research in recent years has been on the search for lesser-known and underutilised crops, many of which have the potential to be useful as human nourishment. The jackfruit, *Artocarpus heterophyllus*, is an Indian native and member of the mulberry family. It is widely distributed over the Indian states of Assam, West Bengal, Uttar Pradesh, Maharashtra, Kerala, Tamil Nadu, and Karnataka [16]. Jackfruit seeds possess important minerals such as magnesium, potassium, phosphorous, calcium, sodium, iron, copper, zinc, and manganese [13]. They perform a wide range of tasks, including serving as our bones' building blocks, affecting muscle and nerve activity, and balancing the body's water levels [35]. Jackfruit seeds are rich in dietary fibre and B-complex vitamins, and their high fibre content helps to lower the risk of heart disease, avoid constipation, and limit adipogenesis [34]. Lignans, flavones, and saponins found in jackfruit seeds exhibit antioxidant, anticancer, antiulcer, antihypertensive, and antiaging properties [30]. Every year, vast numbers of seeds are squandered due to challenges in processing and storing them. Jackfruit is considered an underutilized fruit due to a lack of understanding about effective usage, a lack of post-harvest equipment, and supply chain network deficiencies [33]. The seed flour can be added to baked goods for value addition because of their high carbohydrate content and other nutrients, without altering the functional and sensory qualities of the finished product. The tray dried seeds can be ground into flour and incorporated to various food products to increase their value and enhance the shelf life. The objective of this study therefore is to examine the impact of different soaking techniques on the functional properties, sensory attributes and proximate composition of jackfruit seed flour.

## 2 MATERIALS AND METHODS

### 2.1 Raw materials

The jackfruit cultivars *Koozha* (k) and *Varikka* (v), which are widely available in Kerala, were selected for the study. Whole jackfruits were procured from Instructional farm, College of Agriculture, Vellayani, Trivandrum.

#### 2.1.1 Processing of jackfruit seed flour

In the present investigation to obtain good-quality jackfruit seed flour, the jackfruit seeds of cultivars *Koozha* and *Varikka* were subjected to various soaking treatments. The white arils (seed coat) were peeled off and the seeds were soaked for 30 minutes in various soaking media, viz., S<sub>1</sub> - water, S<sub>2</sub> - 4% sodium metabisulphite, and S<sub>3</sub> - 3% sodium hydroxide. After soaking, the brown spermoderm were removed from the seeds using a stain-less steel knife. The seeds underwent cleaning and washing under running water. The seeds were gelatinized at 65°C for 10 minutes. Following gelatinization, the seeds were sliced and tray-dried at 110 °C for 12 hours. After tray drying, seeds from both cultivars were ground into powder, sieved through a fine wire mesh (150 micron), stored in polyethylene pouches, and used for further investigation.

## **2.2 Analytical methods**

### **2.2.1 Functional quality analysis of jackfruit seed flour from different treatments**

Food's functional properties are determined by its organoleptic, physical, and/or chemical characteristics. The functional qualities of foods and flours are impacted by the food's components, particularly carbohydrates, proteins, lipids and oils, moisture, fiber, ash, and other substances or food additives added to the food (flour), as well as the structures of these components [6]. Water absorption capacity and oil absorption capacity of the jackfruit seed flour was determined by the method of Niba et al., [22] and Beuchat [7]. Solubility index and swelling power were determined according to the method of Oladele and Aina [25]. Bulk density and flour dispersibility were determined by the method suggested by Okaka and Potter [24] and Islam et al., [14].

### **2.3 Evaluation of Sensory quality attributes of jackfruit seed flour samples**

The organoleptic evaluation the samples was performed by 25 semi trained panelists. To develop the sensory sample, 10 g of jackfruit seed flour from various treatments was combined with 200 ml of cold milk. The sensory attributes of the jack fruit seed flour, such as their colour and appearance, taste, texture, flavour, and overall acceptability, are evaluated using a 9-point hedonic scale (1= dislike extremely and 9= like extremely). The difference in the scores was analyzed using Kruskal – Wallis test.

#### **2.3.1 Selection of the best jackfruit seed flour samples**

Based on the results of organoleptic evaluation, the best jackfruit seed flour samples were chosen for quality analysis.

### **2.4 Determination of proximate composition of jackfruit seed flour samples**

Moisture, ash, crude protein and crude fat of the samples were determined by the AOAC [2] method. Crude fiber content was estimated by the method of AOAC [3]. The carbohydrate content was calculated by difference, (100- sum of the values for moisture, crude protein, crude fiber, crude fat and crude ash).

### **2.5 Statistical analysis**

All data obtained from various analysis were pooled and subjected to Completely Randomized Design (One-way ANOVA) using KAU-GRAPES software. The sensory scores were analysed using Kruskal – Wallis test.

## **3. RESULT AND DISCUSSION**

### **3.1 Functional properties of the jackfruit seed flour samples**

Functional properties of jackfruit seed flour samples from various treatments are depicted in Table. 1.

Water absorption capacity (WAC), also known as water absorption, is the amount of water (moisture) absorbed by food or flour in order to attain the desired consistency and produce a quality food product [6]. In the present investigation the WAC of Jackfruit seed flour ranged between 194.8 – 203.1 ml/100g. Water absorption capacity was found to be higher for the cultivar *Varikka* ( $VS_1$  – 202.1ml/100g,  $VS_2$  – 198.7 ml/100g and  $VS_3$  – 197.7 ml/100g). It may be ascribed to the greater concentration of carbohydrates (namely starch) and fiber in the *Varikka* cultivar compared to the *Koozha* cultivar. The increase in water absorption capacity of flour could also be attributed to an increase in amylose solubility and leaching, as well as the degradation of crystalline structure of starch [15]. The results are comparable with the values reported by Reddy et al.,[28] and Chowdary et al., [10] who reported the water absorption capacity of jackfruit seed flour as 184.27 ml/100g and 203.4%, respectively.

Oil absorption capacity is the binding of fat by the non-polar side chain of proteins. The rate of oil absorption is very high in foods with high protein content. The oil binding capacity of protein in food depend on the intrinsic factors such as protein conformation, amino acid composition, and surface polarity or hydrophobicity [32]. Oil absorption capacity was significantly higher for the cultivar *Varikka* ( $VS_1$  – 87.80 ml/100g,  $VS_2$  – 85.80 ml/100g and  $VS_3$  – 86.50 ml/100g). According to Suresh and Samsher [32], rate of oil absorption is high in food with high protein content. The findings abide by the study done by Borgis and Bharati [8] and Reddy et al., [28] who reported oil absorption capacity of jackfruit seed flour as 89.93 ml/100g and 87.65 ml/100g respectively.

The swelling capacity is the measure of the starch ability to absorb water and swell. The extent of swelling is determined by water absorption, temperature and availability of water. It is considered a quality requirement in high quality food compositions, such as bakery products [26]. The swelling power of the jackfruit seed flour samples ranged between 3.99 – 4.82 g/g. The highest swelling power was observed for the cultivar *Varikka* ( $VS_1$  – 4.82 g/g,  $VS_2$  – 4.28 g/g and  $VS_3$  – 4.18 g/g) than the cultivar *Koozha*. It could be attributed to the high starch content of the cultivar *Varikka*. The swelling capacity (index) of flours are also influenced by the particle size, species variety and method of processing [32]. The obtained values are comparable with the values reported by Butool and Butool [9] 3.62 g/g and Ocloo et al., [23] 4.77 g/g.

In the food system, solubility is the ability of solid, liquid, or gaseous food (chemical) compounds known as solutes to dissolve in a liquid, gaseous, or solid solvent. It is measured and determined in terms of the maximum quantity of solute dissolved in a given solvent at equilibrium [6]. In the present investigation solubility index of jackfruit seed flour ranged from 1.84 – 2.05%. Solubility index was found to be significantly higher for the cultivar *Varikka* ( $VS_1$  – 2.05%,  $VS_2$  – 1.97% and  $VS_3$  – 1.92%). The obtained values are similar with the values reported by Akter and Haque [1] 1.80% and Reddy et al., [28] 1.46%.

Bulk density may change depending on how the food material is handled; it is not an intrinsic property of a food material. Bulk density depends on the size of the particles in the samples and measures the heaviness of the flour. In the food processing industry, bulk density is important in determining to package and handling materials and used in the application of wet processes in the food industry [6]. The bulk density of the jackfruit seed flour samples ranged from 0.66 – 0.73 g/cm<sup>3</sup>. It was observed that the cultivar *Varikka* exhibited highest bulk density ( $VS_1$  – 0.73 g/cm<sup>3</sup>,  $VS_2$  – 0.71 g/cm<sup>3</sup> and  $VS_3$  – 0.70 g/cm<sup>3</sup>) than the cultivar *Koozha*. The variation in bulk density could be attributed to the cultivar *Varikka*'s higher starch content. The higher the starch content the more likely the increase in bulk density [15]. The findings abide by the study done by Nabubuya et al., [21] 0.70 g/cm<sup>3</sup> and Kamal et al., [18] 0.64 g/cm<sup>3</sup>.

Dispersibility describes flour's tendency to separate from water molecules and displays its hydrophobic effect. The higher dispersibility, the better is the reconstitution property (Kulkarni et al., 1991). According to Eke-Ejiofor et al., [12] the dispersibility percentage is a measure of flours strong water absorption capacity as well as the good quality of gel. In the present investigation the dispersibility of seed flour samples ranged

between 28.33% - 33.66%. The highest flour dispersibility was observed for the treatment VS<sub>1</sub> – 33.66%, while the treatments VS<sub>3</sub> (30.66%), KS<sub>2</sub> (30.33%) and KS<sub>1</sub> (28.33), KS<sub>3</sub> (28.66) were statistically found to be on par. The obtained values correspond to the values reported by Sultana et al.,[31] 28% and Arya et al., [5] 32.67%.

**Table no: 1 Functional properties of jackfruit seed flour from different treatments**

Treatments	WAC (ml/100g)	OAC (ml/100g)	SP (g/g)	SI (%)	BD (g/cm <sup>3</sup> )	Dispersibility (%)
VS <sub>1</sub>	202.1 <sup>a</sup>	87.80 <sup>a</sup>	4.82 <sup>a</sup>	2.05 <sup>a</sup>	0.73 <sup>a</sup>	33.66 <sup>a</sup>
VS <sub>2</sub>	198.7 <sup>b</sup>	85.80 <sup>c</sup>	4.28 <sup>b</sup>	1.97 <sup>b</sup>	0.71 <sup>b</sup>	31.64 <sup>b</sup>
VS <sub>3</sub>	197.7 <sup>c</sup>	86.50 <sup>b</sup>	4.18 <sup>c</sup>	1.92 <sup>c</sup>	0.70 <sup>b</sup>	30.66 <sup>c</sup>
KS <sub>1</sub>	196.7 <sup>d</sup>	84.62 <sup>f</sup>	4.09 <sup>d</sup>	1.84 <sup>e</sup>	0.66 <sup>e</sup>	28.33 <sup>d</sup>
KS <sub>2</sub>	195.0 <sup>e</sup>	85.11 <sup>d</sup>	4.01 <sup>e</sup>	1.87 <sup>d</sup>	0.67 <sup>d</sup>	30.33 <sup>c</sup>
KS <sub>3</sub>	194.8 <sup>e</sup>	84.90 <sup>e</sup>	3.99 <sup>e</sup>	1.85 <sup>e</sup>	0.69 <sup>c</sup>	28.66 <sup>d</sup>
±SE(m)	0.133	0.008	0.008	0.004	0.004	0.333
CV%	0.117	0.016	0.315	0.388	1.069	1.89

Values are means of triplicates. Values with different superscripts (a,b,c,d) within the same column are significantly different ( $P \leq 0.05$ ). VS<sub>1</sub>- Varikka soaked in water, VS<sub>2</sub>- Varikka soaked in 3% Sodium hydroxide, VS<sub>3</sub>- Varikka soaked in 4% Sodium metabisulphite, KS<sub>1</sub>- Koozha soaked in water, KS<sub>2</sub>- Koozha soaked in 3% Sodium hydroxide, KS<sub>3</sub>- Koozha soaked in 4% Sodium metabisulphite. WAC= Water absorption capacity, OAC= Oil absorption capacity, SI= Solubility index, SP= Swelling power, BD = Bulk density.

### 3.2 Evaluation of Sensory quality attributes of jackfruit seed flour samples

The sensory quality attributes of jackfruit seed flour are depicted in the following Table 2.

#### 3.2.1 Colour and Appearance

Colour and appearance aspects of products should not be overlooked because these features may render the product acceptable or unacceptable. The appearance is an attribute which a decision is taken to purchase or consume [29]. The mean value for colour and appearance ranged from 6.2 – 8.4. highest score was observed for the samples VS<sub>1</sub> (8.4) and KS<sub>1</sub> (8.1), they were found to be on par.

#### 3.2.2 Taste

Taste is a key aspect in determining whether a consumer would accept a certain food product. When food is ingested, its taste provides the consumer with crucial information about its quality and thus its acceptability. There is a strong correlation between the quality of taste and product's palatability [20]. The mean value for taste ranged from 5.4 – 8.1. There was a significant difference between the scores. The treatments VS<sub>1</sub> (8.1) and KS<sub>1</sub> (7.9) were statistically found to be on par.

### 3.2.3 Texture

Texture is a multimodal, multisensory food attribute. It is defined as the sensory, functional manifestation of the surface, mechanical, and structural qualities of foods that are perceived by touch, vision, hearing, and kinesthetic senses [11]. The mean value for the attribute texture ranged between 5.8 – 7.8. There was a significant difference between the jackfruit seed flour samples for the attribute texture.

### 3.2.4 Flavour

Flavor is defined as the sum of perceptions resulting from stimulation of the sense ends that are grouped together at the entrance of the alimentary and respiratory tracts. The mean value for flavour ranged from 5.5 – 7.9. For the attribute flavour highest score was obtained for the treatments VS<sub>1</sub> (7.9) and KS<sub>1</sub> (7.5), and they were statistically found to be on par.

### 3.2.5 Overall acceptability

Finding the treatment that performs the best across all of these criteria, including colour, appearance, taste, texture, and flavour, makes up overall acceptability. Food's sensory qualities, including taste, texture, flavour, and appearance, have distinct and significant effects on the overall acceptability of food [32]. The sensory assessment revealed that the treatments VS<sub>1</sub> (8.0) and KS<sub>1</sub> (7.7) had maximum score for overall acceptability and they were statistically found to be on par.

**Table no: 2 Sensory quality attributes of jackfruit seed flour samples**

Treatments	Colour and Appearance	Taste	Texture	Flavour	Overall Acceptability
VS <sub>1</sub>	8.4 <sup>a</sup>	8.1 <sup>a</sup>	7.8 <sup>a</sup>	7.9 <sup>a</sup>	8.0 <sup>a</sup>
VS <sub>2</sub>	6.6 <sup>b</sup>	5.4 <sup>b</sup>	6.2 <sup>b</sup>	5.6 <sup>b</sup>	5.9 <sup>bc</sup>
VS <sub>3</sub>	6.9 <sup>b</sup>	6.5 <sup>c</sup>	6.4 <sup>b</sup>	6.1 <sup>b</sup>	6.4 <sup>c</sup>
KS <sub>1</sub>	8.1 <sup>a</sup>	7.9 <sup>a</sup>	7.5 <sup>a</sup>	7.5 <sup>a</sup>	7.7 <sup>a</sup>
KS <sub>2</sub>	6.2 <sup>b</sup>	5.5 <sup>b</sup>	5.8 <sup>b</sup>	5.5 <sup>b</sup>	5.7 <sup>b</sup>
KS <sub>3</sub>	6.6 <sup>b</sup>	6.1 <sup>bc</sup>	5.9 <sup>b</sup>	6.0 <sup>b</sup>	6.1 <sup>bc</sup>
$\chi^2$	44.331	48.768	42.223	40.8	46.342
p_value	0	0	0	0	0

Values with different superscripts (a,b,c,d) within the same column are significantly different ( $P \leq 0.05$ ). VS<sub>1</sub>- Varikka soaked in water, VS<sub>2</sub>- Varikka soaked in 3% Sodium hydroxide, VS<sub>3</sub>- Varikka soaked in 4% Sodium metabisulphite, KS<sub>1</sub>- Koozha soaked in water, KS<sub>2</sub>- Koozha soaked in 3% Sodium hydroxide, KS<sub>3</sub>- Koozha soaked in 4% Sodium metabisulphite.

### 3.2.6 Selection of the best jackfruit seed flour samples

We chose the best seed flour samples based on the results of the sensory assessment. The sensory evaluation results demonstrated a substantial difference between the six jackfruit seed flour samples in terms of colour and appearance, taste, texture, flavour, and overall acceptability. The treatments VS<sub>1</sub> and KS<sub>1</sub> had the highest score for all the sensory attributes; hence, these two samples were selected for further quality analysis.

### 3.3 Proximate composition of jackfruit seed flour samples

The proximate composition of jackfruit seed flour samples is depicted in Table 3.

Moisture content is one of the most important parameters which determine the shelf-life quality of food product. In the present investigation moisture content of the jackfruit seed flour samples ranged from 7.79 – 7.81%. There was no significant difference observed between the jackfruit seed flour samples for moisture content. The results obtained were in agreement with the values reported by Nabubuya et al., [21] and Sultana et al., [31] who reported 7.8% and 8.1% of moisture content in jackfruit seed flour.

In the current investigation, the ash content of jackfruit seed flour samples ranged between 2.55 and 2.61%. It was found that the ash content was substantially higher for the cultivar *Varikka*. The ash content of jackfruit seed flour was reported as 2.57% and 2.60% by Arefin et al., [4] and Nabubuya et al., [21]. The obtained values correspond to the reported values.

Protein is a necessary ingredient for the body's vital processes. The protein content of the jackfruit seed flour samples ranged between 12.58 g/100g to 13.17 g/100g. The protein content was found to be significantly higher in the cultivar *Varikka*. The results are comparable with the values reported by Juárez-Barrientos et al., [17] 13.86 g/100 g and Eke-Ejiofor et al., [12] 12.45 g/100 g.

In the present investigation the fat content of jackfruit seed flour samples ranged between 1.68 – 1.72%. The fat content was found to be higher in the cultivar *Varikka*. Jackfruit seed flour has low lipid content, preventing rancidity and extending its shelf life. The results are consistent with the findings of Islam et al., [14] and Palamthodi et al., [27], who reported the fat content of jackfruit seed flour as 1.77% and 1.44%, respectively.

The crude fiber content of the jackfruit seed flour samples ranged between 2.87% - 2.96%. The crude fiber content of the jackfruit seed flour samples was found to be substantially higher in the cultivar *Varikka*. The findings abide by the study done by Sultana et al., [31] and Ocloo et al., [23], who reported the crude fiber content of jackfruit seed flour as 2.8% and 3.19%, respectively.

Carbohydrates are one of the most abundant and widespread organic substance in nature. The carbohydrate content of the jackfruit seed flour samples ranged from 74.30 g/100g to 76.82 g/100g. It was found that the *Varikka* jackfruit seed flour samples had more carbohydrates than the *Koozha* jackfruit seed flour samples. The findings are comparable with the values reported by Eke-ejiofor et al [12] and Ocloo et al. [23], who reported 72.16 g/100 g and 79.34 g/100 g of carbohydrate content in jackfruit seed flour, respectively.

**Table no: 3 Proximate composition of jackfruit seed flour samples**

Treatments	Moisture (%)	Ash (%)	Protein (g/100g)	Fat (%)	Crude fiber (%)	Carbohydrate (g/100g)
------------	--------------	---------	------------------	---------	-----------------	-----------------------

<b>VS<sub>1</sub></b>	7.81	2.61 <sup>a</sup>	13.17 <sup>a</sup>	1.72 <sup>a</sup>	2.96 <sup>a</sup>	76.82 <sup>a</sup>
<b>KS<sub>1</sub></b>	7.79	2.55 <sup>b</sup>	12.58 <sup>b</sup>	1.68 <sup>b</sup>	2.87 <sup>b</sup>	74.30 <sup>b</sup>
<b>±SE(m)</b>	0.005	0.003	0.011	0.005	0.005	0.05
<b>CV%</b>	0.105	0.223	0.142	0.478	0.28	0.114

Values are means of triplicates. Values with different superscripts (a,b,c,d) within the same column are significantly different ( $P \leq 0.05$ ). VS<sub>1</sub>- Varikka soaked in water, KS<sub>1</sub>- Koozha soaked in water.

## CONCLUSION

The purpose of the study was to assess the functional properties, Sensory qualities and proximate composition of jackfruit seed flour from various soaking treatments. This study used gelatinization and tray drying as the primary processing techniques for the production of the jackfruit seed flour. The functional parameters such as WAC, OAC, SP, SI, BD, and dispersibility of the cultivars *Koozha* and *Varikka* jackfruit seed flour were documented. The functional properties of two different cultivars of jackfruit seed flour were found to vary significantly. The cultivar *Varikka* had higher WAC, OAC, SI, SP, BD and dispersibility than cultivar *Koozha*. The statistical analysis of sensory evaluation revealed that the treatments VS<sub>1</sub> and KS<sub>1</sub> had the highest scores for all the attributes, hence these two samples were chosen for proximal analysis. The quality analysis revealed that the jackfruit seed flour is a high-quality source of carbs, protein, and fiber. Jackfruit seed flour had minimal fat content. Reduced fat prevents rancidity and increases flour shelf life. The tray drying process can alter the colour, aroma, and textural properties of the seed samples. This procedure makes it feasible to preserve jackfruit seeds by converting them into flour, extending their shelf life. The findings revealed that jackfruit seed flour can be employed as a key ingredient in a variety of nutrient-dense food products.

## REFERENCE

1. Akter B, Haque MA. Utilization of jackfruit (*Artocarpus heterophyllus*) seed's flour in food processing: A review. *The Agriculturists*. 2018;16(2):131-142.
2. AOAC. Official methods of analysis of association of official analytical chemists. 18th Edition, Washington, DC; 2010.
3. AOAC. Official Methods of Analysis: 18th Edition. Association of Official Analytical Chemists. Washington, DC. 2005;210.
4. Arefin P, Ahmed S, Habib MS, Sadiq ZA, Bobby F, Dey SS, Abdurrahim M, Ashraf T, Islam S, Arefin A, Arefin MS. Assessment and Comparison of Nutritional Properties of Jackfruit Seed Powder with Rice, Wheat, Barley, and Maize Flour. *Current Research in Nutrition and Food Science*. 2022;10(2):544.
5. Arya AM, Singh BR, Samsher SC, Neelesh Chauhan V, Singh S. Functional properties of jackfruit seed flour. *FOOD SCIENCE*. 2020;11(2):115-119.
6. Awuchi CG, Igwe VS, Echeta CK. The functional properties of foods and flours. *International Journal of Advanced Academic Research*. 2019;5(11):139-160.
7. Beuchat LR. Functional and electrophoretic characteristics of succinylated peanut flour protein. *Journal of Agricultural and Food chemistry*. 1977;25(2):258-261.
8. Borgis S, Bharati P. Processing characteristics and acceptability of jackfruit (*Artocarpus heterophyllus* Lam.) seeds, physical and functional properties of its flour. *EPRA International Journal of Research and Development*. 2020;5(10):193-202.
9. Butool S, Butool M. Nutritional quality on value addition to jack fruit seed flour. *Int. J. Sci. Res*. 2015;4(4):2406-2411.
10. Chowdhury AR, Bhattacharyya AK, Chattopadhyay P. Study on functional properties of raw and blended jackfruit seed flour (a non-conventional source) for food application. 2012;3(3):347-353.
11. De Barros SF, Cardoso MA. Adherence to and acceptability of home fortification with vitamins and minerals in children aged 6 to 23 months: a systematic review. *BMC Public Health*. 2016; 16:1-11.
12. Eke-Ejiofor J, Beleya EA, Onyenorah NI. The effect of processing methods on the functional and compositional properties of jackfruit seed flour. *Int. J. Nutr. Food Sci*. 2014;3(3):166-173.
13. Hajj VF, Lopes AP, Visentainer JV, Petenuci ME, Fonseca GG. Physicochemical properties, mineral and fatty acids composition of Jackfruit seeds flour of two varieties from Brazilian Midwest. *Acta Scientiarum. Technology*. 2022;44:e60187.
14. Islam MS, Begum R, Khatun M, Dey KC. A study on nutritional and functional properties analysis of jackfruit seed flour and value addition to biscuits. *Int J Eng Res Technol*. 2015;4(12):139-147.
15. Iwe MO, Onyeukwu U, Agiriga AN. Proximate, functional and pasting properties of FARO 44 rice, African yam bean and brown cowpea seeds composite flour. *Cogent Food & Agriculture*. 2016; 31(1):1142409.

16. Jagadeesh SL, Reddy BS, Basavaraj N, Swamy GS, Gorbal K, Hegde L, Raghavan GS, Kajjidoni ST. Inter tree variability for fruit quality in jackfruit selections of Western Ghats of India. *Scientia Horticulturae*. 2007;112(4):382-387.
17. Juárez-Barrientos JM, Hernández-Santos B, Herman-Lara E, Martínez-Sánchez CE, Torruco-Uco JG, Ramírez-Rivera EJ, Pineda-Pineda JM, Rodríguez-Miranda J. Effects of boiling on the functional, thermal and compositional properties of the Mexican jackfruit (*Artocarpus heterophyllus*) seed jackfruit seed meal properties. *Emirates Journal of Food and Agriculture*. 2017;29(1):1.
18. Kamal MM, Chowdhury MG, Shishir MR, Sabuz AA, Islam MM, Khan MH. Impacts of drying on physicochemical properties, bioactive compounds, antioxidant capacity, and microstructure of jackfruit seed flour. *Biomass Conversion and Biorefinery*. 2023; 21:1-16.
19. Kulkarni KD, Kulkarni DN, Ingle UM. Sorghum malt-based weaning food formulations: preparation, functional properties, and nutritive value. *Food and nutrition bulletin*. 1991;13(4):1-7.
20. Li XE, Jervis SM, Drake MA. Examining extrinsic factors that influence product acceptance: a review. *Journal of food science*. 2015;80(5):R901-909.
21. Nabubuya A, Mugabi R, Kaggwa A, Ainebyona P, Nalugya R. Influence of roasting on the proximate, functional and sensory properties of Jackfruit seeds and amaranth grain composite complementary flours. *Tanzania Journal of Science*. 2022;48(1):156-169.
22. Niba LL, Bokanga MM, Jackson FL, Schlimme DS, Li BW. Physicochemical properties and starch granular characteristics of flour from various *Manihot esculenta* (cassava) genotypes. *Journal of food science*. 2002 ;67(5):1701-1705.
23. Ocloo FC, Bansa D, Boatın R, Adom T, Agbemavor WS. Physico-chemical, functional and pasting characteristics of flour produced from Jackfruits (*Artocarpus heterophyllus*) seeds. *Agriculture and biology journal of North America*. 2010;1(5):903-908.
24. Okaka JC, Potter NN. Physico-chemical and functional properties of cowpea powders processed to reduce beany flavor. *Journal of Food science*. 1979;44(4):1235-1240.
25. Oladele AK, Aina JO. Chemical composition and functional properties of flour produced from two varieties of tigernut (*Cyperus esculentus*). *African journal of biotechnology*. 2007;6(21): 2473-2476.
26. Osungbaro TO, Jimoh D, Osundeyi E. Functional and pasting properties of composite Cassava-Sorghum flour meals. *Agriculture and Biology Journal of North America*. 2010;1(4):715-7120.
27. Palamthodi S, Shimpi S, Tungare K. A study on nutritional composition and functional properties of wheat, ragi and jackfruit seed composite flour. *Food Science and Applied Biotechnology*. 2021; 4(1):63-75.
28. Reddy SS, Devi GN, Lakshmi K, Lakshmi KB. Physico-chemical and functional properties of jackfruit (*Artocarpus heterophyllus*) seed flour. *The Journal of Research ANGRAU*. 2022;50(4):45-56.

29. Romagny S, Ginon E, Salles C. Impact of reducing fat, salt and sugar in commercial foods on consumer acceptability and willingness to pay in real tasting conditions: A home experiment. *Food Quality and Preference*. 2017; 56:164-72.
30. Shedge MS, Haldankar PM, Ahammed Shabeer TP, Pawar CD, Kasture VV, Khandekar RG, Khapare LS. Jackfruit: functional component related with human health and its application in food industry. *Pharma Innov J*. 2022; 11:824-830.
31. Sultana A, Parvin R, Alam MK, Akter F, Alim MA. Physico-chemical, functional properties and storage characteristics of jackfruit seed flour. *Bangladesh Journal of Veterinary and Animal Sciences*. 2015;3(1):20-25.
32. Suresh C, Samsher S. Assessment of functional properties of different flours. *African Journal of Agricultural Research*. 2013;8(38):4849-4852.
33. Vijayaraghavan K, Ahmad D, Ibrahim MK. Biohydrogen generation from jackfruit peel using anaerobic contact filter. *International Journal of Hydrogen Energy*. 2006;31(5):569-579.
34. Waghmare R, Memon N, Gat Y, Gandhi S, Kumar V, Panghal A. Jackfruit seed: an accompaniment to functional foods. *Brazilian Journal of Food Technology*. 2019;22:e2018207.
35. Weyh C, Krüger K, Peeling P, Castell L. The role of minerals in the optimal functioning of the immune system. *Nutrients*. 2022;14(3):644.