

Comparative study of different rapid multiplication methods in black pepper (*Piper nigrum* L.) cv. Panniyur - 1

Abstract

The present investigation entitled, “Comparative study of different rapid multiplication methods in black pepper (*Piper nigrum* L.) cv. Panniyur - 1” was conducted during the year 2023-2024 at the Department of Plantation, Spices, Medicinal and Aromatic Crops, College of Horticulture, Dapoli, Dr. B. S. K. K. V., Dapoli, Dist. Ratnagiri (M.S.), India. The experiment was conducted in randomized block design with seven treatments namely T₁: Raised bed method, T₂: Soil mound method, T₃: Wooden log method, T₄: Serpentine method, T₅: Split halves of PVC method, T₆: Modified serpentine method, T₇: Trench method and each treatment ~~was~~ were replicated four times. Among the various treatments studied, the treatment T₄ (Serpentine method) recorded maximum number of cuttings per harvest, total number of cuttings harvested per year, sprouting percentage at 30 days after planting (DAP), survival percentage at 45 DAP and the highest B:C ratio (2.37) was also observed in serpentine method.

Keywords: Black pepper, rooted cuttings, rapid propagation, multiplication, techniques

Introduction

Black pepper (*Piper nigrum* L.) belongs to the family Piperaceae and is a perennial ~~export~~ export-oriented spice crop in India. It is gaining popularity worldwide as an indispensable food adjunct due to its unique pungency and associated flavour. It is regarded as “King of Spices” and “Black Gold” it having enormous socio-economic importance (Bhat *et al.*, 1995)^[3]. Pepper crop is native to South Asia and historical records reveal that pepper ~~is~~ originated in South India. Black pepper has spicy taste is mainly due to the ~~present~~ presence of a compound piperine ~~that~~ is a pungent alkaloid that enhances the bioavailability of various structurally and therapeutically diverse drugs (Khajuria *et al.*, 2002)^[6]. Black pepper includes 5-9% of piperine, a pungent alkaloid, as its main bioactive chemical compound (Jiang, 2019)^[4]. Piperine comprises of four isomers namely piperine, chavicine, isopiperine, and isochavicine (Ahmad *et al.*, 2012)^[1].

Black pepper ~~has the ability to can~~ reproduce both vegetatively and through seeds. Seed propagation requires a significant amount of time and effort due to ~~their~~ its high sterility and low viability in the post-fertilization stages. They have a modest number of offspring (Ravindran, 2000)^[10]. Furthermore, plants derived from different vines must exhibit variability in growth patterns and productivity (Khan *et al.*, 2021; Philip *et al.*, 1992)^[7] ^[9]. As a result, commercial production is boosted by the adoption of vegetative propagation techniques. Although black

Commented [BK1]: Recheck it

pepper can be propagated through layering, cuttings, rooted cutting, grafting, and budding for industrial cultivation (Thangaselvabal *et al.*, 2008)^[11].

The different rapid multiplication methods such as raised bed method, soil mound method, wooden log method, serpentine method, split halves of PVC method, modified serpentine method, and trench method of black pepper used in [the](#) present study for quicker propagation of black pepper plants, leading to mass multiplication production within a shorter time frame. It can reduce the overall cost of production by minimizing the time and resources required for propagation, such as labour and materials. By rapidly multiplying plants, growers can preserve the genetic diversity of black pepper varieties, ensuring a more resilient and adaptable crop in the face of pests, diseases, and environmental changes. These methods enable growers to scale up their operations more efficiently, meeting the demands of a growing market for black pepper. With rapid multiplications, growers can maintain a steady supply of black pepper throughout the year, reducing seasonal fluctuations in availability and prices. Controlled environments used in rapid multiplication methods can help minimize the risk of disease transmission, ensuring that propagated plants are healthy and disease-free. These rapid multiplication methods allow for the propagation of black pepper in diverse environmental conditions, making cultivation feasible in regions where traditional propagation methods ~~not beare not~~ suitable. Overall, rapid multiplication methods play [a](#) vital role in enhancing the efficiency, sustainability, and profitability of black pepper cultivation.

Material and methods

The field experiment was carried out at the Department of Plantation, Spices, Medicinal and Aromatic Crops, College of Horticulture, Dapoli, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli, Dist. Ratnagiri during the year 2023-2024. The experiment was executed in Randomized Block Design (RBD) with seven treatments and four replications. Five black ~~pepper~~ [pepper](#)-rooted cuttings were selected randomly from [the](#) replication of each treatment to record observations. ~~Average~~ [The average](#) of five cuttings was computed for further statistical analysis. In each treatment, 20 cuttings were planted in every replication. The maximum number of cuttings per harvest, [the](#) total number of cuttings obtained per year, [the](#) sprouting percentage at 30 DAP, and [the](#) survival percentage at 45 DAP were recorded at appropriate stages during [the](#) investigation period. The data was statistically ~~analysed~~ [analyzed](#) as [the](#) method suggested by Panse and Sukhatme (1995)^[9].

Results and Discussion

Number of cuttings per harvest

The data ~~pertaining to about~~ the number of cuttings in different rapid multiplication methods in black pepper are presented in Table 1 and graphically depicted in Fig. 1. From the data, it is observed that there was a significant difference among the treatments ~~with respect to the~~ number of cuttings per harvest.

The number of cuttings obtained at the first and second harvests differed significantly among the various treatments. The highest number of cuttings at the first harvest (55.60) and second (63.37) harvest were observed in treatment T₄ (Serpentine method) which was significantly superior over the rest of the treatments at the first and second harvest. The minimum number of cuttings of at first (30.61) and second (35.94) harvest were recorded in treatment T₇ (Trench method).

In the present experiment, the highest number of cuttings per harvest ~~were was~~ observed in the serpentine method. It might be due to each node ~~is being~~ gently pressed into the polybags with the help of a clip, as the rooted cuttings grow and produce nodes these nodes eventually come in contact with soil and potting media such as vermicompost, allowing for roots at that specific point.

Similar results reported by Kadake (2019)^[5] observed that the serpentine method produced the maximum numbers of black pepper cuttings at first (52.35) and second (52.95) harvest. Whereas, the raised bed method produced the minimum number of cuttings at first (33.15) and second (35.80) harvest, respectively.

Total number of cuttings

The perusal of data on the total number of cuttings obtained in different rapid multiplication methods of black pepper ~~are is~~ presented in Table 1 and depicted in Fig 1.

From the data, it is observed that there was a significant difference among the different treatments ~~in with~~ respect ~~of to the~~ total number of cuttings obtained per year. Significantly the highest numbers of cuttings (118.97) harvested in a year ~~were~~ recorded in treatment T₄ i.e. serpentine method which was significantly superior over the rest of the treatments. However, the lowest total number of cuttings ~~were was~~ recorded in treatment T₇ i.e. Trench method (66.55).

In the present study, the highest total number of cuttings per plant ~~were was~~ obtained in the serpentine method. It might be due to the rapid development of roots at each node when pressed and fixed in (soil + vermicompost) media ~~that which~~ results in a greater number of cuttings every year.

The present findings were similar to the research findings reported by Thapa *et al.* (2017)^[12] stated that the serpentine technique of multiplication produced ~~an~~ an average of 60

cuttings ~~from~~ per mother plant per year. Bhai *et al.* (2018)^[2] concluded that the maximum numbers of black pepper cuttings produced in nine months from a single plant in variety "IISR Malabar Excel" was produced (59.60) cuttings followed by "IISR Thevam" (51.00) and "IISR Shakthi" (23.00) by serpentine method. Kadake (2019)^[5] observed that the serpentine method produced the highest number of cuttings per year (105.30). The lowest number of cuttings ~~were was~~ obtained from ~~the~~ raised bed method (68.95).

Table 1: Effect of different rapid multiplication methods in black pepper on ~~the~~ production of ~~several number of~~ rooted cuttings

| Treatments | Number of cuttings per harvest | | Total number cuttings obtained per year |
|---------------------------------------|---------------------------------|--|---|
| | 1 st harvest 150 DAP | 2 nd harvest 120 days after 1 st harvest | |
| T ₁ - Raised bed (Control) | 39.50 | 46.68 | 86.18 |
| T ₂ - Soil mound | 45.65 | 49.19 | 94.84 |
| T ₃ - Wooden log | 34.53 | 37.40 | 71.93 |
| T ₄ - Serpentine | 55.60 | 63.37 | 118.97 |
| T ₅ - Split halves of PVC | 35.74 | 42.46 | 78.20 |
| T ₆ - Modified serpentine | 50.16 | 52.98 | 103.14 |
| T ₇ - Trench method | 30.61 | 35.94 | 66.55 |
| Mean | 41.68 | 46.86 | 88.54 |
| Range | 30.61-55.60 | 35.94-63.37 | 66.55-118.97 |
| S. Em.± | 1.65 | 2.40 | 5.20 |
| C. D. at 5% | 4.92 | 7.14 | 15.46 |
| 'F' test | SIG | SIG | SIG |

Commented [BK2]: Add a Note about SIG, C.D., S. Em. ±

Commented [BK3]: Remove the value, add the lettering or error bar. If you any help, you will contact with me.




Fig. 1 Effect of different rapid multiplication methods of black pepper on the production of rooted cuttings

Sprouting percentage

The data regarding on sprouting percentage in various rapid multiplication methods in black pepper are presented in Table 2 and graphically represented in Fig 2.

At 30 DAP sprouting percentage at the first and second harvests differed significantly among treatments. Significantly the highest sprouting percentage was recorded in treatment T₄ - Serpentine method at first (93.22 %) and second (97.45 %) harvest which was statistically superior over the rest of treatments. However, the lowest sprouting percentage were-was observed in treatment T₇ - Trench method (63.64 % and 69.70 %) at the first and second harvest, respectively.

In the present investigation, the highest sprouting percentage was observed in the serpentine method. It may be due to the root volume plays-playing an important role in the sprouting of cuttings. Use-The use of soil and vermicompost media resulted in increasing porosity, water holding capacity, low shrinkage, low bulk density, and slow biodegradation of the medium along with the availability of nutrients resulted-resulting in an increase in root volume and dry matter content of root. The weather conditions during the experimental period was-were also congenial for increasing the cell activity for the formation of roots, which resulted in a higher sprouting percentage.


The results are following accordance with the findings of Kadake (2019)^[5] revealed that the highest percentage of sprouting in black pepper cuttings was recorded at first (96.59 %) and

second (94.32 %) harvest in treatment T₄ - Serpentine method. Whereas, the lowest percentage of sprouting was recorded in treatment T₁ - Raised bed method at first (77.73 %) and second (78.37 %) harvest.

Table 2: Effect of different rapid multiplication methods on sprouting percentage per rooted cuttings in black pepper

| Sprouting Percentage (30 DAP) | | |
|---------------------------------------|---------------------------------|--|
| Treatments | 1 st harvest 150 DAP | 2 nd harvest 120 days after 1 st harvest |
| T ₁ - Raised bed (Control) | 76.74 | 81.39 |
| T ₂ - Soil mound | 78.72 | 82.97 |
| T ₃ - Wooden log | 66.19 | 74.64 |
| T ₄ - Serpentine | 93.22 | 97.45 |
| T ₅ - Split halves of PVC | 67.94 | 78.20 |
| T ₆ - Modified serpentine | 82.52 | 84.46 |
| T ₇ - Trench method | 63.64 | 69.70 |
| Mean | 75.56 | 81.25 |
| Range | 63.64-93.22 | 69.70-97.45 |
| S. Em.± | 3.49 | 4.37 |
| C. D. at 5% | 10.38 | 12.98 |
| 'F' test | SIG | SIG |

Commented [BK4]: Add a Note about SIG, C.D., S. Em. ±



Commented [BK5]: Remove the value, add the lettering or error bar. If you any help, you will contact with me.

Fig. 2 Effect of different rapid multiplication methods on sprouting percentage per rooted cuttings of black pepper

Survival percentage

The data ~~pertaining to~~about the survival percentage of rooted cuttings of black pepper in different rapid multiplication methods are presented in Table 3 and graphically represented in Fig. 3.

The highest survival percentage of rooted cuttings at ~~the~~ first harvest (92.37%) and second harvest (96.61%) were observed in treatment T₄ (Serpentine method). Whereas, the lowest survival percentage (60.61%) and (68.18%) was recorded in treatment T₇ (Trench method) at first and second harvest, respectively.

In ~~the~~ present investigation, the serpentine method (T₄) had the highest survival percentage at both ~~the~~ first and second harvest. ~~It is possible that vermicompost supplied physical conditions and adequate nutrients to rooted cuttings, resulting in improved metabolic and physiological activities along with good development of~~ ~~the~~ root system.

The results are in conformity with findings reported by Kadake (2019)^[5] that the maximum survival percentage of black pepper cuttings at first harvest (93.58) and second harvest (91.83) in serpentine method. Thapa *et al.* (2021) and Mustakim *et al.* (2022) ^[13,14] reported that the maximum survivability (93.00 % and 88.33 %, respectively) at 30 and 90 days after cuttings in Panniyur -1 by serpentine layering method; these findings are in agreement with present results.


Commented [BK6]: Recheck

Table 3: Effect of different rapid multiplication methods in black pepper on survival percentage

| Survival Percentage (45 DAP) | | |
|---------------------------------------|---------------------------------|--|
| Treatments | 1 st harvest 150 DAP | 2 nd harvest 120 days after 1 st harvest |
| T ₁ - Raised bed (Control) | 69.76 | 75.58 |
| T ₂ - Soil mound | 73.40 | 77.65 |
| T ₃ - Wooden log | 63.38 | 70.42 |
| T ₄ - Serpentine | 92.37 | 96.61 |
| T ₅ - Split halves of PVC | 65.38 | 71.79 |
| T ₆ - Modified serpentine | 78.64 | 83.49 |
| T ₇ - Trench method | 60.61 | 68.18 |
| Mean | 71.91 | 77.67 |
| Range | 60.61-92.37 | 68.18-96.61 |
| S. Em.± | 3.40 | 4.11 |
| C. D. at 5% | 10.10 | 12.22 |
| 'F' test | SIG | SIG |

Commented [BK7]: Add a Note about SIG, C.D., S. Em. ±

UNDER PEER REVIEW



Commented [BK8]: Remove the value, add the lettering or error bar. If you any help, you will contact with me.

Fig. 3 Effect of different rapid multiplication methods on survival percentage per rooted cuttings in black pepper

Commented [BK9]: Write anything in your paper without box

Conclusion

On the basis of the current study, it can be concluded that; the maximum number of cuttings per harvest, the total number of cuttings obtained per year, the sprouting percentage, and the survival percentage was/were observed in treatment T₄ (Serpentine method). So far as growth performance, quality, and production of the total number of sampling samples per year is are concerned, the serpentine method hence, which was found best for rapid multiplication in black pepper.

References

- Ahmad, N., Fazal, H., Abbasi, B. H., Farooq, S., Ali, M. and Khan, M. A. (2012). Biological role of *Piper nigrum* L. (Black pepper). A review: *Asian Pacific J. Tropical. Biomedicine*. 2(3): 1945-1953.
- Bhai, R. S., Subila, K. P., Eapen, S. J., Reshma, A., Pervez, R., Ishwara Bhat, A. and Srinivasan, V. (2018). Effect of biocontrol agents on production of rooted black pepper cuttings by serpentine method. *J. Spices and Aromatic Crops*, 27(1): 59-65.

3. Bhat, S. R. Chandel, K. P. S. and Malik, S. K. (1995). Plant regeneration from various explants of cultivated Piper species. *Plant Cell Rep.* 14: 398-402.
4. Jiang, T. (2019). Health benefits of culinary herbs and spices. *National Centre for Biotechnology Information*, **102**(2): 395-411.
5. Kadake (2019). Comparative performance of rapid multiplication methods in black pepper (*Piper nigrum* L.) M.Sc. thesis submitted to Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli, Maharashtra. Dist. Ratnagiri (M. S.).
6. Khajuria, A., Thusu, N. and Zutshi, U. (2002). Piperine modulates permeability characteristics of intestine by inducing alteration in membrane dynamics influence on brush border membrane fluidity, ultra-structure and enzyme kinetics. *Phytomedicine*, **9**(3): 224-231.
7. Khan, A. U. Talucder, M. S. A., Das, M., Noreen, S. and Pane, Y. S. (2021). Prospect of the black pepper (*Piper nigrum* L.) as natural product used to an herbal medicine. *Open Access Maced J. Med. Sci.*, **9**(5): 563-573.
8. Panse, V. G. and Sukhatme, P. V. (1995). Statistical methods for agricultural workers, ICAR, New Delhi. Rev. Ed., pp. 97-156.
9. Philip, V. J., Joseph, D., Triggs, G. S. and Dickinson, N. M. (1992). Micropropagation of black pepper (*Piper nigrum* Linn) through shoot tip cultures. *Plant Cell Report*, (12): 41-44.
10. Ravindran, P. N., Nirmal Babu, K., Sasikumar, B. and Krishnamurthy, K. S. (2000). Botany and crop improvement of black pepper, fruit set, fruit development and fruit maturity in Black pepper (*Piper nigrum* L.). India: Hardwood Academy Publishers, p. 87.
11. Thangaselvabal, T., Gailce Leo Justin C. and Leelamathi, M. (2008). Black pepper (*Piper nigrum* L.) 'the king of spices'. A review. *Agricultural Reviews*, **29**(2): 89-98.
12. Thapa, A., Datta, S., Dey, A. N. and Baisare, P. (2017). Advance propagation techniques in important spice crops. *International J. Current Microbiology and Applied Sci.*, **6**(9): 1979-1985.
13. Thapa, A., Rai, U. and Dat, S. (2021). Response of black pepper varieties for propagation by serpentine layering. *J. Spices and Aromatic Crops*, (30).
14. Mustakim, M.T., Talucder, M.S.A., Ruba, U.B., Islam, F., Rahman, A., Uddin, M.S., and Khan, A.U. (2022). Growth Performance of Black Pepper (*P. nigrum*) Cuttings in Different Rooting Media and Growth Regulators. *J. Agrof. Env.* 15(2): 63-68.