

Effect of Soil Application of Organic Amendments on Flowering and yield parameters of chilli (*Capsicum annuum* L.)

ABSTRACT

The present investigation was undertaken to evaluate the effect of soil application of organic amendments on flowering and yield parameters of chilli (*Capsicum annuum* L.). The chilli variety, Vellayani Athulya was selected for the study and the soil was applied with different organic amendments comprising of FYM and FYM substituted with vermicompost, goat manure and combinations of those three with biochar. The results obtained revealed that the treatment of FYM substituted with goat manure (1t ha⁻¹) as per KAU Organic POP(OPOP) (2017)+ rice husk biochar (0.5% w/w) resulted in better flowering and yield attributes in chilli. Also the same treatment showed superior performance in terms of number of flowers plant⁻¹ (44.67), days to first flowering (22), days to first harvest (41), percentage of fruit setting (64.18%), number of fruits plant⁻¹ (28.67), fruit length (15.30cm), fruit girth (9.30cm), fruit weight (17.70 g fruit⁻¹) and yield (507.41 g plant⁻¹). Thus, the combined application of goat manure (1t ha⁻¹) as per KAU OPOP + rice husk biochar (0.5% w/w) proved to be a better soil organic amendment for chilli variety Vellayani Athulya .

Keywords: Soil Application, Organic amendment, Flowering, Yield, Chilli

1. INTRODUCTION

Chilli (*Capsicum annuum* L.) is one of the major spice cum vegetable grown extensively in India. It comes under the family Solanaceae. Solanaceae has major role in balanced diet and the family consists of around 3000–4000 perennial and annual species which comes under 90 genera [1]. Across the world, the key solanaceous members mainly consumed are potatoes, tomatoes, eggplants, and chillies. Overuse of chemical fertilizers results in a number of problems such as air, water, and soil contamination, which is harmful to both human and animal health. But organic manures can replace chemical fertilizers without much reduction in crop yields, if applied effectively. By enhancing the microflora and fauna in the soil, organic manures enhance the quality of the soil and hence increase overall production of crops [2]. Also application of organic manures significantly enhance microbial activities in the soil and improve soil enzyme activities, specifically dehydrogenase and phosphatase, which are crucial for soil health and nutrient cycling [3].

Organic manures such as farm yard manure, sheep manure, pressmud and chicken manure are thought to be storehouses of several nutrients that are necessary for plant growth [4]. Biochar, a carbon-rich product obtained through the pyrolysis of organic matter, has gained attention as a soil amendment due to its potential benefits in plant growth and soil health. Applying biochar increases plant nutrient absorption and enhance physical and chemical characteristics of the soil, including pH, adsorption of plant nutrients, cation exchange capacity (CEC) and water holding capacity. Also, it maintains crop production and improves soil fertility [5]. Organic fertilizers contain some of the nutrients required for better development, and they provide quick growth with higher quality in crops. According to

Commented [W1]: delete

Commented [W2]: check it cm

reports, using vermicompost and chicken manure increases the yield of okra and its contributing characteristics [6]. Goat manure is a valuable organic amendment that can enhance soil fertility and plant growth. The application of goat manure increase the soil organic matter content, which improves soil structure and water-holding capacity and it provides essential nutrients like nitrogen, phosphorus and potassium, supporting plant growth and development. It also enhances soil microbial activity, promoting nutrient cycling and availability of nutrients to plants[7].

2. MATERIAL AND METHODS

2.1 Experimental Setup

A pot culture experiment was carried out at the College of Agriculture, Vellayani from January 2023 to May 2023. Rice husk biochar was prepared at the biochar preparation unit at the department of soil science and agricultural chemistry at College of Agriculture, Vellayani. The chilli variety Vellayani athulya was selected to carry out the experiment and the seedlings were raised in trays and one month old seedlings were transplanted to the pots. Before transplanting, the pots were filled with soil subjected to respective organic treatments as indicated. There were eight treatments comprising of T₁= Control-1 [KAU OPOP (2017) with FYM (25 t ha⁻¹), T₂ = Control -2 [KAU POP (2016)], T₃ = FYM substituted with vermicompost (12.5 t ha⁻¹) as per KAU OPOP (2017), T₄ = FYM substituted with goat manure (1t ha⁻¹) as per KAU OPOP (2017), T₅ = T₁+ Rice husk biochar (0.5% w/w), T₆ = T₂+ Rice husk biochar(0.5% w/w), T₇= T₃+ Rice husk biochar (0.5% w/w) and T₈= T₄+ Rice husk biochar (0.5% w/w). For each treatment, three replications were carried out and the design followed for the experiment was CRD.

OPOP - Basal [Based on soil testing, lime was applied to the soil and phosphorous and potash were supplemented as rock phosphate (217 kg ha⁻¹) and potassium sulphate(62.5 kg ha⁻¹ respectively). The top dressing was done with soil application of groundnut cake (1 kg per 10 litres of water applied at 10 days interval) .It was given for all the treatments except T2. For T2 as per KAU POP, N, P and K were given (N : P₂O₅ : K₂O 75: 40: 25 kg ha⁻¹).

2.2 Flowering parameters

The following flowering parameters were taken at reproductive stage. Representative samples from chilli plants were taken for the observation. Ideally, plants that were uniform in age and health were used for consistency in the data. Tags and markers were used to identify the plants to be observed. This helped in tracking the same plants over time in understanding the flowering stages of the chilli plants. Chilli flowers typically appeared in clusters, and it became important to note the difference between flower buds and fully opened flowers.

2.2.1 Number of flowers plant⁻¹

At the beginning of the flowering period, the number of flower buds and open flowers which appeared on the tagged plants were marked and counted. This was done daily or on every few days, depending on the flowering rate. A record of number of flowers observed on each plant was maintained. A simple table format was used to log the data such as date and the number of flowers. It was continued to observe and count the flowers at regular intervals throughout the flowering period in all the tagged plants of each replication. After the

Commented [W3]: delete

Commented [W4]: delete

Commented [W5]: space adjusted

flowering period, the data was analysed to determine the average number of flowers per plant.

Commented [W6]: space gape

2.2.2 Days to first flowering

The dates when the seeds were sown and the seedlings transplanted were carefully recorded. Thus the starting point for recording the days to first flowering was obtained. The plants were observed regularly for signs of flowering. Flower buds which appeared before the flowers started to open were noted and the first flower opening date was recorded. This was the reference point for calculating the days to first flowering. The observations in each of the plants were recorded to gather the comprehensive data and the data were compiled to analyse the average days to first flowering for the variety under the specific growing conditions.

2.3 Yield parameters

Following yield parameters were taken at harvest stage. A representative sample of chilli plants were used for the observation. Plants that were uniform in age and health were selected to ensure consistency in the data. Tags or markers were used to identify the plants under observation. This helped in tracking the same plants over time. Analysis of various parameters were then carried out.

2.2.2 Days to first harvest

Number of days taken from transplanting to the formation of first fruit was noted from the tagged plants from each replication and the mean value was taken into account.

2.2.2 Percentage of fruit setting

It is the ratio of number of fruits produced to that of total number of flowers produced which gave the fruiting percentage. Observations were taken from the tagged plants from each replication and the mean value was taken into account.

$$\text{Fruit setting (\%)} = \frac{\text{Number of fruits} \times 100}{\text{Number of flowers}}$$

2.2.2 No. of fruits plant⁻¹

The total number of fruits produced by each plant was counted. Observations were taken from tagged plants in each replication and the mean value was taken into account.

2.2.2 Fruit length (cm)

A measuring scale was used to measure the end to end distance and the mean value from each replication was taken into account and it was expressed in cm.

2.2.2 Fruit girth (cm)

A thread was used to measure the fruit girth and the mean value from each replication was taken into account and it was expressed in cm.

2.2.2 Fruit weight (g fruit⁻¹)

An electric weighing balance was used to measure the fruit weight and the mean value from each replication was taken into account and it was expressed in g fruit⁻¹.

2.2.2 Yield (g plant⁻¹)

Weight of all the fruits in individual plants were added to get the total yield per plant, and it was expressed in g plant⁻¹

3. RESULTS AND DISCUSSION

The experimental data was statistically analyzed by completely randomized design using GRAPES 1.0.0 software [8]. Results for flowering and yield parameters are presented in Table 1 to 4 and Figures 1 to 2.

3.1 Number of flowers plant⁻¹

The impact of soil application of different organic amendments on the Number of flowers plant⁻¹ in chilli are illustrated in (Table 1). T8 resulted in highest number of flowers plant⁻¹ (44.67) followed by T7 (41.67) and the lowest value was recorded in the treatment T2(38). T8 (combined application of goat manure and rice husk biochar (0.5% w/w) improved the number of flowers per plant. Biochar enhances physical, chemical and biological properties of soil and there by it enhances nutrient supply in plants [9]. Similarly goat manure also enhances the nutrient availability by means of enhanced microbial activity and there by gives sufficient supply of essential nutrients required for flowering [10]. The result is shown in Table 1.

3.2 Days to first flowering

Table 1 shows the days to first flowering as influenced by soil application of various organic amendments. The lowest value was recorded for the treatment T8 (22) which was found to be on par with T7(23.67). The highest value was recorded for the treatment T2 (30.33). It indicates that T8 [combined application of goat manure and rice husk biochar (0.5% w/w)] resulted in early flowering and it took lesser number of days for the appearance of first flower. In another study application of goat manure (100%), took around 104.13 days only for 50% flowering in chilli [11]. It indicates that flowering occurs more faster by application of goat manure. Also another study has shown that application of 3.2 tons of biochar per acre resulted in highest chilli yield (0.95 tons per acre) compared to control (0.346 tons per acre) [12]. This increased yield could be correlated with flowering as well as improvement in chemical properties of soil.

3.3 Days to first harvest

Table 1. shows the days to first harvest which was found to be influenced by soil application of various organic amendments. The lowest days to first harvest was recorded for the treatment T8 (41.00 days) followed by T3 (44.33 days). The highest days to first harvest was recorded for the treatment T1 (51.00 days) followed by T2 (57.33 days). It indicates that T8 (combined application of goat manure and rice husk biochar (0.5% w/w) resulted in early flowering and it took lesser number of days for the appearance of fruits which resulted in

earlier harvest. Combined application of biochar and goat manure results in enhanced nutrient supply and there by promoting early flowering and fruiting as well as early harvest in chilli [12 & 13].

3.4 Percentage of fruit setting

The impact of soil application of different organic amendments on the percentage of fruit setting in chilli was examined and T8 resulted in highest percentage of fruit setting .Also T8 (64.18%) was followed by T7 (57.46%) and the lowest value was recorded in the treatment T2 (57.46%) followed by T1 (57.89%). It indicates that T8 [combined application of goat manure and rice husk biochar (0.5% w/w)] induced formation of more number of flowers per plant and better development of fruits [14 &15] and resulted in more yield. Thus the percentage of fruit setting was also found to be higher in T8 . The result is shown in Table 1.

Table 1. Effect of soil application of organic amendments on Number of flowers plant⁻¹, Days to first flowering and Days to first harvest and Percentage of fruit setting in chilli

Treatments	Number of flowers plant ⁻¹	Days to first flowering	Days to first harvest	Percentage of fruit setting (%)
T1	39.00	29.33	51.00	57.89
T2	38.00	30.33	57.33	57.46
T3	39.33	28.67	47.33	59.32
T4	38.33	29.00	45.67	58.26
T5	39.00	29.33	47.67	58.97
T6	38.33	26.67	45.67	58.26
T7	41.67	23.67	44.33	58.97
T8	44.67	22.00	41.00	64.18
SE(m) (±)	0.782	0.816	1.08	1.384
C.D(0.05)	2.344	2.448	3.238	4.151

CD=critical difference SE=standard error

3.5 No. of fruits plant⁻¹

Table 2. and Figure 1 shows the impact of soil application of different organic amendments on the number of fruits per plant in chilli. T8 resulted in highest number of fruits per plant (28.67) followed by T7 (25.67cm) and the lowest value was recorded in the treatment T2 (22.33) which was on par with T1 (12.13 g), T4 (22.33) and T6 (22.33). It indicates that T8

[(combined application of goat manure and rice husk biochar (0.5% w/w)] resulted in better fruiting of chilli. Goat manure is one of the important soil amendments which enhance water retention in soil there by enhances nutrient solubility and availability [16]. Biochar enhances soil porosity, organic matter content as well as nutrient availability and thereby it increases fruit setting and yield also [17].

Table 2. Effect of soil application of organic amendments on Number of fruits plant⁻¹ of chilli

Treatments	Number of fruits plant ⁻¹
T1	23.00
T2	22.00
T3	23.33
T4	22.33
T5	23.00
T6	22.33
T7	25.67
T8	28.67
SE(m) (±)	0.782
C.D(0.05)	2.344

CD=critical difference SE=standard error




Figure 1. Effect of soil application of organic amendments on Number of fruits plant⁻¹ of chilli

3.6 Fruit length (cm), Fruit girth(cm), and Fruit weight(g fruit⁻¹)

Table 3. shows the impact of soil application of different organic amendments on yield parameters such as fruit length (cm), fruit girth(cm), and fruit weight(g fruit⁻¹) in chilli. T8 was found as the best treatment for fruit length (cm), fruit girth (cm), and fruit weight(g fruit⁻¹). The fruit length was highest for T8 (15.30cm) followed by T7 (14.10cm) and lowest for T2 (10.83 cm) followed by T1 (11.33cm). Similarly fruit girth was highest for T8 (9.30cm) followed by T7(8.60 cm) and T6 (8.56cm) which were found on par. However the lowest was for T2(7.33 cm) which was on par with T1 (7.50 cm). Fruit weight was highest for T8 (17.70 gfruit⁻¹) followed by T7 (16.20 gfruit⁻¹) and lowest for T2 (11.63 g fruit⁻¹) which was on par with T1 (12.13 gfruit⁻¹). It indicates that T8 (combined application of goat manure and rice husk biochar (0.5% w/w) resulted in better yield parameters in chilli. Biochar contributes to better soil fertility by enhancing soil structure, increasing cation exchange capacity (CEC), and improving moisture retention. These changes create a more favorable environment for root development and nutrient uptake, which are crucial for fruit development [12]. The application of goat manure has been linked to increased fruit length and fruit weight in chili plants. Studies indicate that higher doses of goat manure such as 40 tons per hectare give the best results in terms of fruit dimensions and overall quality [16]

Table 3. Effect of soil application of organic amendments on Fruit length (cm), Fruit girth (cm) and Fruit weight (g) of chilli

Treatments	Fruit length (cm)	Fruit girth (cm)	Fruit weight (g)
T1	11.33	7.50	12.13
T2	10.83	7.33	11.63
T3	11.73	7.87	13.33
T4	12.33	8.07	13.10
T5	13.00	8.57	15.13
T6	11.67	8.17	14.00
T7	14.10	8.60	16.20
T8	15.30	9.30	17.70
SE(m) (±)	0.38	0.12	0.474
C.D(0.05)	1.138	0.36	1.421

CD=critical difference SE=standard error

3.6 Yiled(g plant⁻¹)

Table 4. and Figure 2 shows the impact of soil application of different organic amendments on chilli yield. T8 is reported as the best treatment for yield (507.41 g plant⁻¹). However the lowest yield was obtained in T2 (259.80 g plant⁻¹). It indicates that T8[(combined application of goat manure and rice husk biochar (0.5% w/w)] resulted in better yield, compared to other treatments. Goat manure improves soil structure and nutrient availability, which is crucial for the healthy growth of chili plants. Enhanced soil fertility leads to better nutrient uptake by the plants, which directly correlates with increased yield [14]. The application of biochar has

been shown to enhance soil fertility, which is crucial for plant growth. It improves the physical and chemical properties of soil, leading to better nutrient retention and availability. This results in healthier plants that can produce more fruits [18]

Commented [W7]: Recent reference may include 2020 onwards

Table 4. Effect of soil application of organic amendments on yield of chilli

Treatments	Yield (g plant ⁻¹)
T1	279.06
T2	259.80
T3	311.10
T4	290.33
T5	332.93
T6	322.00
T7	415.81
T8	507.41
SE(m) (±)	0.816
C.D(0.05)	2.448

CD=critical difference SE=standard error




Figure 2. Effect of soil application of organic amendments on yield of chilli

4. CONCLUSION

Organic farming minimizes the use of synthetic fertilizers, reducing the risk of nutrient leaching, greenhouse gas emissions, and water pollution. It promotes the recycling of on-farm resources and making the nutrient cycling system more sustainable. It overcomes the ill effects of indiscriminate use of fertilizers in the environment as well as in soil. Organic matter from sources like manure, compost and plant residues including biochar enhances soil structure, porosity and water-holding capacity. It increases the activity and diversity of beneficial soil microorganisms, promoting nutrient cycling and disease suppression. Organic nutrients are released slowly and steadily, matching the nutrient demands of crops throughout their growth cycle. They improve the availability of essential nutrients like nitrogen (N), phosphorus (P), and potassium (K) for plant uptake. Overall, the use of organic amendments in soil can improve flowering and yield parameters in chilli. The present study revealed that soil application of organic amendments (goat manure along with biochar) had significant effect on flowering and yield parameters of chilli (*Capsicum annuum* L.) and contributed to the sustainable production in chilli.

REFERENCES

1. Knapp, S.; Bohs, L.; Nee, M. and Spooner, D.M. (2004). Solanaceae—a model for linking genomics with biodiversity. *Comparative and Functional Genomics*, 5(3): 285–291
2. Greenspan, A. (2009). Organic alternatives to chemical fertilizer. In *Chemical fertilizer Alternatives*. p. 33-39.
3. Singh, Y.V.; Singh, B.V.; Pabbi, S. and Singh, P.K. (2007). Impact of organic farming on yield and quality of Basmati Rice and soil properties. Paper at: Zwischen Tradition und Globalisierung - 9. Wissenschaftstagung Ökologischer Landbau, Universität Hohenheim, Stuttgart, Deutschland, 20.-23.03.2007
4. Gokul, D., Poonkodi, P., and Angayarkanni, A. 2020. Effect of integrated nutrient management on the growth and nutrient content of chilli (*Capsicum annuum* L.). *IJCS* 8(4): 2647-2651.
5. Alkharabsheh, H.M., Seleiman, M.F., Battaglia, M.L., Shami, A., Jalal, R.S., Alhammad, B.A., Almutairi, K.F. and Al-Saif, A.M., 2021. Biochar and its broad impacts in soil quality and fertility, nutrient leaching and crop productivity: A review. *Agronomy*, 11(5), p.993
6. Sameera, D. L., V. Shankaraiah and D. Srihari, 2005. Effect of packaging and storage on organic manures grown okra (*Abelmoschus esculentus* L. Moench). *Journal of Research, ANGRAU*, 33 (4): 30-35
7. Washaya, S. and Washaya, D.D., 2023. Benefits, concerns and prospects of using goat manure in sub-Saharan Africa. *Pastoralism*, 13(1), p.28.
8. Gopinath P, Parsad R, Joseph B, Adarsh VS. grapesAgri1: Collection of Shiny Apps for Data Analysis in Agriculture. *Journal of Open-Source Software*. 2021; <https://doi.org/6.10.21105/joss.03437>
9. Budiyanto, G. and Achmad, F., 2022, February. Application of Goat Manure Briquettes on Red Chilli Cultivation in Coastal Sandy Soil. In *IOP Conference Series: Earth and Environmental Science* (Vol. 985, No. 1, p. 012030). IOP Publishing
10. Conversa, G., Bonasia, A., Lazzizzera, C. and Elia, A., 2015. Influence of biochar, mycorrhizal inoculation, and fertilizer rate on growth and flowering of Pelargonium (*Pelargonium zonale* L.) plants. *Frontiers in plant science*, 6, p.429.

11. Tripathi, A., Singh, J.K., Mourya, P.K., Chaudhary, P., Chaudhary, A.K. and Kannaujiya, A.K., 2023. Effect of Different Sources of Organic Manures on Growth and Yield of Chilli. *Biological Forum – An International Journal* 15(11): 355-360.
12. Wangmo, T., Dorji, S., Tobgay, T. and Pelden, T., 2022. Effects of biochar on yield of chilli, and soil chemical properties. *Asian Journal of Agricultural Extension, Economics & Sociology*, 40(9), pp.64-77.
13. Uwah, D.F. and Eyo, V.E., 2014. Effects of number and rate of goat manure application on soil properties, growth and yield of sweet maize (*Zea mays L. saccharata* Strut). *Sustainable Agriculture Research*, 3(4).
14. Ojeniyi, S.O., Awodun, M.A. and Akanni, D.A., 2007. Effect of goat manure on some soil properties and growth, yield and nutrient status of tomato. *University of Khartoum Journal of Agricultural Sciences*, 15(3), pp.396-406.
15. Rematwa, A., Situmeang, Y.P. and Suarta, M., 2022. Application of Biochar and Poschar from Several Types of Animal Manure on the Growth and Yield of Red Chili Plants (*Capsicum annum L.*). *Agriwar Journal*, 2(1), pp.1-6
16. Budiyanto, G. and Achmad, F., 2022, February. Application of Goat Manure Briquettes on Red Chili Cultivation in Coastal Sandy Soil. In *IOP Conference Series: Earth and Environmental Science* (Vol. 985, No. 1, p. 012030). IOP Publishing.
17. Hartley, W., Riby, P. and Waterson, J., 2016. Effects of three different biochars on aggregate stability, organic carbon mobility and micronutrient bioavailability. *Journal of Environmental Management*, 181, pp.770-778.
18. HONG, S., LOR, L., NUT, N., THENG, D., VUN, V., UNG, V., HIN, L. and PHUONG, D., 2016. The comparison of liquid bio-slurry and rice husk biochar application on the production yield of daineang chili pepper (*Capsicum annum L.*). *International Journal of Environmental and Rural Development*, 7(1), pp.142-147.

ABBREVIATIONS

OPOP =Package of practices recommendations (Organic), 2017, Kerala Agriculture University
 KAU POP =Package of practices recommendations, 2016, Kerala Agriculture University
 FYM= Farm Yard Manure
 w/w= Weight by weight basis
 CRD=Completely Randomized Design