

## Original Research Article

### Impact of integrated nutrient management on soil nutrient status and yield in garlic (*Allium sativum* L.)

#### ABSTRACT

A two-year experiment was undertaken during rabi seasons of ~~the year~~ 2021 and 2022 to investigate the effect of integrated nutrient management on growth and yield in garlic. The experiment consisted of ~~eighteen~~ treatments with ~~varying combination of inorganic fertilizers along with biofertilizers~~ combinations of inorganic fertilizers, biofertilizers, and organic manures like ~~j~~eevamrit and ~~beejamrit~~Beejamrit. Treatment T<sub>12</sub> [(75 % Recommended dose of NPK (N: P<sub>2</sub>O<sub>5</sub>: K<sub>2</sub>O-125:75:60) + 40 kg S/ha + Azotobacter + PSB + FYM (250 q ha<sup>-1</sup>)] resulted in minimum soil pH (6.82) and electrical conductivity (0.181 dSm<sup>-1</sup>) as well as highest organic carbon (0.86 %), available N (258.84 kg ha<sup>-1</sup>), P (26.72 kg ha<sup>-1</sup>) contents. Further, the treatment T<sub>17</sub> [(100 % Recommended dose of NPK + FYM (250 q ha<sup>-1</sup>)] recorded maximum available K (180.35 kg ha<sup>-1</sup>) and treatment T<sub>16</sub> [(100 % Recommended dose of NPK + 40 kg S/ha + FYM (250 q ha<sup>-1</sup>)] recorded highest S (44.98 kg ha<sup>-1</sup>) contents. The highest gross income (₹ 13,71,900/ha), net income (₹ 10,23,204.48/ha) and B:C ratio (2.93) was recorded by the treatment T<sub>12</sub> [(75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM (250 q ha<sup>-1</sup>)). Thus, it can be concluded that integration of inorganic fertilizers with biofertilizers and organic manures helps in improving the yield and soil nutrient status.

**KEYWORDS:** Biofertilizers; garlic; ~~j~~eevamrit; Jeevamrit; manures; soil fertility

#### INTRODUCTION

Garlic (*Allium sativum* L.) is popularly grown as Allium species after onion belonging to the family Alliaceae. It is an important vegetable crop of India since ancient times. It is originated in Central Asia and used as a vegetable as well as for medicinal purposes. It contains high amount of carbohydrates (29 %), proteins (6.3 %), minerals (0.3 %) and essential oils (0.1- 0.4 %) and contains appreciable quantities of fat, vitamins C and sulphur (Memane et al. 2008). It has antiviral, antibacterial, antifungal and antiprotozoal properties. It is beneficial to the cardiovascular and immune system due to its antioxidant and anti-cancerous properties. It is described as a hot stimulant, carminative and antirheumatic and its oil is a powerful antiseptic mainly due to the presence of allicin an important organosulfur compound. It is used as a

Commented [SS1]:

Commented [SS2]: Restructure sentence

Commented [SS3]:

Commented [SS4]:

Commented [SS5R4]:

Commented [SS6]:

Commented [SS7R6]:

Formatted: Highlight

Commented [SS8]: PSB???

Commented [SS9]: 25 mt/ha :: use metric unit

Formatted: Highlight

Formatted: Highlight

Commented [SS10]: Benefit cost ratio:

Commented [SS11]: Not need to explain since already mentioned in before

Commented [SS12]: Follow citation methods according to journal author guideline

<https://journaljabb.com/index.php/JABB/about/submission>

ey appear in the text. Every reference number in brackets [3].

vermifuge for expelling round worms and has been recommended for cure of several ailments viz., wounds, ulcers, pneumonia, bronchitis and gastro- intestinal disorders (Hazra et al. 2011). Modern agriculture mainly depends upon inorganic fertilizers to fulfil-fulfill the nutrient demand of the crops. Application of fertilizers by farmers without information regarding soil-~~ity and crop information~~ ~~fulfill-fulfill the~~ ~~nutrient~~ ~~demand~~ ~~of~~ ~~the~~ ~~crops~~ ~~Application~~ ~~of~~ ~~fertilizers~~ ~~by~~ ~~farmers~~ ~~without~~ ~~information~~ ~~regarding~~ ~~soil-~~ ~~ity~~ ~~and~~ ~~crop~~ ~~information~~ ~~fulfill-fulfill~~ ~~the~~ ~~nutrient~~ ~~demand~~ ~~of~~ ~~the~~ ~~crops~~ (Ray et al. 2000). Application of synthetic chemical fertilizers alone can have deleterious effects on the soil health, water, and ecosystem. Plant nutrition is one of the key factors influencing growth and yield of crop plants. Nutrients play an important role in internal metabolic activities in plant body. It is well known fact that long time inorganic fertilizers application leads to harmful effect on soil fertility, resulting in poor yield and quality of crops (Meena et al. 2016). Therefore, integration of inorganic fertilizers, organic manures and biofertilizers is capable to maintain the good soil health, productivity and fertility status of soil (Priyanshu et al. 2020).

**Commented [SS13]:** Write concisely

**Commented [SS14]:** Why it is important in garlic production specially in the research site? Provide problems related to garlic production why does it need integration specifically in garlic production???

Is there any nutrient deficiency issues?  
Is there any nutrient leaching issues?  
Is there decline in garlic production?

This question should be answer to show the importance of nutrient integration in garlic production.

Organic manure is an eco-friendly, economically viable and ecologically sound that also played a significant role in improving physical, chemical and biological properties of soil (Acharya and Kumar 2018). It acts as an excellent substrate for soil microbes and increases the proportion of carbon and nitrogen, directly stimulating the population and activity of microorganisms. Jeevamrit is one of the major liquid manures which is prepared from cow urine and dung. Use of Jeevamrit promotes higher growth, yield and quality of crop (Krishna et al. 2017).

**Commented [SS15]:** Provided recent highlights about the role of organic manure in soil health and garlic production

Biofertilizers containing living cells of different types of microorganisms which when applied to seed, plant surface or soil, colonize the rhizosphere or the interior of the plant and promote growth by converting nutritionally important elements like nitrogen and phosphorous from unavailable to available form through biological processes such as nitrogen fixation and solubilization of rock phosphate. It increases the soil fertility, improve soil structure, porosity, water holding capacity and enhance seed germination. Azotobacter is a free living non-symbiotic nitrogen fixing bacteria and it produces auxins, gibberellins, cytokinin and some antibiotic metabolites role for benefit of the plant growth, yield and quality. Phosphate Solubilizing Bacteria solubilizes phosphorus to increase soil fertility and biological activities (Bhushan et al. 2020). Therefore, the importance of the integrated nutrient supply in sustaining productivity is emphasized to restore and sustain soil health and productivity in the long run which otherwise is likely to deteriorate due to continuous and intensive cultivation without adequate nutrient management.

**Commented [SS16]:** Add important role of biofertilizer and type of species use in agriculture, importantly recent highlights related to garlic production and nutrient uptake etc with proper citations.....

## MATERIALS AND METHODS

The experiment was conducted at the Experimental Farm of the Department of Vegetable Science, College of Horticulture & Forestry, Neri, Hamirpur, HP, ~~India during July, 2021 to October, from July 2021 to October 2021 and~~ 2022. Geographically, Neri is located at an altitude of 650 m above mean sea level between 31°41'47.6" N & 72°28'6.3" E. ~~Experimental site is 11 km away from Hamirpur city of Himachal Pradesh.~~ The climate of the region is characterized as subtropical, with hot summers and mild to cool winters. ~~Generally, December and January are coldest months, while May and June are hottest month.~~ Majority of precipitation is received during monsoon period i.e. from June to September. The soil ~~structure~~ of experimental ~~farm site is was~~ to loam to clay loam with pH ranging from 6.8 to 7.0. Before planting of the crop, random soil samples were collected from different spots of the experimental site from a depth of 0-15 cm and the composite sample was prepared. These samples were air-dried, crushed and passed through a 2 mm sieve and stored in cloth bags for chemical analysis of parameters such as soil pH, electrical conductivity, organic carbon and for available nitrogen, available phosphorous and available potassium. The pH and EC of soil samples were measured using a digital pH meter and an electrical conductivity meter, respectively. Organic carbon content of the samples was determined using the Chromic and Titration method proposed by (Walkley and Black 1934). The Alkaline Potassium Permanganate Method was used to determine available N (Subbiah and Asija 1956), ~~phosphorus~~ (P) was measured by the method given by Olsen (Olsen et al. 1954), ~~potassium~~ (K) was measured by Normal Neutral Ammonium Acetate Method (Merwin and Peech 1951) and ~~S~~ was determined by 0.15% CaCl<sub>2</sub> extractant and turbidimetric method (Chesnin and Yien 1950). ~~Mean values of data were subjected to analysis of variance as described by Gomez and Gomez (1984) for Randomized Complete Block Design.~~ The initial value of soil pH (7.03), electrical conductivity (0.248 dSm<sup>-1</sup>), organic carbon (0.64 %), available N (208.43 kg/ha), P (13.52 kg/ha), K (140.24 kg/ha) and S (31.06 kg/ha) contents in soil before the start of the experiment.

### Experimental design and crop management

Experiment was laid out in Randomized Complete Block Design with eighteen treatments and three replications at a spacing of 20 X 10 cm in a plot size of 1X 1m accommodating 50 plants in each plot. The experiment comprised of eighteen treatments viz., T<sub>0</sub> = Control, T<sub>1</sub> = Cow urine + Jeevamrit + FYM (250 q/ha), T<sub>2</sub> = Beejamrit + Jeevamrit + FYM (250 q/ha), T<sub>3</sub> =

**Commented [SS17]:** Not necessary

**Commented [SS18]:** Soil texture

**Commented [SS19]:** Why soil texture is different range?

**Commented [SS20]:** Write full form when you use first time , later you can use short form.

**Commented [SS21]:** Not need to analysis initial sample as anova

**Commented [SS22]:** No fertilizer and organic manures??????

**Commented [SS23]:** Use metric unit

Azotobacter + PSB(Phosphate solubilizing bacteria) + FYM (250 q/ha), T<sub>4</sub> = 75 % Recommended dose of NPK + Azotobacter + FYM (250 q/ha), T<sub>5</sub> = 75 % Recommended dose of NPK + PSB + FYM (250 q/ha), T<sub>6</sub> = 75 % Recommended dose of NPK + Azotobacter + PSB + FYM (250 q/ha), T<sub>7</sub> = 50 % Recommended dose of NPK + Azotobacter + FYM (250 q/ha), T<sub>8</sub> = 50 % Recommended dose of NPK + PSB + FYM (250 q/ha), T<sub>9</sub> = 50 % Recommended dose of NPK + Azotobacter + PSB + FYM (250 q/ha), T<sub>10</sub> = 75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + FYM (250 q/ha), T<sub>11</sub> = 75 % Recommended dose of NPK + 40 kg S/ha + PSB + FYM (250 q/ha), T<sub>12</sub> = 75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM (250 q/ha), T<sub>13</sub> = 50 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + FYM (250 q/ha), T<sub>14</sub> = 50 % Recommended dose of NPK + 40 kg S/ha + PSB + FYM (250 q/ha), T<sub>15</sub> = 50 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM (250 q/ha), T<sub>16</sub> = 100 % Recommended dose of NPK + 40 kg S/ha + FYM (250 q/ha), T<sub>17</sub> = 100 % Recommended dose of NPK (125:75:60 kg/ha) + FYM (250 q/ha).

Sowing was done on 19<sup>th</sup> October 2021. Seeds were sown in the lines at a spacing of 20 X 10 cm. The soil structure of experimental farm is loam to clay loam with pH ranging from 6.8 to 7.0. The experimental field was thoroughly ploughed 1 to 2 times with the help of tractor followed by planking. All the stubble and weeds were removed. Agrifound Parvati variety developed by National Horticulture Research and Development Foundation, Nasik, Maharashtra was used for the present study. This variety [Agrifound Parvati](#) is long day type and bulbs are bigger in size (5 – 6.5 cm diameter) and creamy white colour with pinkish tinge [with maturity duration of 220-240 days. It reaches maturity in 220-240 days. Calculated-The calculated](#) amount of inorganic fertilizers Nitrogen, Phosphorous(P<sub>2</sub>O<sub>5</sub>), Potassium(K<sub>2</sub>O) and Sulphur (125:75:60:40 kg/ha) were applied in the form of urea (203.8 kg/ha), SSP (Single super phosphate) [356.25 kg/ha], MOP (Muriate of potassium) [75 kg/ha] and Bentonite Sulphur (44.45 kg/ha) [were applied](#) in respective treatments before sowing of seed. One-third dose of N along with the fulldoses of P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O were applied as basal dose. Remaining one-third dose of N applied after a month of planting and one-third dose applied after 50 days of planting. Organic manures such as [FYM \(250 q/ha\)](#) were applied during field preparation in the respective treatments. Cloves were treated with cow [urine](#) and [beejamrit-Beejamrit](#) as per treatments and planted. Biofertilizers viz., Azotobacter and Phosphate Solubilizing Bacteria were applied after 15 days of sowing. No fertilizer was applied in the control plot. Cloves were soaked in the solution of 10 % cow urine and 10 % solution of [beejamrit-Beejamrit](#), sole application for one hour before

**Commented [SS24]:** List the treatment details in tabular form to show clearly  
For example:

T0: control (no amendment)  
T1: Organic sources (cow urine, Jeevamrit, FYM)  
T2: Beejamrit + Jeevamrit + FYM  
T3: Biofertilizer (Azob., PSB)+ FYM  
T4: 75% Fert.+ Azo+ FYM  
T5: 75% Fert.+PSB+FYM  
T6: 75% Fert.+ T3  
T7: 50% Fert.+ Azo+ FYM  
T8: 50% Fert.+PSB+FYM  
T9: 50% Fert.+ T3  
T10: T4+ Sulfur  
T11: T5+ Sulfur  
T12: T6+ Sulfur  
T13: T7+ Sulfur  
T14: T8+ Sulfur  
T15: T9+ Sulfur  
T16: 100% ferti + FYM+ sulfur  
T17: 100% ferti + FYM

I think this type of annotation helps to describe the results and better understanding.

**Commented [SS25]:** Texture??

Why texture is different, if you are doing in the same location, soil texture should be same, usually soil texture does not change and has less variability unlike soil pH

sowing as per treatments. Azotobacter @ 200 g/ha, Phosphate Solubilizing Bacteria @ 200 g/ha and mixture of Azotobacter + Phosphate Solubilizing Bacteria @ 200 g/ha were applied along with FYM as per the treatment combination. These biofertilizers were mixed thoroughly with FYM and kept for overnight, and applied the next day as per the different treatment combinations.

### Preparation of ~~jeevamrit~~ Jeevamrit and ~~beejamrit~~ Beejamrit

#### Jeevamrit

Jeevamrit is a mixture of cow dung, cow urine, jaggery, pulse flour and living soil. For preparation of ~~jeevamrit~~ Jeevamrit, required quantities of ingredients were thoroughly mixed in water and allowed to ferment for 7 days. Ingredients were stirred once in ~~morning and once in evening in the morning and once in the evening in a~~ clockwise direction. Jeevamrit @ 10 % drenching was given at fortnight interval in the respective treatments. Table 1 shows the standardized technique for preparation of jeevamrit as suggested by Sreenivasa et al. (2010).

**Table 1** Ingredients used for preparation of Jeevamrit

Ingredient	Quantity
Cow dung	10 kg
Cow urine	10 L
Jaggery	2 kg
Pulse Flour	2 kg
Fertile Soil	1 kg
Water	200 L

#### Beejamrit

Palekar (2007) suggested the standardized technique of preparing beejamrit. This is a mixture of cow dung, cow urine, lime and living soil. For preparation of beejamrit take 5 kg fresh cow dung in cloth & tie it with rope. Arrange to dip this cow dung in the bucket containing 20 litres of water up to 12 hours. In other pot, add 50 gm of lime in one liter of water, let it stable for night. Next morning, squeeze the bundle of cow dung in same water thrice continuously, so that all essence of cow dung will get accumulated in it. Add handful of soil from bund of field in that water and stir well. Lastly add 5 litres of cow urine & lime water and stir well. Beejamrit is ready to use.

**Commented [SS26]:** Describe the measured parameter methodology  
Soil sampling process, depth, soil processing etc

Plant sampling process, measured parameters  
Statistical methodology :

How do you calculate cost of experiment, which parameter did you consider for cost prospect???

## RESULTS AND DISCUSSION

### Physiochemical properties of soil

#### pH of soil

Data presented in Table 2 revealed that maximum soil pH (7.07) was obtained by the treatment T<sub>0</sub> (Control) and minimum soil pH (6.82) was recorded in treatment T<sub>12</sub> (75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM). This might be due to release of organic acids during the process of decomposition may be attributed to decline in soil pH (Kurrey et al. 2018).

**Commented [SS27]:** It might be due to application of sulfur in addition to FYM. Provide how does soil chemistry works on reduction on soil ph with sulfur and FYM application

#### Electrical conductivity of soil (dSm<sup>-1</sup>)

Maximum electrical conductivity (0.244 dSm<sup>-1</sup>) was observed in treatment T<sub>17</sub> (100 % Recommended dose of NPK + FYM). While, minimum electrical conductivity (0.181 dSm<sup>-1</sup>) was recorded in treatment T<sub>12</sub> (75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM).

#### Soil Organic Carbon (%)

The highest soil organic carbon (0.86 %) was observed in treatment T<sub>12</sub> (75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM). Whereas, lowest soil organic carbon (0.62 %) was recorded in treatment T<sub>0</sub> (Control). Organic manures and [biofertilizers application might have created environment favourable for the formation of humic acid which stimulated the activity of soil microorganisms resulting in an increase in biofertilizer application might have created an environment favourable for the formation of humic acid, which stimulated the activity of soil microorganisms, increasing the](#) organic carbon of soil (Kurrey et al. 2018).

**Commented [SS28]:** How do the azotobacter, PSB, and FYM help to increase organic carbon in the soil? Provide more discussion with citation

#### Available Nitrogen (kg/ha)

The data presented in Table 2 revealed that maximum available nitrogen (258.84 kg/ha) was recorded in treatment T<sub>12</sub> (75 % [Recommended](#) dose of NPK + 40 kg S/ha + Azotobacter

**Commented [SS29]:** Why? Better nutrient uptake and better nutrient holding capacity of soil due to PSB, azotobacter and organic manure

+ PSB + FYM) and minimum (197.80 kg/ha) was observed in treatment T<sub>0</sub> (Control). The increase in available nitrogen might be due to direct absorption of nitrogen by the soil which enhanced microbial activity and consequent released to organic complexing substances (Choudhary et al. 2015).

#### **Available Phosphorus (kg/ha)**

Maximum available phosphorus (26.72 kg/ha) was recorded in treatment T<sub>12</sub> (75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM) and minimum (12.70 kg/ha) was observed in treatment T<sub>0</sub> (Control). This might be due to activity of phosphate solubilizing bacteria which resulted into release of organic acids that are responsible for conversion of unavailable P to available P (Singh and Singh 2017).

Commented [SS30]: More discussion

#### **Available Potassium (kg/ha)**

The data presented in Table 2 revealed that the maximum available potassium (180.35 kg/ha) was observed in treatment T<sub>17</sub> (100 % Recommended dose of NPK + FYM) and minimum (138.68 kg/ha) was recorded in treatment T<sub>0</sub> (Control). Greater availability of nutrients from inorganic sources might have increased available K in soil.

Commented [SS31]: More discussion

#### **Available Sulphur (kg/ha)**

Maximum available sulphur (44.98 kg/ha) was observed in treatment T<sub>16</sub> (100 % Recommended dose of NPK + 40 kg S/ha + FYM) and minimum (34.68 kg/ha) was observed in treatment T<sub>0</sub> (Control). The increase in the available sulphur content might be due to the application of sulphur which increased the number of sulphur consuming microorganism and accelerate the conversion of sulphur to SO<sub>4</sub><sup>2-</sup> (Solanki et al. 2020).

#### **Economics of Garlic Cultivation**

Perusal of data given in Table 3, indicated that maximum bulb yield (228.65 q/ha) was recorded in treatment T<sub>12</sub> (75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM) and minimum (90.95 q/ha) was observed in treatment T<sub>0</sub> (Control). Maximum gross income (₹ 13,71,900 /ha) was recorded in treatment T<sub>12</sub> (75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM) and minimum (₹ 5,45,700 /ha) was observed in treatment T<sub>0</sub> (Control). Maximum total cost of cultivation (₹ 3,49,967.40 /ha) was recorded in treatment T<sub>16</sub> (100 % Recommended dose of NPK + 40 kg S/ ha + FYM) and minimum (₹ 2,95,280 /ha) was

observed in treatment T<sub>0</sub> (Control). Net income (₹ 10,23,204.48 /ha) was recorded maximum in treatment T<sub>12</sub> (75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM) and minimum net income (₹ 2,50,420 /ha) was observed in treatment T<sub>0</sub> (Control). Highest B:C ratio (2.93) was recorded in treatment T<sub>12</sub> (75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM) and minimum (0.78) was observed in treatment T<sub>2</sub> (Beejamrit + Jeevamrit + FYM).

**Table 2 Effect of integrated nutrient management on pH, EC (dSm<sup>-1</sup>) and organic carbon (%), available N (kg/ha), P (kg/ha), K (kg/ha) and S (kg/ha) in soil**

Treatment	pH	EC (dSm <sup>-1</sup> )	Organic Carbon (%)	Available N (kg/ha)	Available P (kg/ha)	Available K (kg/ha)	Available S (kg/ha)
T <sub>0</sub>	7.07	0.232	0.62	197.80	12.70	138.68	34.68
T <sub>1</sub>	7.01	0.236	0.68	216.74	15.76	144.00	37.34
T <sub>2</sub>	7.03	0.235	0.67	220.44	15.73	144.74	36.67
T <sub>3</sub>	6.99	0.193	0.73	225.70	17.27	144.20	39.02
T <sub>4</sub>	6.98	0.224	0.78	229.47	19.41	152.00	39.34
T <sub>5</sub>	6.96	0.223	0.75	227.22	21.00	145.68	37.69
T <sub>6</sub>	6.88	0.183	0.82	253.95	24.73	167.77	39.35
T <sub>7</sub>	6.97	0.208	0.74	234.14	19.76	144.07	36.69
T <sub>8</sub>	7.01	0.218	0.75	228.11	19.86	146.00	38.68
T <sub>9</sub>	6.95	0.192	0.81	235.41	21.46	154.68	38.68
T <sub>10</sub>	7.02	0.193	0.80	243.77	20.39	158.83	42.50
T <sub>11</sub>	6.97	0.211	0.79	238.41	20.36	163.74	42.85
T <sub>12</sub>	6.82	0.181	0.86	258.84	26.72	170.00	43.00
T <sub>13</sub>	7.03	0.195	0.77	240.81	22.54	155.70	42.01
T <sub>14</sub>	7.01	0.186	0.76	230.54	21.38	147.37	41.34
T <sub>15</sub>	6.90	0.184	0.83	250.70	22.71	150.68	41.16
T <sub>16</sub>	7.02	0.242	0.72	242.40	22.30	177.71	44.98
T <sub>17</sub>	7.04	0.244	0.70	247.90	22.42	180.35	38.35
Mean	6.98	0.210	0.75	234.57	20.36	154.78	39.68
CD <sub>(0.05)</sub>	0.03	0.004	0.02	3.09	1.25	2.46	1.56
SE(m)	0.01	0.001	0.01	1.07	0.43	0.85	0.54
C.V.	0.33	1.018	2.06	0.79	3.69	0.95	2.3

T<sub>0</sub> = Control, T<sub>1</sub> = Cowurine + Jeevamrit + FYM (250 q/ha), T<sub>2</sub> = Beejamrit + Jeevamrit + FYM (250 q/ha), T<sub>3</sub> = Azotobacter + PSB + FYM (250 q/ha), T<sub>4</sub> = 75 % Recommended dose of NPK + Azotobacter + FYM (250 q/ha),

Formatted: Highlight

T<sub>5</sub> = 75 % Recommended dose of NPK + PSB + FYM (250 q/ha), T<sub>6</sub> = 75 % Recommended dose of NPK + Azotobacter + PSB + FYM (250 q/ha), T<sub>7</sub> = 50 % Recommended dose of NPK + Azotobacter + FYM (250 q/ha), T<sub>8</sub> = 50 % Recommended dose of NPK + PSB + FYM (250 q/ha), T<sub>9</sub> = 50 % Recommended dose of NPK + Azotobacter + PSB + FYM (250 q/ha), T<sub>10</sub> = 75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + FYM (250 q/ha), T<sub>11</sub> = 75 % Recommended dose of NPK + 40 kg S/ha + PSB + FYM (250 q/ha), T<sub>12</sub> = 75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM (250 q/ha), T<sub>13</sub> = 50 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + FYM (250 q/ha), T<sub>14</sub> = 50 % Recommended dose of NPK + 40 kg S/ha + PSB + FYM (250 q/ha), T<sub>15</sub> = 50 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM (250 q/ha), T<sub>16</sub> = 100 % Recommended dose of NPK + 40 kg S/ha + FYM (250 q/ha), T<sub>17</sub> = 100 % Recommended dose of NPK (125:75:60 kg/ha) + FYM (250 q/ha).

**Table 3 Effect of integrated nutrient management on economics of garlic cultivation**

Treatment	Total bulb yield (q/ha)	Total cost of cultivation (₹/ha)	Gross Income(₹/ha)	Net Income (₹/ha)	B:C ratio
T <sub>0</sub>	90.95	2,95,280	5,45,700	2,50,420	0.84
T <sub>1</sub>	136	3,77,080	8,16,000	4,38,920	1.16
T <sub>2</sub>	112.20	3,77,085	6,73,200	2,96,115	0.78
T <sub>3</sub>	137.37	3,41,180	8,24,220	4,83,040	1.41
T <sub>4</sub>	144.50	3,47,295.52	8,67,000	5,19,704.48	1.49
T <sub>5</sub>	146.20	3,47,195.52	8,77,200	5,30,004.48	1.52
T <sub>6</sub>	207.40	3,47,695.52	12,44,400	8,96,704.48	2.57
T <sub>7</sub>	152.15	3,45,123.64	9,12,900	5,67,776.36	1.64
T <sub>8</sub>	146.20	3,45,023.64	8,77,200	5,32,176.36	1.54
T <sub>9</sub>	175.10	3,45,523.64	10,50,600	7,05,076.36	2.04
T <sub>10</sub>	206.55	3,48,295.52	12,39,300	8,91,004.48	2.55
T <sub>11</sub>	172.55	3,48,195.52	10,35,300	6,87,104.48	1.97
T <sub>12</sub>	228.65	3,48,695.52	13,71,900	10,23,204.48	2.93
T <sub>13</sub>	157.25	3,46,123.64	9,43,500	5,97,376.36	1.72
T <sub>14</sub>	168.30	3,46,023.64	10,09,800	6,63,776.36	1.91
T <sub>15</sub>	223.55	3,46,523.64	13,41,300	9,94,776.36	2.87
T <sub>16</sub>	169.15	3,49,967.40	10,14,900	6,64,932.60	1.89
T <sub>17</sub>	152.15	3,48,967.40	9,12,900	5,63,932.60	1.61

T<sub>0</sub> = Control, T<sub>1</sub> = Cowurine + Jeevamrit + FYM (250 q/ha), T<sub>2</sub> = Beejamrit + Jeevamrit + FYM (250 q/ha), T<sub>3</sub> = Azotobacter + PSB + FYM (250 q/ha), T<sub>4</sub> = 75 % Recommended dose of NPK + Azotobacter + FYM (250 q/ha), T<sub>5</sub> = 75 % Recommended dose of NPK + PSB + FYM (250 q/ha), T<sub>6</sub> = 75 % Recommended dose of NPK + Azotobacter + PSB + FYM (250 q/ha), T<sub>7</sub> = 50 % Recommended dose of NPK + Azotobacter + FYM (250 q/ha), T<sub>8</sub> = 50 % Recommended dose of NPK + PSB + FYM (250 q/ha), T<sub>9</sub> = 50 % Recommended dose of NPK + Azotobacter + PSB + FYM (250 q/ha), T<sub>10</sub> = 75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + FYM (250 q/ha), T<sub>11</sub> = 75 % Recommended dose of NPK + 40 kg S/ha + PSB + FYM (250 q/ha), T<sub>12</sub> = 75 %

Formatted: Highlight

Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM (250 q/ha), T<sub>13</sub> = 50 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + FYM (250 q/ha), T<sub>14</sub> = 50 % Recommended dose of NPK + 40 kg S/ha + PSB + FYM (250 q/ha), T<sub>15</sub> = 50 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM (250 q/ha), T<sub>16</sub> = 100 % Recommended dose of NPK + 40 kg S/ha + FYM (250 q/ha), T<sub>17</sub> = 100 % Recommended dose of NPK (125:75:60 kg/ha) + FYM (250 q/ha).

### Conclusion

Based on the results of this study, it can be concluded that overall treatment T<sub>12</sub> (75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM) performed best in terms of yield and nutrient status of soil in garlic. Highest B:C ratio (2.93) was also recorded in treatment T<sub>12</sub> (75 % Recommended dose of NPK + 40 kg S/ha + Azotobacter + PSB + FYM) and minimum (0.78) was observed in treatment T<sub>2</sub> (Beejamrit + Jeevamrit + FYM). Hence, the combined application of organic manures and inorganic fertilizers with biofertilizers helps ~~in improving the yield and physiochemical properties of soil~~ improve soil yield and physiochemical properties.

## References

- Acharya S, and Kumar H. Effect of some organic manure on growth and yield of garlic in greenhouse condition at cold desert high altitude Ladakh region. *Defence Life Science Journal*. 2018;3(2):100-104.
- Bhushan C, Yadav AK, Gangwar HK, Kumar B, Katiyar SK, and Vikram N. Effect of biofertilizers on growth and yield of garlic. *International Journal of Current Microbiology and Applied Sciences*. 2020;9(11):739-743.
- Chesnin L, and Yien CH. Turbidimetric determination of available sulphates. *Soil Sciences Society of America Proceeding*. 1950;15(1):149-151.
- Choudhary MK, Kavita A, Maurya IB, and Hatwal PK. Effect of biofertilizers and micronutrients on yield and quality of garlic var. 'G-282' under black cotton soils. *Green Farming*. 2015;6(4):829-832.
- Gomez KA, and Gomez AA. 1984. *Statistical Procedures for Agricultural Research*. 2<sup>nd</sup> ed. John Wiley & Sons Inc. New York. 427p.
- Hazra P, Chattopadhyay A, and Karmakar K. *Modern technology in vegetable production*. New India Publishing Agency. 2011; pp.288-289.
- Krishna D, Shivaprasad M, Tatagar MH, Kareem MA and Sweta K. Response of garlic for graded level of fertilizers and jeevamrit application. *Research Journal of Chemical and Environmental Science*. 2017;5(6):2321-2340.
- Kurrey DK, Sharma R, Lahre MK and Kurrey RL. Effect of Azotobacter on physio-chemical characteristics of soil in onion field. *The Pharma Innovation Journal*. 2018;7(2):108-113.
- Memane PG, Tomer RS, Kulkarni GU and Chovatia RS. Effect of clove weight and plant growth regulators on growth and yield of garlic cv. Gujrat garlic-3. *The Asian Journal of Horticulture*. 2008;3(1):82-86.
- Meena SL, Bairwa HL, Mahawer LN, Meena S, Chittorz A and Meena S. Effect of integrated nutrient management on growth, yield and quality of onion cv. N-53. *Research on Crops*. 2016;17(3):550-554.

Merwin HD and Peech M. Exchange ability of soil potassium in the sand, silt and clay fraction as influenced by the nature and complementary exchangeable cations. Soil Science American Proceedings. 1951;15(1):125-128.

Olsen SR, Cole CV, Watenable DS and Dean LA. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. USDS Circular. 1954; 939p.

Palekar S. 2007. Zero budget spiritual farming. Research development and extension movement, Amir Subhash Palekar publication, India.100p.

Priyanshu ABM, Singh K, Kumar M, Kumar V, Malik S, Sahahi UP and Lodhi SK. Effect of integrated nutrient management on yield and quality of garlic. Journal of Agrisearch. 2020;7(4):251-254.

Ray PK, Jana AK, Mitra DN, Saha MN, Choudhary J, Saha S and Saha AR. Fertilizer prescription on soil test basis for jute, rice and wheat in typic ustochrept. Journal of the Indian Society of Soil Science. 2000;48(2):79-84.

Singh G and Singh SK. Effect of biofertilizers and NPK on yield of garlic and nutrient availability of soil. Agriways. 2017;5(2):91-96.

Solanki SS, Chaurasiya A, Mudgal A, Mishra A and Singh AK. Effect of soil application of sulphur, farm yard manure and vermicompost on soil fertility, growth and yield of garlic (*Allium sativum* L.). International Journal of Chemical Studies. 2020;8(1):1370-1373.

Sreenivasa MN, Naik N, Bhat SN and Nekar MM. Effect of organic liquid manures on growth, yield and quality of chilli. Green Farming. 2010;1(1):282-284.

Subbiah BV and Asija GL. A rapid procedure for the estimation of the available nitrogen in soils. Current Science. 1956;25(2):259-260.

Walkley A and Black TA. An estimation of soil organic matter and proposed modification of the chromic acid titration method. Soil Science. 1934;37(2):29-38.