

Original Research Article

"Impact of Sowing Methods, Weed Management, and Growth Promoters on the Economics and Nutrient Dynamics of Kharif Maize (*Zea mays* L.)"

Abstract

The present investigation was carried out during two consecutive *kharif* seasons in years 2022 and 2023 at ~~students~~Students, Instructional Farm, Department of Agronomy, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur (U.P). The experiment was laid out in Split Plot Design keeping sowing methods in main plot and weed management practices in sub plots with growth promoters in sub – sub plots with three replications. There ~~was were~~ two sowing methods viz; Conventional methods (S₁) and Ridge methods (S₂). Whereas ~~five~~ weed management practices ~~were five~~, viz; ~~Weed-weed Free-free~~ (W₁), ~~Weedy-weedy Check-check~~ (W₂), Atrazine ~~Prepre~~-emergence @ 1.25 Kg /ha (W₃), Halosulfuron methyl ~~Postpost~~-emergence @ 65g a.i./ha (W₄), Atrazine ~~Prepre~~-emergence @ 0.75 l/ha + Halosulfuron methyl ~~Postpost~~-emergence @ 35g ai./ha (W₄). and ~~there were~~ ~~was~~ ~~Growth-growth~~ promoters' practices ~~were three~~ viz; ~~Gibberellic gibberellic~~ acid (Sayish) (G₁), ~~Amino-amino~~ acid + ~~Humic-humic~~ acid (Spring ~~everEver~~) (G₂), ~~Cytokinins-cytokinins~~ + ~~Enzymes-enzymes~~ (Ambition) (G₃); ~~were there~~. The ~~experiment-crop~~ was shown on 10th July during 2022 and 13rd July during 2023 using hybrid maize DKC-9144. The crop was harvested at full ripe stage on 11th ~~October~~ and 14th October, ~~in during~~ 2022 and 2023, respectively. ~~The best results in economics of various treatments and available nutrients in soil was found under the treatment of S₂ (Ridge method) in sowing method, treatment W₂ (weed free) in weed control methods and treatment G₃ (Cytokinins + Enzymes) in plant growth promoter.~~

Keywords: Humic acid, Cytokinins, Gibberellic acid, Conventional methods, weed management, Economics

1. Introduction

Maize (*Zea mays* L.) is a highly adaptable and versatile crop, known for its exceptional genetic yield potential among food grains. It plays a crucial role in agriculture, serving ~~as~~ both ~~as~~ a staple food for humans and feed for livestock. Often referred to as the "Queen of cereals," maize is integral to numerous industrial applications, including the production of starch, oil, protein, beverages, food sweeteners, pharmaceuticals, cosmetics, textiles, gums, packaging, and paper. Globally, maize is grown on approximately 207.25 million hectares across 160 countries, with a total production of 1217.30 million tons and an average yield of 5.87 metric tons per hectare. In India, maize ranks as the third most important cereal crop after rice and wheat, covering 10.10 million hectares and yielding 33.60 million tons, with an average productivity of 3.33 metric tons per hectare. In Uttar Pradesh, ~~maize-it~~ is cultivated on around 0.73 million hectares, producing approximately 1.78 million tons with a

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Superscript

Formatted: Superscript

Comment [WU1]: Give quantifiable salient results

Comment [WU2]: Very lengthy: reduce by giving only the importance/ need of study

productivity of about 2448 kg per hectare. This productivity is below the national average (Anonymous, 2020), despite the state's significant role in maize cultivation.

The method of sowing plays a crucial role in boosting maize yields. Traditionally, farmers use the broadcast method, which has several drawbacks including uneven seed distribution, inconsistent planting depth, and seeds being scattered and potentially consumed by birds. Improving planting techniques could lead to higher maize productivity, contributing to food and feed self-sufficiency. Research has shown that maize and sorghum planted on slopes can yield 14 to 106 percent and 6 to 59 percent more, respectively, compared to planting on level beds. Additionally, ridge sowing has been found to enhance plant fresh weight, seedling emergence, and ultimately, grain yield. (Bakht *et al.*, 2006). Various planting methods, including broadcasting, drilling, and dibbling, are used on different terrains. Techniques like flat and ridge sowing are particularly effective (Singh *et al.*, 2018). Moreover, appropriate planting methods can improve drainage, helping to reduce flooding during irrigation and heavy pre-monsoon rains. Using methods such as sowing on beds and ridges can help achieve these benefits. Weeds usually reduces crop yield up to 31.5% (22.7% in *Rabi-rabi* and 36.5% in *kharif* 36.5%). But as farmers adopt some kind of weeding on their crop field, a conservative estimate of 10% loss in crop yields may be taken as more realistic, hand weeding is most effective if done in time, though it is costly and time consuming. Occurrence of a variety of weeds especially during *kharif* season makes it cumbersome and uneconomical owing to non availability of labour. Apart from this, labourers are not available for weeding sowing to other agricultural operation going on simultaneously. Additionally, manual method of weed control cannot be put into practice until weeds have achieved certain heights. In *kharif* maize, problem of severe weed infestation level combined with various species of weeds. Almost every type of weeds namely grassy, broad leaved and sedges infest the maize fields. As they compete for moisture, nutrients, space, light, shelter for many diseases and pest ultimately disturbs the growth of the plants, reduce the yield and deteriorates the quality of crop, hence reduce the protein and starch content. (Sharma *et al.*, 2000).

Gibberellins are probably one of the growth regulators that have a significant effect on flowering. Dwarfing depends upon gibberellin deficiency and dwarfing gene effects on gibberellin biosynthesis. Thus, dwarf maize mutants were treated with gibberellic acid, and after hormone treatment, their growth returned to normal. Tall stems also contain more bioactive gibberellin than short stems do. (Naghashzadeha *et al.*, 2009). The method of spraying plant growth regulators (PGRs) is used to promote the formation of maize biomass; therefore, high-quality maize can be produced. By controlling the transmission and metabolism of plant endogenous hormone signals, PGRs can improve plant shape and yield. (Naeem *et al.*, 2012).

The land configuration practices of furrow sowing reduced maximum gross income (Rs. 78923.50 ha), ~~net~~ net income (Rs. 58966.9 ha) and BC ratio (2.96 ha). Ridge bed sowing and BBF sowing, on the other hand, recorded higher gross income (Rs. 77292.50 ha, Rs. 76630.50 ha), net income (Rs. 57337.9 ha, Rs. 56675.9 ha), and BC ratio (2.87, 2.84),

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

Comment [WU3]: reference

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

Comment [WU4]: ????

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

respectively, minimum gross return (Rs. 75301.50 ha), (Rs. 55546.9 ha), and BC ratio (2.81) with flatbed broad casting. Kumar *et al.*, (2018).

Comment [WU5]: Need not to give results in introduction

2. Methods and materials

Experimental site: - A field experiment was conducted during two consecutive *kharif* seasons of 2022 and 2023 at Student's Instructional Farm, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur ~~which was served as the experimental site. Kanpur Nagar is a city in central Uttar Pradesh that is at a height of 125.9 meters above sea level on the alluvial tract of the Gangetic plains. Its coordinates are 25° to 28° North latitude and 79° to 80° East longitude. The semi arid climate and rich alluvial soil characterize this northern zone. About 935 mm of rain falls on the region each year on average.~~ The soil of experiment plot was sandy loam in texture having 0.45% organic carbon, 190.7 kg/ha available N, 11.85 kg/ha available P and 171.16kg/ha K in both the years. ~~To assess the most suitable hybrid maize variety for effect of sowing methods, weed management and growth promoters on weed dynamics, growth, and economics.~~ The experiment was shown on 10th July during 2022 and 13rd July during 2023 using hybrid maize DKC-9144. The crop was harvested at full ripe stage on 11th October and 14th October, in 2022 and 2023, respectively.

Comment [WU6]: Were samples analyzed separately for both the years

Treatment details: - The experiment was laid out in ~~split~~ Split plot-Plot Design (SPD) in 3 factors with 3 replications. **Main plot:** Sowing ~~Method~~method-2 i.e. (S₁) ~~Conventional~~conventional ~~Method~~method, (S₂) Ridge ~~Method~~method. **Sub Plot:** Weed ~~Management~~management- 5. (W₁) Weed ~~Free~~free, (W₂) Weedy ~~Check~~check, (W₃) Atrazine ~~Prepre~~emergence @ 1.25 ~~Kg~~kg/ha, (W₄) Halosulfuron methyl ~~Postpost~~emergence @ 65g a.i./ha, (W₅) Atrazine ~~Prepre~~emergence @ 0.75 l/ha + Halosulfuron methyl ~~Postpost~~emergence @ 35g ai./ha and **Sub-Sub Plot Treatments:-** Growth ~~Promoters~~promoters- 3 (G₁) Gibberellic acid (Sayish), (G₂) Amino acid + Humic acid (Spring ever), (G₃) Cytokinins + ~~Enzymes~~enzymes (Ambition).

Formatted: Superscript

Formatted: Superscript

Formatted: Font: Not Bold

Fertilizer Application: - The recommended dose of fertilizer (NPK: 120:60:40 ~~Kg~~kg ha⁻¹) was applied uniformly in each plot. ~~Nitrogen was applied as treatments through urea, half as basal dose and remaining half at 45 days after sowing. Phosphorus and potassium were applied respectively.~~

Economics: - The gross monetary returns in rupees per hectare were worked out on the basis of maize yield and green biomass yield. The prevailing market price of maize grains and stover were considered.

Cost of cultivation (ha⁻¹): - Cost of cultivation for different treatments were worked out by considering all the expenses incurred in the cultivation of experimental crop and added with variable cost due to treatment.

Gross monetary returns (ha⁻¹): - The gross monetary returns were calculated by considering the prices of maize cobs and stover yield prevailing in the market at the time of harvest. The money value of both grain and stover yield was added together in order to achieve gross monetary return ha⁻¹.

Net monetary returns (ha⁻¹): -The net monetary return was calculated by deducting the cost of cultivation from the gross monetary returns.

Net monetary returns = Gross monetary income - Total cost of cultivation

Benefit: cost ratio (B: C ratio): -The benefit cost ratio was calculated as follows;

$$\text{B: C ratio} = \text{Gross monetary returns (ha}^{-1}\text{)} / \text{Cost of cultivation (ha}^{-1}\text{)}$$

Comment [WU7]: Give details of treatments and observations recorded

Comment [WU8]: Give details of statistical analysis

Available Nutrients

Available nitrogen: - Total nitrogen was determined by alkaline potassium permanganate as advocated by **Subbiah and Asija (1956)**.

Available phosphorus: - Available phosphorous was determined calorimetrically extracting by 0.5 M NaHCO₃ (pH 8.3) extracting as given by **Olsen et al., (1954)**.

Available potassium: - Available potassium was first extracted by using 1 N NH₄OAC (pH 7.0) Morgan's solution and estimated by Flame photometer as described by **Jackson (1967)**.

3. Result and discussion

3.1. Cost of cultivation

The data was depicted in Table no-1. Effect of different sowing methods on cost of cultivation of maize during kharif season recorded of non-significant. At different days after sowing the ridge method recorded highest cost of cultivation in both the year 2022 and 2023 of experiment over the conventional method of sowing. In 2022, the ridge method (S₂) recorded 51475 (Rs. ha⁻¹) of cost of cultivation which is greater than conventional methods (S₁) method of sowing. In 2023 the ridge method recorded 54454 (Rs. ha⁻¹) of cost of cultivation which is also higher than conventional method of sowing. Similarly, in pooled data of both the year the ridge method recorded higher cost of cultivation in comparison of conventional method of sowing which are 55181 (Rs. ha⁻¹).

In a study on weed management practices, the cost of cultivation varied significantly across treatments over two years. In 2022, the weed-free treatment (W₂) had the highest cost at 53,690 Rs ha⁻¹, while the lowest cost was observed in the weedy check (W₁) at 46,556 Rs ha⁻¹. In 2023, the weed-free treatment (W₂) again recorded the highest cost at 56,673 Rs ha⁻¹, with the weedy check (W₁) at 49,533 Rs ha⁻¹. The pooled data confirmed that the weed-free treatment (W₂) had the highest average cost of 55,181 Rs ha⁻¹. Among herbicide treatments, Atrazine pre-emergence at 0.75 l/ha combined with Halosulfuron methyl post-emergence at 35 g ai/ha (W₃) had the next highest costs, reporting 50,852 Rs ha⁻¹ in 2022 and 53,829 Rs ha⁻¹ in 2023, with a pooled average of 52,340 Rs ha⁻¹. These findings underscore the financial implications of different weed management strategies in maize cultivation, highlighting the trade-offs between cost and weed control effectiveness.

Among all the growth promoters applied treatment the treatment Cytokinins + Enzymes (Ambition) (G₃) recorded superior cost of cultivation 50210 (Rs. ha⁻¹) in 2022. And 53187 (Rs. ha⁻¹) in 2023 respectively. The pooled data of cost of cultivation of experiment also showed superior in Cytokinins + Enzymes (Ambition) (G₃) treatment which is 51698 (Rs. ha⁻¹) in comparison of other treatment. While statistically the effect of all the growth promoters on cost of cultivation of maize at days after sowing recorded significant.

The interaction of all effect of sowing methods, weed management and growth promoters on yield to produce non-significant variation in cost of cultivation (primary and secondary) in both year and pooled data of study.

3.2. Gross return (Rs/ha)

The effect of different sowing methods on gross return of maize during kharif season recorded of non-significant (Table 1.). At different days after sowing the ridge method recorded greater gross return in both the year 2022 and 2023 of experiment over the conventional method of sowing. In 2022, the ridge method (S_2) recorded 141107 (Rs/ha) of gross return which is higher than conventional methods (S_1) method of sowing. In 2023 the ridge method recorded 129131 (Rs/ha) gross return which is also higher than conventional method of sowing. Similarly, in pooled data of both the year the ridge method recorded higher gross return in comparison of conventional method of sowing which are 135119 (Rs/ha). **Kumar *et al.* (2018), CRIDA (2010), Meena *et al.* (2015).**

In a study assessing various weed management practices in maize during the kharif season, the weed-free treatment (W_2) consistently achieved the highest gross returns. In 2022, W_2 recorded a gross return of 142,834 Rs/ha, which decreased to 131,224 Rs/ha in 2023, resulting in a pooled average of 137,029 Rs/ha. Among herbicide treatments, Atrazine pre-emergence at 0.75 l/ha combined with Halosulfuron methyl post-emergence at 35 g ai/ha (W_5) showed notable performance, with gross returns of 141,943 Rs/ha in 2022 and 129,681 Rs/ha in 2023, averaging 135,812 Rs/ha. In contrast, the lowest gross returns were recorded for the Atrazine pre-emergence at 1.25 kg/ha (W_3) treatment. These findings highlight the effectiveness of weed-free management and specific herbicide combinations in maximizing gross returns in maize cultivation, emphasizing the importance of selecting appropriate weed management strategies for improved economic outcomes. **Nagdeote *et al.* (2016)**

Among all the growth promoters applied treatment the treatment Cytokinins + Enzymes (Ambition) (G_3) recorded superior gross return of 140587 (Rs/ha) in 2022. Similarly, in 2023 the treatment Cytokinins + Enzymes (Ambition) (G_3) recorded greater gross return of 129513 (Rs/ha) in comparison of rest of the treatment. The pooled data of gross return of experiment also showed superior in Cytokinins + Enzymes (Ambition) (G_3) treatment which is gross return of 135050 (Rs/ha) in comparison of other treatment. The least gross return of maize under among growth promoters' practices during both the year of experiment recorded from Amino acid + Humic acid (Springever) (G_2) treatment. While statistically the effect of all the growth promoters on gross return of maize at days after sowing recorded significant. Further the data also reverent that the gross return was lower during second years of study as compared to first years. **Giannakoula *et al.* (2012)**

The interaction of all effect of sowing methods, weed management and growth promoters on weed dynamics, growth and yield (Appendices) to produce non-significant variation in gross return (primary and secondary) in both year and pooled data of study.

Table No. 1. Effect of sowing methods, weed management and growth promoters on economics return (Rs/ha) of maize during 2022 and 2023

Treatment	Cost of cultivation			Gross return (Rs/ha)		
	2022	2023	Pooled	2022	2023	Pooled
Sowing Method						
S ₁	49710	52726	51218	137641	126994	132317
S ₂	51475	54454	52964	141107	129131	135119
SE(d)	-	-	-	1348.63	623.67	415.39
CD (P= 0.05)	-	-	-	2697.26	1247.34	1812.13
Weed management						
W ₁	46556	49533	48089	131530	120806	126168
W ₂	53690	56673	55181	142834	131224	137029
W ₃	49364	52341	50852	138381	126658	132519
W ₄	50098	53075	51586	193682	127994	160838
W ₅	50852	53829	52340	141943	129681	135812
SE(d)	-	-	-	1061.16	1138.05	1187.87
CD (P= 0.05)	-	-	-	2126.20	2280.30	2380.12
Growth Promoters						
G ₁	50111	53092	51601	139422	127556	133489
G ₂	50015	52992	51503	136613	124819	130716
G ₃	50210	53187	51698	140587	129513	135050
SE(d)	-	-	-	821.97	881.53	920.12
CD (P= 0.05)	-	-	-	1646.96	1766.31	1843.63
CD (p=0.05) for interactions	NS	NS	NS	NS	NS	

Comment [WU10]: Give details

3.3. Net return (Rs/ha)

The effect of different sowing methods on Net return of maize during kharif season recorded of non-significant (Table No. 2). At different days after sowing the ridge method recorded greater Net return in both the year 2022 and 2023 of experiment over the conventional method of sowing. In 2022, the ridge method (S₂) recorded 78655 (Rs/ha) of Net return which is higher than conventional methods (S₁) method of sowing. In 2023 the ridge method recorded 86652 (Rs/ha) Net return which is also higher than conventional method of sowing. Similarly, in pooled data of both the year the ridge method recorded higher Net return in comparison of conventional method of sowing which are 82653 (Rs/ha).

Kumar et al. (2018), Nagdeote et al. (2016), CRIDA (2010), Meena et al. (2015)

In a study on weed management practices in maize during the kharif season, the weed-free treatment (W₂) consistently achieved the highest net returns. In 2022, W₂ recorded a net return of 80,101 Rs/ha, increasing to 88,868 Rs/ha in 2023, resulting in a pooled average of 84,484 Rs/ha. Among herbicide treatments, Atrazine pre-emergence at 0.75 l/ha combined with Halosulfuron methyl post-emergence at 35 g ai/ha (W₅) yielded the next highest net returns, with 79,582 Rs/ha in 2022 and 88,847 Rs/ha in 2023, averaging 84,214 Rs/ha. Conversely, the lowest net returns were recorded for the treatment using Atrazine pre-emergence at 1.25 kg/ha (W₃). These findings indicate that effective weed management practices significantly influence net returns in maize cultivation, highlighting the benefits of both weed-free treatments and specific herbicide combinations. **Duary et al. (2015), Sheela et al. (2016)**

Among all the growth promoters applied treatment the treatment Cytokinins + Enzymes (Ambition) (G₃) recorded superior Net return of 79303 (Rs/ha) in 2022. Similarly, in 2023 the treatment Cytokinins + Enzymes (Ambition) (G₃) recorded greater Net return of 87399 (Rs/ha) in comparison of rest of the treatment. The pooled data of Net return of experiment also showed superior in Cytokinins + Enzymes (Ambition) (G₃) treatment which is Net return of 83351 (Rs/ha) in comparison of other treatment. The least Net return of maize under among growth promoters' practices during both the year of experiment recorded from Amino acid + Humic acid (Springever) (G₂) treatment. While statistically the effect of all the growth promoters on Net return of maize at days after sowing recorded significant. Further the data also revert that the Net return was lower during second years of study as compared to first years. **Giannakoula et al. (2012)**

The interaction of all effect of sowing methods, weed management and growth promoters on weed dynamics, growth and yield (Appendices) to produce non-significant variation in Net return (primary and secondary) in both year and pooled data of study.

3.4. B: C ratio (%)

The effect of different sowing methods on B:C ratio of maize during kharif season recorded of non-significant (Table 2.). At different days after sowing the ridge method recorded greater B:C ratio in both the year 2022 and 2023 of experiment over the conventional method of sowing. In 2022, the ridge method (S₂) recorded 2.54 (%) of B:C

ratio which is higher than conventional methods (S_1) method of sowing. In 2023 the ridge method recorded 2.62 (%) B:C ratio which is also higher than conventional method of sowing. Similarly, in pooled data of both the year the ridge method recorded higher B:C ratio in comparison of conventional method of sowing which are 2.58 (%). **Kumar et al. (2018), Nagdeote et al. (2016), CRIDA (2010), Meena et al. (2015) Swetha et al. (2015)**

In a study evaluating various weed management practices in maize during the kharif season, the weed-free treatment (W_2) consistently achieved the highest benefit-to-cost (B:C) ratio. In 2022, (W_2) recorded a B:C ratio of 2.44, which increased to 2.51 in 2023, resulting in a pooled average of 2.74. Among the herbicide treatments, Atrazine pre-emergence at 0.75 l/ha combined with Halosulfuron methyl post-emergence at 35 g ai/ha (W_5) showed the next best results, with B:C ratios of 2.44 in 2022 and 2.52 in 2023, averaging 2.48. Conversely, the treatment using Atrazine pre-emergence at 1.25 kg/ha (W_3) recorded the lowest B:C ratio across both years. Overall, these findings highlight the effectiveness of the weed-free treatment and specific herbicide combinations in enhancing the economic viability of maize cultivation. **Duary et al. (2015), Sheela et al. (2016). Swetha et al. (2015).**

The treatment using Cytokinins + Enzymes (Ambition) (G_3) achieved the highest benefit-to-cost (B:C) ratio, recording 2.49 in 2022 and 2.57 in 2023, with a pooled average of 2.53. In contrast, the Amino acid + Humic acid (Springever) (G_2) treatment had the lowest B:C ratio across both years. Significant differences in B:C ratios were observed among all growth promoters, with overall ratios lower in the second year compared to the first. **Giannakoula et al. (2012)**

The interaction of all effect of sowing methods, weed management and growth promoters on weed dynamics, growth, and yield (Appendices) to produce non-significant variation in B:C ratio (primary and secondary) in both year and pooled data of study.

Comment [WU11]: Be brief. Thorough improvement in language and presentation is required

3.5. Available nitrogen in soil (kg ha⁻¹)

The effect of different sowing methods on available nitrogen in soil ~~of maize~~ during ~~kharif~~ season recorded of non-significant (Table 3.). At different days after sowing the ridge method recorded greater available nitrogen in soil in both the year 2022 and 2023 of experiment over the conventional method of sowing. In 2022, the ridge method (S₂) recorded 218.99 (kg ha⁻¹) of available nitrogen in soil which is higher than conventional methods (S₁) method of sowing. In 2023 the ridge method recorded 221.41 (kg ha⁻¹) available nitrogen in soil which is also higher than conventional method of sowing. Similarly, in pooled data of both the year the ridge method recorded higher available nitrogen in soil in comparison of conventional method of sowing which are 220.19 (kg ha⁻¹).

In a two-year study on weed management in maize during the kharif season, the weed-free treatment (W₂) consistently showed the highest available nitrogen in soil, recording 219.53 kg ha⁻¹ in 2022 and 222.26 kg ha⁻¹ in 2023, with a pooled average of 220.89 kg ha⁻¹. Among herbicide treatments, Atrazine pre-emergence at 0.75 l/ha combined with Halosulfuron methyl post-emergence at 35 g ai/ha (W₅) also performed well, yielding 216.50 kg ha⁻¹ in 2022 and 219.27 kg ha⁻¹ in 2023, with a pooled average of 217.88 kg ha⁻¹. Conversely, the treatment using Atrazine pre-emergence at 1.25 kg/ha (W₃) resulted in the lowest available nitrogen levels. These results indicate that effective weed management practices significantly influence nitrogen availability in maize cultivation.

The treatment using Cytokinins + Enzymes (Ambition) (G₃) consistently recorded the highest available nitrogen in soil, with 219.78 kg ha⁻¹ in 2022 and 222.61 kg ha⁻¹ in 2023, yielding a pooled average of 221.20 kg ha⁻¹. Conversely, the Amino acid + Humic acid (Springever) (G₂) treatment had the lowest nitrogen levels. Overall, significant differences in nitrogen availability were noted among the growth promoters, with lower levels observed in the second year.

The interaction of all effect of sowing methods, weed management and growth promoters on weed dynamics, growth, and yield (Appendices) to produce non-significant variation in available nitrogen in soil (primary and secondary) in both year and pooled data of study.

3.6. Available phosphorus in soil (kg ha⁻¹)

The effect of different sowing methods on available phosphorus in soil of maize during kharif season recorded of non-significant (Table 3.). At different days after sowing the ridge method recorded greater available phosphorus in soil in both the year 2022 and 2023 of experiment over the conventional method of sowing. In 2022, the ridge method (S₂) recorded 14.91 (kg ha⁻¹) of available phosphorus in soil which is higher than conventional methods (S₁) method of sowing. In 2023 the ridge method recorded 14.31 (kg ha⁻¹) available phosphorus in soil which is also higher than conventional method of sowing. Similarly, in pooled data of both the year the ridge method recorded higher available phosphorus in soil in comparison of conventional method of sowing which are 14.16 (kg ha⁻¹).

Different weed management practices significantly influenced the available phosphorus in soil during the experiment. The weed-free treatment (W₂) consistently showed the highest

Comment [WU12]: ???

Formatted: Font: Italic

phosphorus levels, recording 15.90 kg ha⁻¹ in 2022 and 14.61 kg ha⁻¹ in 2023, leading to a pooled average of 14.45 kg ha⁻¹. Among herbicide treatments, Atrazine Pre-emergence at 0.75 l/ha + Halosulfuron methyl post-emergence at 35 g ai/ha (W₅) had the next highest phosphorus levels, with 14.54 kg ha⁻¹ in 2022 and 14.20 kg ha⁻¹ in 2023, resulting in a pooled average of 14.30 kg ha⁻¹. The least available phosphorus was found in the Atrazine Pre-emergence at 1.25 kg/ha (W₃) treatment. The data indicate that the weed-free treatment was the most effective for maintaining higher available phosphorus levels in maize soil, while W₅ provided a significant alternative among herbicide applications. Overall, the findings highlight the importance of effective weed management practices for optimizing soil nutrient availability.

The treatment of Cytokinins + Enzymes (Ambition) (G₃) consistently recorded the highest available phosphorus in soil, with 14.65 kg ha⁻¹ in 2022 and 14.46 kg ha⁻¹ in 2023, resulting in a pooled average of 14.29 kg ha⁻¹. In contrast, the lowest phosphorus levels were observed in Amino acid + Humic acid (Springever) (G₂). Overall, available phosphorus decreased in the second year.

The interaction of all effect of sowing methods, weed management and growth promoters on weed dynamics, growth, and yield (Appendices) to produce non-significant variation in available phosphorus in soil (primary and secondary) in both year and pooled data of study.

3.7. Available potassium in soil (kg ha⁻¹)

The effect of different sowing methods on available potassium in soil of maize during kharif season recorded of non-significant (Table 3.). At different days after sowing the ridge method recorded greater available potassium in soil in both the year 2022 and 2023 of experiment over the conventional method of sowing. In 2022, the ridge method (S₂) recorded 176.59 (kg ha⁻¹) of available potassium in soil which is higher than conventional methods (S₁) method of sowing. In 2023 the ridge method recorded 177.90 (kg ha⁻¹) available potassium in soil which is also higher than conventional method of sowing. Similarly, in pooled data of both the year the ridge method recorded higher available potassium in soil in comparison of conventional method of sowing which are 177.21 (kg ha⁻¹).

Various weed management practices significantly affected the available potassium in soil during the experiment. The weed-free treatment (W₂) consistently showed the highest potassium levels, recording 177.98 kg ha⁻¹ in 2022 and 179.61 kg ha⁻¹ in 2023, with a pooled average of 178.68 kg ha⁻¹. Among herbicide treatments, Atrazine Pre-emergence at 0.75 l/ha + Halosulfuron methyl post-emergence at 35 g ai/ha (W₅) achieved the next highest potassium levels, with 176.38 kg ha⁻¹ in 2022 and 178.09 kg ha⁻¹ in 2023, resulting in a pooled average of 177.19 kg ha⁻¹. The least available potassium was recorded in the Atrazine Pre-emergence at 1.25 kg/ha (W₃) treatment. Overall, the weed-free treatment was the most effective for maintaining higher available potassium levels in maize soil, while W₅ provided a significant alternative among herbicide applications. These findings highlight the importance of weed management practices on soil nutrient availability.

The treatment of Cytokinins + Enzymes (Ambition) (G₃) consistently recorded the highest available potassium in soil, with values of 177.35 kg ha⁻¹ in 2022 and 179.44 kg ha⁻¹ in 2023, leading to a pooled average of 178.45 kg ha⁻¹. In contrast, Amino acid + Humic acid (Springever) (G₂) showed the lowest levels. Available potassium was lower in the first year compared to the second.

The interaction of all effect of sowing methods, weed management and growth promoters on weed dynamics, growth, and yield (Appendices) to produce non-significant variation in available potassium in soil (primary and secondary) in both year and pooled data of study

UNDER PEER REVIEW

Conclusion

The two years of experimentation in sandy loam soil of central Uttar Pradesh, ridge sowing methods were most effective in minimizing weeds and enhancing economics, resulting in the highest cost of cultivation, gross return, net return, available nitrogen, available phosphorus and available potassium ~~but though recording~~ the lowest B:C ratio. The weed-free treatment (W₂) also significantly reduced weed competition and improved returns. Among herbicides, Atrazine ~~as~~ pre-emergence ~~at @~~ 0.75 l/ha combined with Halosulfuron methyl (post-emergence) (W₅) performed well. Additionally, ~~Cytokinins-cytokinins~~ + ~~Enzymes-enzymes~~ (Ambition) (~~G₃~~) proved effective as a growth promoter, enhancing yield and economic returns.

References

- Anonymous (b) DAC & FW (2019). Agricultural Statistics Division. Estimates of Production of Foodgrains for Directorate of Economics and Statistics; Department of Agriculture, Cooperation and Farmers Welfare; Ministry of Agriculture and Farmers Welfare, Government of India.
- Bakht, J., Ahmad S., Tariq M., Akbar, M and Shafi, M. (2006).** Response of maize to planting methods and fertilizers. *N J Agri Bio Sci* 1(3): 8-14.
- CRIDA. (2010).** Central Research Institute for Dryland Agriculture, Hyderabad, Annual Report.
- Duary, B., Pusal Sharma, P.S. and CharanTeja, K., 2015.** Effect of tank-mix application of tembotrione and atrazine on weed growth and productivity of maize.
- Jackson, W.A. and Volk, R.J., (1968).** Role of potassium in photosynthesis and respiration. *The role of potassium in agriculture*, pp.109-145.
- Kumar, M., (2018).** Halosulfuron Methyl 75% WG (Sempra) – A New Herbicide for the Control of *Cyperus rotundus* in Maize (*Zea mays* L.) Crop in Bihar *Int. J. Curr. Microbiol. App. Sci* (2018) 7(3): 841-846
- Kumar, M., (2018).** Halosulfuron Methyl 75% WG (Sempra) A New Herbicide for the Control of *Cyperus rotundus* in Maize (*Zea mays* L.) Crop in Bihar *Int.J.Curr.Microbiol. App.Sci* (2018) 7(3): 841-846.
- Meena, K.L. Idnani, L.K. Kumar, A., Khanna, M., Shokin, L and Cholhaty. R1 (2015)** Water economization in rabi maize (*Zea mays* L.) to enhance productivity through land configuration and irrigation scheduling in the Indo-Gangetic plains of India. *Journal of Soil and Water Conservation*. 14 (1) 49-55.
- Naeem, S., Duffy, J.E. and Zavaleta, E., (2012).** The functions of biological diversity in an age of extinction. *science*, 336(6087), pp.1401-1406.
- Nagdeote, V.G., Mangala, G., Mhaske, A.R., Balpande, S.S. and Ghodpage, R.M. (2016).** Effect of land configuration, plant population and nitrogen management on

productivity of *sweet corn* in vertisol. *International Journal of Agricultural Sciences*. 8(61): 3428-3433.

Naghashzadeha, M., Rafieeb, M and Khorgamyb, A. (2009). Evaluation of effects of gibberellic acid on maize (*Zea mays* L.) in different planting dates. *Plant Ecophysiology* 3: 159-162

Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of available phosphorus in soils by extraction with NaHCO₃, USDA Cir.939. U.S. Washington; 1954.

Comment [WU13]: Year??

Sharma, Toor. and Sur. (2000). Effect of interculture operation and scheduling of atrazine application on weed control and productivity of maize (*Zea mays* L.) in shiwalik foothills of Punjab. *Indian Journal of Agricultural Science*; 70:757-761

Sheela Barla, S.B., Upasani, R.R., Puran, A.N. and Thakur, R., (2016). Weed management in maize.

Singh, P., Singh, H. Janeja. and Rani, S. (2018). Effect of Sowing Methodology on Diverse Hybrid Maize (*Zea mays* L.) Cultivars in Two Contrasting Environments *International Journal of Current Microbiology and Applied Sciences*.

Subbiah BV, Asija GL. A Rapid Procedure for the Estimation of Available Nitrogen in Soils. *Current Science*. 1956; 25:259- 260.

Comment [WU14]: Year?

Swetha, K., Madhavi, M., Pratibha, G. and Ramprakash, T., (2015). Weed management with new generation herbicides in maize

UNDER PEER REVIEW

UNDER PEER REVIEW

