

Original Research Article

**BIOPHYSICOCHEMICAL PROPERTIES AND IN-VITRO  
ANTIOXIDANT INDICES OF *Cnidoscolus aconitifolius* LEAVES**

---

**ABSTRACT**

*Cnidoscolus aconitifolius* also called Chaya leaf or spinach tree, commonly found in West Africa especially Nigeria, was carefully investigated for its biophysicochemical properties, and in-vitro antioxidant potential. The well dried and pulverized sample was assayed for its flavonoids, phenolics, alkaloids, nitrate, moisture, ash and crude content as well as peroxidase, superoxide dismutase (SOD) and its scavenging activity against 2,2-diphenyl-1-picrylhydrazyl (DPPH) following the standard assay methodologies. The result showed that the sample contains  $79.12 \pm 0.16\%$  moisture content,  $6.61 \pm 0.27\%$  ash and  $13.35 \pm 0.27\%$  crude fiber. The result recorded flavonoid content, at  $132.83 \pm 9.25$  mg/100gQE, alkaloid ( $105.91 \pm 1.52$  mg/100g), phenol ( $351.53 \pm 2.33$  mg/100gCE), and nitrate concentration of  $461.29 \pm 5.83$  mg/100g. The in-vitro antioxidant activity results revealed  $0.442 \pm 0.027$  activity/min/unit/g for peroxidase, and  $0.076 \pm 0.004$  units/g Units/g fresh weight for SOD. Meanwhile, the inhibitory concentration (IC<sub>50</sub>) value of the sample against DPPH activity was 12.95 as against 0.13 recorded by the standard (ascorbic acid), indicating that the sample has a lower DPPH scavenging potential when compared to ascorbic acid. In conclusion, these findings highlight the nutritional richness and antioxidant properties of *Cnidoscolus aconitifolius*, indicating its potential health benefits especially in countering the deleterious actions of reactive nitrogen and oxygen species, however the antioxidant activity should be examined in-vivo.

**Keywords:** Antioxidant, Biophysicochemical, *Cnidoscolus aconitifolius*, Chaya leaf, DPPH, Flavonoid, and Nitrate.

**1. INTRODUCTION**

The bioactive components of plants including vegetables are known to contribute significantly to the nutritional and therapeutic potentials and general wellbeing and survival of humans and animals. According to [1], *Cnidoscolus aconitifolius* (family Euphorbiaceae), commonly called chaya or spinach tree is a leafy perennial shrub native of Yucatan peninsula of Mexico in Central America. In Southwest Nigeria, it is commonly known as 'Efo Jerusalem' or 'Efo lyana Ipaja'; in Niger Delta region (South-South) of Nigeria it is known as 'Hospital Too Far'. It is mostly consumed as vegetable in soups, salads and therapeutically used as anti-diabetic, hepatoprotective, anti-microbial, amongst others [1]. According to [2], it is used to treat rheumatism, gastrointestinal disorders and inflammatory diseases. The therapeutic potentials highlight the ethnobotanical significance of the plant within indigenous cultures, requiring the necessity for scientific investigation to elucidate and validate the bioactive compounds responsible for that effect. Chaya leaves is rich in protein, vitamins, calcium, and iron; and antioxidants [3]. According to [4], its shoot and leaves are used as laxative, diuretics, circulation aid, and lactation stimulants. Researches have shown that the presence of some parameters such as moisture content, ash, crude fiber, and nitrate levels

contributes significantly in the variation of the bioactive and nutritional value of vegetables. The moisture content, indicating the water content in plant materials, impacts the overall quality and shelf life [5]; ash content, indicates the presence of inorganic elements [6]; crude fiber, consisting mainly of cellulose, hemicellulose, and lignin, adds to dietary fiber content, impacting digestive health and nutrient absorption [7]; while nitrate, though present in many leafy greens, can be lethargic due to their potential health implications when consumed in high amounts [8], but has been proven to have cardioprotective effects [9]. Abundant antioxidant compounds like coumarin, flavonoids, phenolics, tannin, anthraquinone, triterpenoid, flavobanin, kaempferol, and cyanogenic glycosides, in addition to nutrients like protein, carbohydrates, fiber, vitamins, calcium, iron, sodium, potassium, magnesium, zinc, and copper in Chaya leaves was reported by [10]. The anti-nutritional contents of Chaya leaves such as oxalate, phytate, hydrogen cyanide, saponin and tannins was reported by [11]. Although some previous studies have reported the nutritional and therapeutic value of *C. aconitifolius*, there are limited scientific information on the specimens harvested at Choba locality of Rivers State, Nigeria. Considering geographical area of previous studies, regional differences may contribute to variations in the plant's overall quality and potential in contributing to human health. Exploring and understanding the biophysicochemical composition and in-vitro antioxidant capacity of plants in their specific region is essential for providing accurate information on dietary planning and nutritional interventions to the local population, health practitioners, and policymakers. This study encompassed a detailed examination of the chemical composition, nutritional content, and in-vitro antioxidant capacity of *Cnidocolus aconitifolius*.

## **2. MATERIAL AND METHODS**

### **2.1 Equipment and Reagents**

The study applied biochemical reagents, chemicals and materials of standard analytical grade. They include phosphate buffer, deionized water, ascorbic acid, bromocresol green (BCG) solution, chloroform, atropine, sodium nitrite, aluminum chloride, sodium hydroxide, Folin-Ciocalteu's reagent (FCR), catechol solution, carbonate buffer, adrenalin, sulphuric acid, acetone, potassium hydroxide, salicylic acid, pyrogallol [0.05 M in 0.1M phosphate buffer (pH 6.5)], H<sub>2</sub>O<sub>2</sub> (1% in 0.1M phosphate buffer, pH 6.5). The equipment used include UV-2450 Spectrophotometer (Schimadzu, Japan), Laboratory Oven or incubator (Genlab, E8A64403), water bath (UT4301E), weighing balance (DS Hortitrade), measuring cylinder, crucibles, pair of tongs, beakers, burette, pipette, Whatman paper, funnel, test tubes, and ceramic mortar.

### **2.2 Collection and Preparation of the Sample Extract**

Fresh sample of *Cnidocolus aconitifolius* was harvested from a vegetable garden at Choba, Rivers State, and was identified at the Department of Plant Science and Biotechnology, University of Port Harcourt, Rivers State, Nigeria. For the extraction, 10 g of the leaves sample was weighed out, and 100 mL of phosphate buffer was added before crushing in a ceramic mortar, after which it was left to macerate for 24 h in a shaker at room temperature. The sample was afterwards filtered with grade 1 Whatman paper (Whatman International Ltd, Maidstone, England) and the extract was further analyzed for phytochemicals (flavonoids, alkaloids and phenols), nitrate, moisture content, ash content, and crude fibre; and antioxidant capacity [2,2-diphenyl-1-picrylhydrazyl (DPPH) scavenging assay, superoxide dismutase (SOD) and peroxidase activity (PD)].

### **2.3 Methods of Analyses**

The alkaloid, flavonoid and phenolic content was assayed using the method described by [12]. Nitrate content was assayed by salicylic acid nitration method as described by [13]. The

nutritional composition was examined using the method described by the Association of Analytical Chemists [14]. PD activity was estimated using the method described by [15]. SOD activity was assayed based on the rapid auto-oxidation of adrenaline in the presence of superoxide anions as described by [16], while the DPPH scavenging activity was determined using the method proposed by [17].

## 2.4 Data analysis

The gathered data was analysed primarily using descriptive statistics to summarize the values obtained from the experimental procedures. The results were presented as means  $\pm$  standard deviation of the results for each of the tests. Meanwhile, the 50 percent inhibitory concentrations of samples and standards were obtained by plotting the percentage inhibitions against the standards and generating the R-square and regression equations from the plots for the necessary calculations.

## 3. RESULTS AND DISCUSSION

### 3.1 RESULTS

The current study investigated the biophysicochemical and some in-vitro antioxidant parameters of *Cnidoscolus aconitifolus* leaves harvested from a vegetable farm in Choba, Rivers State, Nigeria. The assayed biological active compounds include alkaloid, flavonoid, nitrate, and phenol. Also investigated were ash, crude fibre and moisture content. The result presented in Table 1 revealed that *C. aconitifolus* is rich in nitrate ( $461.29 \pm 5.83$  mg/100g), phenol ( $351.53 \pm 2.33$  mg/100g), flavonoid ( $132.83 \pm 9.25$  mg/100g), and alkaloid ( $105.91 \pm 1.52$  mg/100g). The concentration of crude fibre, ash content, and moisture content in *C. aconitifolus* sample were  $13.35 \pm 0.27$  %,  $6.61 \pm 0.27$  %, and  $79.12 \pm 0.16$  % respectively.

**Table 1. Biophysicochemical properties of *Cnidoscolus aconitifolus***

Parameter	Concentration
Flavonoids (mg/100g)	$132.83 \pm 9.25$
Alkaloids (mg/100g)	$105.91 \pm 1.52$
Phenols (mg/100g)	$351.53 \pm 2.33$
Nitrate (mg/100g)	$461.29 \pm 5.83$
Crude fibre (%)	$13.35 \pm 0.27$
Ash (%)	$6.61 \pm 0.27$
Moisture (%)	$79.12 \pm 0.16$

Values represent Mean  $\pm$  Standard deviation; number of test, n = 3.

The result of the in-vitro antioxidant capacity assayed by estimating the activities of some antioxidant enzymes (superoxide dismutase and peroxidase), and the free radical scavenging ability of the sample against 2, 2-diphenyl-1-picrylhydrazyl (DPPH) revealed that the sample recorded SOD activity of  $0.076 \pm 0.004$  Units/g fresh weight, and peroxidase activity of  $0.442 \pm 0.027$  (activity/min/u/g) (Table 2)

**Table 2. Antioxidant composition of *Cnidoscolus aconitifolus***

Antioxidant	Concentration
Superoxide Dismutase (Units/g fresh weight)	0.076 ± 0.004
Peroxidase (Activity/min/u/g)	0.442 ± 0.027

Values represent Mean ± Standard deviation; number of test, n = 3.

The free radical scavenging ability of the extract against DPPH as presented in Table 3 indicated that the sample recorded an inhibitory concentration (IC<sub>50</sub>) value of 12.95, which is higher than 0.13 recorded by the standard (ascorbic acid).

**Table 3. DPPH Assay of *Cnidioscolus aconitifolius* compared with Ascorbic acid**

Sample	Inhibitory concentration (IC <sub>50</sub> )	R <sup>2</sup>	Equation
Ascorbic acid	0.13397	0.8979	y = 23.151 ln(x) + 96.536
<i>T. triangulare</i>	12.95	0.97	y = 19.663ln(x) - 0.3602

Values represent Mean; number of test, n = 3.

### 3.2 DISCUSSION

Vegetables has been known to contribute significantly in the general wellbeing of man through its nutritional and therapeutic potential. There is hardly any balanced diet that is void of vegetable. This signifies the essential role of vegetables in the supply of important nutrients. According to [9], intake of diet enriched in vegetables and fruits has a significant effect on the prophylaxis and treatment of cardiovascular diseases (CVDs). The current study investigated the biophysicochemical and some in-vitro antioxidant parameters of *Cnidioscolus aconitifolius* leaves harvested from a vegetable farm in Choba, Rivers State, Nigeria. The assayed biological active compounds include alkaloid, flavonoid, nitrate, and phenol. Also investigated were ash, crude fibre and moisture content. The result presented in Table 1 revealed that *C. aconitifolius* is rich in nitrate, phenol, flavonoid, and alkaloid. The result also revealed significant concentrations of ash content and crude fibre, with a very high moisture content. These compounds are known to for their well-established biological properties and activities. For instance, the ability of nitrate to act as an antioxidant and anti-inflammatory compound was reported by [18]. [9] reported that most biologically active phytochemicals including nitrate present in green leafy and root vegetables may confer cardioprotective benefits via various mechanisms. According to [18], nitrate intake has some beneficial effects on the cardiovascular system such as triglyceride reduction, blood pressure regulation, and stroke and atherosclerosis prevention. Nitrate has also been reported by [19], to act as an anti-nutrient which have direct and indirect effects ranging from mild reactions to death. Remarkably, [20], posited that the ecological risks associated with high nitrate intake signifies that high nitrate content in vegetable is beneficial due to high content of natural antioxidants such as polyphenols, betalain pigments, and vitamins that inhibits nitrosamines synthesis. According to [21], flavonoids are a group of natural substances with variable phenolic structures and are present mostly in fruits, vegetables, grains, bark, roots, stems, flowers, tea and wines. Considering its anti-oxidative, anti-

inflammatory, anti-mutagenic and anti-carcinogenic properties coupled with their capacity to modulate key cellular enzyme function, they have become an essential constituent in various industries such as cosmetics, nutraceutical, pharmaceutical, and medicinal firms. According to [22], phenolic compounds are bioactive constituents of plant that possess good health-promoting activities. The phenolic compounds possess anti-inflammatory, anti-clotting, antioxidant, immune enhancers and hormone modulators [23]. [24], reported the antioxidant properties of phenols. Alkaloid on its part has been reported by [25], to act as CNS stimulant, topical anaesthetic in ophthalmology, powerful pain relievers, antipyretic action, among other uses. The high concentration of phenols, flavonoids, alkaloid, and nitrate recorded in this study is a clear indication that *C. aconitifolus* possess various medicinal effect especially antioxidant potential. Also, the appreciable amount of crude fibre present in the sample is an indication that *C. aconitifolus* has the ability to enhance reduction of cholesterol level, maintain body weight, stabilize glucose and minimize the risk of other health problems such as heart disease, hemorrhoids and cancer as well as help to keep the digestive system healthy by adding bulk to faeces and preventing constipation. The aforementioned activities of crude fibre has been reported by [26]. The ash content recorded in this study is an indication that the *C. aconitifolus* sample contains appreciable amount of inorganic mineral. [27], posited that the ash content of food samples can affect different characteristics of food including physiochemical and nutritional properties. This ensures the safety of foods, making sure there are no toxic minerals present. The result also revealed high moisture content in the sample indicating short shelf-life and poor resistance to bacterial activities which can lead to easy perishability of the sample. Also, high moisture content is an indication of significant amount of micronutrients such as carotene, vitamins, and minerals in leafy vegetables, which makes them an essential component of a nutritious diet [28]. The result of the in-vitro antioxidant capacity assayed by estimating the activities of some antioxidant enzymes (superoxide dismutase and peroxidase), and the free radical scavenging ability of the sample against 2, 2-diphenyl-1-picrylhydrazyl (DPPH) revealed that the sample recorded low SOD activity, and significant peroxidase activity (Table 2). This outcome is in line with the position of [29] that vegetables are rich sources of antioxidant enzymes such as SOD, peroxidase, catalase (CAT), and some other antioxidant compounds that ameliorate the oxidative damage caused by accumulation of reactive oxygen species (ROS). [30], reported that SOD activity regulates the concentration of  $H_2O_2$  and  $O_2$  and serves as the initial line of defense against toxicity caused by different stresses in plants. The SOD activity recorded in this study is an indication that *C. aconitifolus* has a potential to mitigate the activities of  $H_2O_2$  and  $O_2$ . The study also recorded a high peroxidase activity indicating a rapid degradation of hydrogen peroxide ( $H_2O_2$ ) by peroxidase through oxidative reaction. The free radical scavenging ability of the extract against DPPH as presented in Table 3 indicated that the sample recorded a higher inhibitory concentration ( $IC_{50}$ ) value than the standard (ascorbic acid). The higher  $IC_{50}$  value recorded by *C. aconitifolus* in this study is an indication that the plant sample has a lesser potent DPPH radical scavenging activity than the ascorbic acid standard. According to [31], an  $IC_{50}$  value is the concentration of the sample required to scavenge 50% of the free radicals present in the system. It is inversely related to the antioxidant activity of crude extracts. Thus, the lower the  $IC_{50}$  value, the higher the antioxidant activity. DPPH radical scavenging activity remains one of the most vital method applied in assaying the antioxidant activity of plant extract [32]. The antioxidant capacity shown by the sample can be attributed to the presence of flavonoids and other phenolic compounds. The results of the obtained in this study almost correlates with some other outcome of the same assay on *C. aconitifolus* sourced in other areas, as geographical location and soil content can impact the primary data.

#### 4. CONCLUSION

In conclusion, findings from this study has revealed the phytochemical, nutritional value, and antioxidant capacity of *C. aconitifolius* leaves harvested from Choba, Rivers State, Nigeria. The sample has high concentration of nitrate, phenol, flavonoids and alkaloid as well as a very high moisture content and a good amount of crude fibre. The sample also showed a relatively high peroxidase and DPPH radical scavenging activities, though not to be compared with that of ascorbic acid. This indicates the sample's capacity to combat oxidative stress and ensure protection against free radicals. It is imperative to state that *C. aconitifolius* leaves harvested from the study location should be considered a good source of antioxidant, and further studies on its in-vivo antioxidant capacity should be investigated to ascertain its potential as a therapeutic remedy for free radical induced disorders. The nutritional and health benefits of *C. aconitifolius* leaves from the study area will go a long way in addressing certain diet-related ailments. Thus, the populace of the research area should be sensitized on the need to grow more of this vegetable and adopt it as a valuable dietary component.

#### REFERENCES

1. Panghal A, Shaji AO, Nain K, Garg MK, Chhikara N. *Cnidoscopus aconitifolius*: Nutritional, phytochemical composition and health benefits—A review. *Bioactive Compounds in Health and Disease-Online* ISSN: 2574-0334; Print ISSN: 2769-2426. 2021;4(11):260-86.
2. Kutshik RJ, Jiyil MK, Mafuyai CE, Ichor LL. Anti-anaemic effect of *Cnidoscopus chayamansa* (Mc Vaugh) leaf extract on phenyl hydrazine-treated rats. *Journal of Advances in Biology & Biotechnology*. 2020;23(11):23-32.
3. Obichi EA, Monago CC, Belonwu DC. Effect of *Cnidoscopus aconitifolius* (Family Euphorbiaceae) aqueous leaf extract on some antioxidant enzymes and haematological parameters of high fat diet and Streptozotocin induced diabetic wistar albino rats. *Journal of Applied Sciences and Environmental Management*. 2015;19(2):201-9.
4. Olatinwo AO, Monsurat Oyeniye Yakubu RN RM. Effect of *Cnidoscopus aconitifolius* (Miller) IM Johnston leaf extract on reproductive hormones of female rats. *International Journal of Reproductive BioMedicine*. 2007;5(3):149.
5. Piergiovanni L, Limbo S. 17 packaging and the shelf life of vegetable oils. *Food packaging and shelf life*. 2010:317.
6. Ukom AN, Obi JA. Comparative evaluation of the nutrient composition and phytochemical content of selected vegetables consumed in Nigeria. *International Letters of Natural Sciences*. 2018;71.
7. Adeyeye EI, Oyeyemi SD. Nutritional composition of *Talinum triangulare*. *Food Science & Nutrition*. 2017;5(3), 678–683.
8. Brkić D, Bošnjir J, Bevardi M, Bošković AG, Miloš S, Lasić D, Krivohlavek A, Racz A, Mojsović-Čuić A, Trstenjak NU. Nitrate in leafy green vegetables and estimated intake. *African Journal of Traditional, Complementary and Alternative Medicines*. 2017;14(3):31-41.
9. Olas B. The cardioprotective role of nitrate-rich vegetables. *Foods*. 2024;13(5):691.
10. Kuri-García A, Guzmán SH. Phenolic profile and antioxidant capacity of *Cnidoscopus chayamansa* and *Cnidoscopus aconitifolius*: A review. *Journal of Medicinal Plants Research*. 2017;11(45):713-27.
11. Adamu HM, Bara PG, Ushie OA. The proximate composition and anti-nutritive content of *Cnidoscopus aconitifolius* leaves. *International Journal of Biochemistry Research & Review*. 2020;29(8):65-73.

12. Madhu M, Sailaja V, Satyadev TN, Satyanarayana MV. Quantitative phytochemical analysis of selected medicinal plant species by using various organic solvents. *Journal of Pharmacognosy and Phytochemistry*. 2016;5(2):25-9.
13. Cataldo DA, Maroon M, Schrader LE, Youngs VL. Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. *Communications in soil science and plant analysis*. 1975;6(1):71-80.
14. Acland J. Association of Official Analytical Chemist AOAC-1990. Official Method of Analysis. *East Africa Crops*. 1985;15:33-8.
15. Reddy KP, Subhani SM, Khan PA, Kumar KB. Effect of light and benzyladenine on dark-treated growing rice (*Oryza sativa*) leaves II. Changes in peroxidase activity. *Plant and cell physiology*. 1985;26(6):987-94.
16. Misra HP, Fridovich I. The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *Journal of Biological chemistry*. 1972;247(10):3170-5.
17. Katalinic V, Milos M, Kulisic T, Jukic M. Screening of 70 medicinal plant extracts for antioxidant capacity and total phenols. *Food chemistry*. 2006;94(4):550-7.
18. Karwowska M, Kononiuk A. Nitrates/nitrites in food—Risk for nitrosative stress and benefits. *Antioxidants*. 2020;9(3):241.
19. Shi M, Gu J, Wu H, Rauf A, Emran TB, Khan Z, Mitra S, Aljohani AS, Alhumaydhi FA, Al-Awthan YS, Bahattab O. Phytochemicals, nutrition, metabolism, bioavailability, and health benefits in lettuce—A comprehensive review. *Antioxidants*. 2022;11(6):1158.
20. Chen L, Zhu Y, Hu Z, Wu S, Jin C. Beetroot as a functional food with huge health benefits: Antioxidant, antitumor, physical function, and chronic metabolomics activity. *Food science & nutrition*. 2021;9(11):6406-20.
21. Panche AN, Diwan AD, Chandra SR. Flavonoids: an overview. *Journal of nutritional science*. 2016;5:e47.
22. Liao DY, Chai YC, Wang SH, Chen CW, Tsai MS. Antioxidant activities and contents of flavonoids and phenolic acids of *Talinum triangulare* extracts and their immunomodulatory effects. *Journal of food and drug analysis*. 2015;23(2):294-302.
23. Doherty VF, Olaniran OO, Kanife UC. Antimicrobial activities of *Aframomum melegueta* (Alligator pepper). *International journal of Biology*. 2010;2(2):126-31.
24. Foti MC. Antioxidant properties of phenols. *Journal of Pharmacy and Pharmacology*. 2007;59(12):1673-85.
25. Chen C, Lin L. Alkaloids in diet. In *Handbook of Dietary Phytochemicals 2021* (pp. 1595-1629). Singapore: Springer Singapore.
26. Thilagavathi T, Pandiyan M, Suganyadevi M, Sivaji M, Yuvaraj M, Sasmitha R. Dietary fibre-health benefits. *Biotica Research Today*. 2020;2(6):519-22.
27. Thibault JF, Lahaye M, Guillon F. Physico-chemical properties of food plant cell walls. In *Dietary fibre—A component of food: Nutritional function in health and disease 1992* (pp. 21-39). London: Springer London.
28. Schönfeldt HC, Pretorius B. The nutrient content of five traditional South African dark green leafy vegetables—A preliminary study. *Journal of food composition and analysis*. 2011;24(8):1141-6.
29. Tao H, Zhao Y, Li L, He Y, Zhang X, Zhu Y, Hong G. Comparative metabolomics of flavonoids in twenty vegetables reveal their nutritional diversity and potential health benefits. *Food Research International*. 2023;164:112384.
30. Dumanović J, Nepovimova E, Natić M, Kuča K, Jačević V. The significance of reactive oxygen species and antioxidant defense system in plants: A concise overview. *Frontiers in plant science*. 2021;11:552969.
31. Babu D, Gurumurthy P, Borra SK, Cherian KM. Antioxidant and free radical scavenging activity of triphala determined by using different in vitro models. *Journal of Medicinal Plants Research*. 2013;7(39):2898-905.
32. Gulcin İ, Alwasel SH. DPPH radical scavenging assay. *Processes*. 2023;11(8):2248.

UNDER PEER REVIEW