

## Effects of seed size on germination, seedling vigour and other seed quality characters of fodder cowpea (*Vigna unguiculata* (L.) Walp) varieties

### Abstract

A study was conducted to investigate how seed size affects germination and vigour of cowpea seeds. The differences due to vigour parameters of graded seeds were significant due to varieties cv. Bundel Lobia-1 showed better vigour than cv. UPC-607. Variety Bundel Lobia-1 recorded significantly higher seed weight (102.6 g) over variety UPC-607 (92.1 g). There was significant difference between seed grades, large seeds having significantly more 1000 seed weight (118.3 g) followed by ungraded seeds (107.2 g), medium seeds (97.1 g) and small seeds (66.8 g), respectively. Interaction Cv- Bundel Lobia-1 (93.0%) recorded significantly superior germination percentage over cv. UPC-607 (86.48 %). However, the differences were non-significant due to seed sizes, large seed (91.03%) recorded more germination percentage followed by medium seeds (90.34%), ungraded seeds (89.44%), small seed (88.16%) respectively.

**Key words:** Seed size, germination, vigour.

### INTRODUCTION

Cowpea (*Vigna unguiculata* (L.) Walp) is well known multipurpose leguminous crop. It is classified under other pulse crop. It is quick growing bulky leguminous crop and high protein content. There is difference of opinion regarding valid name of species. It is synonymously named as *Vigna sinensis* (L.) Savi and *Vigna catjana* (Burn) Walp. Verdcord and Royal botanical gardens prefers *Vigna unguiculata* (L.) Walp with several cowpea throughout world as cultivars. Its common name is 'cowpea'. Occasionally 'Southern pea' and new english name is 'Covalence'. In vernaculars it is called as 'Chavali'. Cowpea for fodder is often grown during Kharif and summer seasons, either as a sole crop or mixed crop in maize or sorghum. Most of fields after harvest of the rabi crop, remain fallow during summer season and are sown only when monsoon set in. This period from February to May is characterized by bright sunshine, high temperature, low relative humidity and thus keeps down the infestation of insects and diseases at the lowest and thus is very much favourable for cowpea crop.

The forage crops are generally shy seed producers and forage cowpea varieties are not exception for this. However, their seed requirement is high. It is therefore, necessary to find out different ways and means to maximize the seed production of this crop. The quality seed is the cheapest input in modern agriculture. Availability of viable and vigorous seed at planting time is very important for achieving target of forage production as it acts as a catalyst for realizing the potential of other inputs. Seed size is observed to contribute increased seed production. In cowpea, seeds are 4 to 8 mm long, 3-4 mm broad and variable in size and colour. The seed indicates the amount of reserve food supply for seedling. Small and shrivelled seeds do not contain as much food to give the plant a vigorous start as the bold and plump seed (Bremner et.al., 1963).

### MATERIALS AND METHODS

Formatted: Font: (Default) Times New Roman, 12 pt

Commented [AS1]: Scientific nomenclature should always be italicized

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, 12 pt

Commented [AS2]: What is the significance of the doing this?

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Commented [AS3]: I would put at least 5 key words and refrain from using the ones that are already in the title page

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

Formatted: Font: (Default) Times New Roman, 12 pt

Commented [AS4]: The notation for genus name can be used as V. henceforth

Commented [AS5R4]: And all scientific name should be italicized

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

The laboratory experiment was conducted in a factorial completely randomized design with three replications in laboratory of Seed Science and Technology Research Unit, Institute of Agricultural Sciences, Bundelkhand University, Jhansi. The data obtained were subjected to statistical analysis of respective designs.

### Treatments

The treatment details are as follows.

Varieties	a) Bundel Lobia -1	: V <sub>1</sub>
	b) UPC 607	: V <sub>2</sub>
Seed sizes		
V <sub>1</sub>	a) Ungraded	: A
	b) Large	: B
	c) Medium	: C
	d) Small	: D
V <sub>2</sub>	a) Ungraded	: A
	b) Large	: B
	c) Medium	: C
	d) Small	: D

### Thousands seed weight (g)

The weight of three replications of 1000 seeds of each grade from both the varieties was recorded on top pan balance with 0.1 g accuracy.

### Germination percentage, root shoot length and vigour index

Four replications of 100 seeds of different grades were germinated at 30°C temperature for eight days using between paper method (Anonymous, 1985). The seedlings were sorted into normal and abnormal seedlings and germination percentage was expressed on the basis of normal seedling only.

Ten normal seedlings from each of treatments in each replication were randomly selected and the shoot and root lengths were measured in centimeters. The average of ten seedlings was calculated and recorded. Vigour index was calculated by using following formula (Abdul baki *et.al.*, 1973):

$$\text{Vigour Index} = (\text{Root length} + \text{Shoot length}) \times \text{germination percentage}$$



**Fig 1. Germination stage of cowpea plants**      **Fig 2. Root and shoot length of cowpea plants**

#### **Fresh and dry weight of seedling (g)**

The fresh as well as oven dry weight of ten randomly selected eight days old normal seedlings were recorded. The average of ten seedlings was calculated for both, fresh and dry weight.

#### **Moisture content of seeds**

Moisture percentage was determined by grinding seeds in a grinding mill and drying the 5 g ground sample at a constant temperature of 130<sup>o</sup>C for one hour (Anonymous, 1985). The percent moisture content was calculated on net weight basis.

### **RESULTS AND DISCUSSION**

#### **Effect of seed size on thousand seed weight**

The variation in 1000- seed weight due to seed sizes was significant. Large seeds had significantly more thousands seed weight (118.3 g) over other grades (107.2 g to 66.8 g). These results were in agreement with those reported by salih (1982) in Fababean, Vadivelu and Ramkrishnan (1983) in bengal gram and Jha *et.al.*(1985) in wheat.

#### **Effect of seed size on germination percentage**

Varieties had significant effect on this trait and cv. Bundel Lobia-1 (93.0%) had more germination percentage than cv. UPC-607 (86.48%).

On contradictory to the report of Abdullahi and Vanderlip (1972) in sorghum, Randhawa *et.al.* (1973) in Kalyansona wheat, Bhor *et.al.* (1987) in gram. Randhawa *et.al.* (1990) in maize reported that seed size did not improve field emergence and germination percentage.

**Table 1 : Effect of seed size on thousand seed weight (g) and germination percentage**

Treatments	Thousand seed weight (g)	Germination percentage
<b>Variety</b>		
V <sub>1</sub>	102.6	93.0
V <sub>2</sub>	92.1	86.48
S.E.±	1.165	0.573
C.D. at 5%	3.494	1.718
<b>Seed size</b>		
A (Ungraded)	107.2	89.44
B (Large)	118.3	91.03
C (Medium)	97.1	90.34
D (Small)	66.8	88.16
S.E.±	1.648	0.810
C.D. at 5%		
<b>Interaction</b>		
(Variety X Seed size)		
S.E.±	2.331	0.115
C.D. at 5%	6.988	N.S.

**Interaction effect due to varieties and seed sizes for thousand seed weight (g).**

	A	B	C	D	Mean	S.E.±	C.D. at 5%
V <sub>1</sub>	111.8	119.0	106.0	73.6	102.6	2.331	6.988
V <sub>2</sub>	102.6	117.6	88.2	60.0	92.1		
Mean	107.2	118.3	97.1	66.8			
S.E.±					2.331		
C.D. at 5%				6.988			

**Effect of seed size on vigour index**

In present investigation vigour index significantly differ due to varieties. But no influence of seed size observed on seedling vigour index. These findings are in conformity with those reported by Marcos *et.al.* (1977). However, China and Phul (1982) in pearl millert and Kalakannawar *et.al.* (1989) in wheat reported that large sized seeds produce more vigorous seedlings. On contrary, Mugnisjah and Nakamura (1986) in soybean reported that plants from small seed produce more vigorous growth.

In present investigation, medium sized seeds produced more vigour index (3044.33) than all other grades.

**Table 2: Effect of seed size on root shoot length and vigour index**

Treatments	Root length (cm)	Shoot length (cm)	Vigour index
<b>Variety</b>			
V <sub>1</sub>	13.10	19.95	3156.75
V <sub>2</sub>	12.01	19.77	2690.17
S.E. at 5%	0.560	0.454	90.275
C.D. at 5%	1.676	N.S.	270.66
<b>Seed size</b>			
A (Ungraded)	12.08	19.33	2696.5
B (Large)	14.12	19.20	3030.17
C (Medium)	12.29	21.35	3044.33
D (Small)	13.52	19.58	2922.83
S.E.±	0.791	0.642	127.67
C.D. at 5%	N.S.	N.S.	N.S.
<b>Interaction</b>			
(Variety X Seed size)			
S.E.±	1.119	0.908	180.55
C.D. at 5%	N.S.	N.S.	N.S.

**Effect of seed size on fresh and dry weight of seedlings**

The fresh and dry weight of seedlings due to varieties was non-significant. Fresh weight of seedlings due to seed sizes showed significant differences as dry weight of seedling showed non-significant differences (Table 3).

Singh et.al. (1972) in soybean, Bishnoi (1977) in triticale, Evans and Bhatt (1977) in wheat, Vadivelu and Ramakrishnan (1983) in gram reported that larger seeds produced seedlings with higher fresh and dry weight.

### Effect of seed size on moisture content of seed

Non-significant differences were obtained due to varieties, seed sizes and their interactions for moisture content of seed. The moisture content of cv. Bundel Lobia-1 was more than that of cv. UPC-607. Large seeds showed highest moisture content (13.07%) followed by medium seeds (12.95%), ungraded seeds (12.90%) and small seeds (12.29%) respectively.

**Table 3. Effect of seed size on moisture content of seed**

Treatments	Moisture content (%)
<b>Variety</b>	
V <sub>1</sub>	13.37
V <sub>2</sub>	12.23
S.E.±	0.433
C.D. at 5%	N.S.
<b>Seed size</b>	
A (Ungraded)	12.90
B (Large)	13.07
C (Medium)	12.95
D (Small)	12.29
S.E.±	0.613
C.D. at 5%	N.S.
<b>Interaction</b> (Variety X Seed size)	
S.E.±	0.866
C.D. at 5%	N.S.

The difference due to variety x seed size interaction were non-significant for all traits except thousands seed weight.

### CONCLUSION

The result indicated that large seeds do not have advantages over small seeds in terms of germination. Therefore, sorting it out will be of no economic return but time consuming. Large seeds size showed better vigour in seedling dry weight.

However, the difference due to variety x seed size interaction were non-significant for all traits except thousands seed weight. Seed size is not therefore important in germination of cowpea seed, sorting it out will be of no economic return but time consuming.

## REFERENCES

Abdual-Baki. A., James. D, and Anderson. D. 1973. Vigour determination in soybean seed by multiple eriteris. *Crop Sei.* 13 (6) : 630-633.

Abdullahi and Vanderlip, R.L. 1972. Relationship of vigour tests and seed source and size to sorghum seedling establishment. *Agron. j.* 64 :143-144.

Anonymous. 1985. international rules for seed testing and Annexes *seed Sei. and Tech.* 13 (2) ; 356-313.

Bhor, S.R. 1987. Effect of seed size on growth, yield and yield contributing characters in gram (*Cicer arietinum L.*). M.Sc. (Agri.) Thesis, M.P.K.V.. Rahuri (MS.) India

Bishnoi, U.R. and Sapra, V.T. 1975. Effect of seed size on seedling growth and yield performance in hexaploid triticale. *Cereal Research Communication.* 3(1) : 49-60.

Bremner, P.M., Eckersall, R.N. and Scott. R.K, 1963. The relative importance of embryo size and endosperm size in causing the effect associated with seed size in wheat, *J. agrie. Sci.* 61(1) : 139-145.

China, B.S. and Phul, P.S. 1982. Association of seed size and seedling vigour with various morphological traits in pearl millet. *Seed Sci. and Tech.* 10(3) ; 541-545.

Evans, L.E. and Bhatt, G.M. 1977. Influence of seed size protein content and cultivar on early seedling vigour of wheat. *Canadian J. PlantSci.* 57(3) : 929-935.

Jha, B.N., Sinha, S.K. and Singh, J.N 1985. Effect of seed Size on yield in wheat (*Triticum aestivum*). *Seed Res.* 13(1) : 24-27.

Kalakannavar, RM-, Shashidhara, S.A. and Kulkarni, G.N. 1989. Effect of grading on quality of wheat seeds. *Seed Res.* 17(2) : 182-185.

Marcos, F.J., Silva, A.F. , Da, Cicero, S.M. and Goncalves, C.A.R. 1977, Effect of seed size on germination, vigour and yield of maize (*Zea mays L*) *Luiz de queiroz.* 34 : .327-337. (Seed Abstr. 1981. 3708).

Mugnisjah, W.Q. and Nakamura, S. 1988. Vigour of soybean seed as influenced by sowing and harvest dates and Seed Size. *Seed Sci. and Tech.* 14: 87-94.

Randhawa, H.S., Dey, S.K., Kaur, J., Sharma, H.L., Hari Singh and Khehra, A.S. 1990. Studies on seed germination, seedling vigour and seed mycoflora of graded maize (*Zea mays L.*) *Seed Res.* 6(1) : 49-52.

Salih, F.A. 1982. Influence of seed size and sowing date on yield components of *Fababenn.* *Fabis News Letter.* 6(4) : 38-39. (From Seed Abstr, 6:1-12) pp. 161.

Singh, J.N., Tripathi, S.K. and Nagi, P.S. 1972. Note on the effect of seed size on germination, growth and yield of soybean, [*Glycine max (L.) Merr.*]. *The Indian J agric. Sci.* 42(1) : 83- 86.

Vadivelu, K.K. and Ramakrishnan, V. 1983. Effect of seed size on quality attributes and yield of seed in Bengal gram. (*Cicer arietinum L.* ). *Seed Res.* 11(2) : 177-181.

UNDER PEER REVIEW

