

Impact of Seed Size on Germination, Seedling Vigor, and Quality Traits in Fodder Cowpea (*Vigna unguiculata* (L.) Walp) Varieties

Abstract

A study was conducted to investigate how seed size affects germination and vigor of cowpea seeds. The differences due to vigour parameters of graded seeds were significant due to varieties cv. Bundel Lobia-1 showed better vigour than cv. UPC-607. Variety Bundel Lobia-1 recorded significantly higher seed weight (102.6 g) than variety UPC-607 (92.1 g). There was a significant difference between seed grades, large seeds having significantly more than 1000 seed weight (118.3 g) followed by ungraded seeds (107.2 g), medium seeds (97.1 g) and small seeds (66.8 g), respectively. Interaction Cv- Bundel Lobia-1 (93.0%) recorded a significantly superior germination percentage over cv. UPC-607 (86.48 %). However, the differences were non-significant due to seed sizes, large seeds (91.03%) recorded more germination percentage followed by medium seeds (90.34%), ungraded seeds (89.44%) and small seeds (88.16%) respectively.

Key words: Seed size, Germination, Vigor, seed grades

INTRODUCTION

Cowpea (*Vigna unguiculata* (L.) Walp) is a well-known multipurpose leguminous crop. It is classified under other pulse crop. It is a quick growing bulky leguminous crop and has a high protein content. There is a difference of opinion regarding valid name of species. It is synonymously named as *Vigna sinensis* (L.) Savi and *Vigna catjana* (Burn) Walp. Rerecord and Royal botanical gardens prefers *Vigna unguiculata* (L.) Walp with several cowpea throughout world as cultivars. Its common name is 'cowpea'. Occasionally 'Southern pea' and new English name is 'Covalence'. In vernaculars it is called as 'Chavali'. Cowpea for fodder is often grown during Kharif and summer seasons, either as a sole crop or mixed crop in maize or sorghum. Most fields after harvest of the rabi crop, remain fallow during summer season and are sown only when typhoon sets in. This period from February to May is characterized by bright sunshine, high temperature, low relative humidity and thus keeps down the infestation of insects and diseases at the lowest and thus is very much favourable for cowpea crop.

The forage crops are generally shy seed producers and forage cowpea varieties are not exception for this. However, their seed requirement is high. It is therefore, necessary to find out different ways and means to maximize the seed production of this crop. The quality seed is the cheapest input in modern agriculture. Availability of viable and vigorous seed at planting time is very important for achieving target of forage production as it acts as a catalyst for realizing the potential of other inputs. Seed size is observed to contribute increased seed production. In cowpea, seeds are 4 to 8 mm long, 3-4 mm broad and variable in size and colour. The seed indicates the amount of reserve food supply for seedling. Small and shrivelled seeds do not contain as much food to give the plant a vigorous start as the bold and plump seed (Bremner et.al., 1963).

MATERIALS AND METHODS

The laboratory experiment was conducted in a factorial completely randomized design with three replications in laboratory of Seed Science and Technology Research Unit,

Institute of Agricultural Sciences, Bundelkhand University, Jhansi. The data obtained were subjected to statistical analysis of respective designs.

Treatments

List 1: The treatment details are as follows.

Varieties	a)Bundel Lobia -1	:V ₁
	b) UPC 607	: V ₂
Seed sizes		
V ₁	a)Ungraded	: A
	b)Large	: B
	c)Medium	: C
	d)Small	: D
V ₂	a)Ungraded	: A
	b)Large	:B
	c)Medium	: C
	d)Small	: D

Thousands seed weight (g)

The weight of three replications of 1000 seeds of each grade from both the varieties was recorded on top pan balance with 0.1 g accuracy.

Germination percentage, root shoot length and vigour index

Four replications of 100 seeds of different grades were germinated at 30⁰C temperature for eight days using between paper method (Anonymous, 1985). The seedlings were sorted into normal and abnormal seedlings and germination percentage was expressed on the basis of normal seedling only.

Ten normal seedlings from each of treatments in each replication were randomly selected and the shoot and root lengths were measured in centimeters. The average of ten seedlings was calculated and recorded. Vigour index was calculated by using following formula (Abdul baki *et.al.*, 1973):

$$\text{Vigour Index} = (\text{Root length} + \text{Shoot length}) \times \text{germination percentage}$$



Fig 1: Germination stage of cowpea plants **Fig 2: Root and shoot length of cowpea plants**

Fresh and dry weight of seedling (g)

The fresh as well as oven dry weight of ten randomly selected eight days old normal seedlings were recorded. The average of ten seedlings was calculated for both, fresh and dry weight.

Moisture content of seeds

Moisture percentage was determined by grinding seeds in a grinding mill and drying the 5 g ground sample at a constant temperature of 130⁰C for one hour (Anonymous, 1985). The percent moisture content was calculated on net weight basis.

RESULTS AND DISCUSSION

Effect of seed size on thousand seed weight

The variation in 1000- seed weight due to seed sizes was significant. Large seeds had significantly more thousands seed weight (118.3 g) over other grades (107.2 g to 66.8 g). These results were in agreement with those reported by salih (1982) in Fababean, Vadivelu and Ramkrishnan (1983) in bengal gram and Jha *et.al.*(1985) in wheat.

Effect of seed size on germination percentage

Varieties had significant effect on this trait and cv. Bundel Lobia-1 (93.0%) had more germination percentage than cv. UPC-607 (86.48%).

On contradictory to the report of Abdullahi and Vanderlip (1972) in sorghum, Randhawa *et.al.* (1973) in Kalyansona wheat, Bhor *et.al.* (1987) in gram. Randhawa *et.al.* (1990) in maize reported that seed size did not improve field emergence and germination percentage.

Table 1 : Effect of seed size on thousand seed weight (g) and germination percentage

Genotypes	Thousand seed weight (g)	Germination percentage
5% (graded)		
5% (medium)		
5% (all)		
5% (medium)		
5% (all)		
5% (medium)		
5% (all)		

Interaction effect due to varieties and seed sizes for thousand seed weight (g).

Genotypes	Thousand seed weight (g)	SE±	t.D. at 5%
5% (graded)		331	8
5% (medium)			
5% (all)			
5% (medium)			
5% (all)			

Effect of seed size on vigour index

In present investigation vigour index significantly differ due to varieties. But no influence of seed size observed on seedling vigour index. These findings are in conformity with those reported by Marcos *et.al.* (1977). However, China and Phul (1982) in pearl millert and Kalakannawar *et.al.* (1989) in wheat reported that large sized seeds produce more vigorous seedlings. On contrary, Mugnisjah and Nakamura (1986) in soybean reported that plants from small seed produce more vigorous growth.

In present investigation, medium sized seeds produced more vigour index (3044.33) than all other grades.

Table 2: Effect of seed size on root shoot length and vigour index

Genotypes	Root length (cm)	Shoot length (cm)	Vigour index
5% (graded)			57
5% (medium)			73
5% (all)			33
5% (medium)			
5% (all)			

Effect of seed size on fresh and dry weight of seedlings

The fresh and dry weight of seedlings due to varieties was non- significant. Fresh weight of seedlings due to seed sizes showed significant differences as dry weight of seedling showed non-significant differences (Table 3).

Singh et.al. (1972) in soybean, Bishnoi (1977) in triticale, Evans and Bhatt (1977) in wheat, Vadivelu and Ramakrishanan (1983) in gram reported that larger seeds produced seedlings with higher fresh and dry weight.

Effect of seed size on moisture content of seed

Non- significant differences were obtained due to varieties, seed sizes and their interactions for moisture content of seed. The moisture content of cv. Bundel Lobia-1 was more than that of cv. UPC-607. Large seeds showed highest moisture content (13.07%) followed by medium seeds (12.95%), ungraded seeds (12.90%) and small seeds (12.29%) respectively.

Table 3. Effect of seed size on moisture content of seed

Treatments	Moisture content (%)
Variety	
V ₁	13.37
V ₂	12.23
S.E.±	0.433
C.D. at 5%	N.S.
Seed size	
A (Ungraded)	12.90
B (Large)	13.07
C (Medium)	12.95
D (Small)	12.29
S.E.±	0.613
C.D. at 5%	N.S.
Interaction (Variety X Seed size)	
S.E. ±	0.866
C.D. at 5%	N.S.

The difference due to variety x seed size interaction were non- significant for all traits except thousands seed weight.

CONCLUSION

The result indicated that large seeds do not have advantages over small seeds in terms of germination. Therefore, sorting it out will be of no economic return but time consuming. Large seeds size showed better vigour in seedling dry weight.

However, the difference due to variety x seed size interaction were non- significant for all traits except thousands seed weight. Seed size is not therefore important in germination of cowpea seed, sorting it out will be of no economic return but time consuming.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

Abdual-Baki. A., James. D, and Anderson. D. 1973. Vigour determination in soybean seed by multiple criteris. *Crop Sei.* 13 (6) : 630-633.

Abdullahi and Vanderlip, R.L. 1972. Relationship of vigour tests and seed source and size to sorghum seedling establishment. *Agron. j.* 64 :143-144.

Anonymous. 1985. international rules for seed testing and Annexes *seed Sei. and Tech.* 13 (2) ; 356-313.

Bhor, S.R. 1987. Effect of seed size on growth, yield and yield contributing characters in gram (*Cicer arietinum L.*). M.Sc. (Agri.) Thesis, M.P.K.V.. Rahuri (MS.) India

Bishnoi, U.R. and Sapra, V.T. 1975. Effect of seed size on seedling growth and yield performance in hexaploid triticale. *Cereal Research Communication.* 3(1) : 49-60.

Bremner, P.M., Eckersall, R.N. and Scott. R.K, 1963. The relative importance of embryo size and endosperm size in causing the effect associated with seed size in wheat, *J. agrie. Sci.* 61(1) : 139-145.

China, B.S. and Phul, P.S. 1982. Association of seed size and seedling vigour with various morphological traits in pearl millet. *Seed Sci. and Tech.* 10(3) ; 541-545.

Evans, L.E. and Bhatt, G.M. 1977. Influence of seed size protein content and cultivar on early seedling vigour of wheat. *Canadian J. PlantSci.* 57(3) : 929-935.

Jha, B.N., Sinha, S.K. and Singh, J.N 1985. Effect of seed Size on yield in wheat (*Triticum aestivum*). *Seed Res.* 13(1) : 24-27.

Kalakannavar, RM-, Shashidhara, S.A. and Kulkarni, G.N. 1989. Effect of grading on quality of wheat seeds. *Seed Res.* 17(2) : 182-185.

Marcos, F.J., Silva, A.F. , Da, Cicero, S.M. and Goncalves, C.A.R. 1977, Effect of seed size on germination, vigour and yield of maize (*Zea mays L*) Luiz de queiroz. 34 : .327-337. (Seed Abstr. 1981. 3708).

Mugnisjah, W.Q. and Nakamura, S. 1988. Vigour of soybean seed as influenced by sowing and harvest dates and Seed Size. *Seed Sci. and Tech.* 14: 87-94.

Randhawa, H.S., Dey, S.K., Kaur, J., Sharma, H.L., Hari Singh and Khehra, A.S. 1990. Studies on seed germination, seedling vigour and seed mycoflora of graded maize (*Zea mays* L.) *Seed Res.* 6(1) : 49-52.

Salih, F.A. 1982. Influence of seed size and sowing date on yield components of Fababenn. *Fabis News Letter.* 6(4) : 38-39. (From *Seed Abstr.* 6:1-12) pp. 161.

Singh, J.N., Tripathi, S.K. and Nagi, P.S. 1972. Note on the effect of seed size on germination, growth and yield of soybean, [*Glycine max* (L.) Merr.]. *The Indian J agric. Sci.* 42(1) : 83- 86.

Vadivelu, K.K. and Ramakrishnan, V. 1983. Effect of seed size on quality attributes and yield of seed in Bengal gram. (*Cicer arietinum* L.). *Seed Res.* 11(2) : 177-181.