

Influence of sowing methods and phosphorus levels on Growth, Yield and Economics of Summer Mungbean (*Vigna radiata* L.)

ABSTRACT: A field experiment was conducted during the *Rabi* season of 2023-24 at the Agricultural Research Farm, School of Agricultural Sciences and Technology, RIMT University, Mandi Gobindgarh, Punjab to study the “Effect of methods of sowing and phosphorus levels on the growth, yield attributes and yield of summer mungbean (*Vigna radiata* L.)”. The experiment consisted of nine treatment combinations with methods of sowing (Flat, Ridge and Bed sowing) and three phosphorus levels (30, 40 and 50 kg/ha P₂O₅) were tested in Randomized Block Design (RBD) with three replications. The results indicated that the crop responded significantly to sowing methods and phosphorus in respect of growth, yield and economics such as plant height (cm), number of nodules/plant, number of branches, dry matter accumulation/plant (g/m²) and yield characteristics viz., number of pods/plant, number of seeds/pod, test weight (g), seed yield (q/ha), straw yield (q/ha), harvest index (%), net return (Rs/ha) and B:C ratio. In the combination of sowing methods and phosphorus, all the parameters were significant by influence. The combination of Bed sowing + 50 kg/ha P₂O₅ treatment had the highest values for growth characteristics, yield characteristics and also performed economically well as compared to other treatments.

Keywords: Summer mungbean, sowing methods, phosphorus levels, growth parameters, yield attributes and yield.

1. INTRODUCTION

Pulses play a significant role in everyday diets, especially on the Asian continent [1]. Pulses offer noteworthy nutritional and health advantages and have been shown to mitigate several non-communicable illnesses, including cardiovascular and colon cancer. Pulses are the main stay of Indian agriculture and have been used as food, fodder and feed. Animals can also be fed high-quality grain made from pulses [2]. India is the largest producer (25% of global production), consumer (27% of world consumption) and importer (14%) of pulses in the world. Pulses account for around 20 percent of the area under food grains and contribute around 7-10 percent of the total food grains production in the country. Pulses are grown in both *Kharif* and *Rabi* seasons; *Rabi* pulses contribute more than 60 percent of the total production [3]. Pulses are grown across the country, in India, the area under pulses is about 28.90 million ha, with production of 26.05 million tonnes and productivity about 902 kg/ha. In Punjab, the area under pulses is about 0.08 lakh ha, with production of 0.08 lakh tonnes and productivity about 947 kg/ha [4].

Mungbean (*Vigna radiata*) world's largest short-duration pulse crop grown in northern India which belongs to Fabaceae family of the genus *Vigna*. It has been known as “moong.” It is widely cultivated throughout the Asia [5]. Mungbean grown in Southeast Asia, Africa, South America and Australia and serves as a major source of dietary protein for the vast majority of vegetarian people [6]. Cultivation of mungbean can improve the physical, chemical and biological properties of soil as well as increase soil fertility status through nitrogen fixation from atmosphere by symbiotic process with the help of micro-symbionts (Rhizobium). In India, area of mungbean is 5.55 million hectares, with the total production of 3.68 million tonnes with a productivity of 663 kg/ha [7].

Commented [H1]: Please check the value

To ensure optimal root system development from the outset of crop growth, efficient sowing techniques maximize the availability of resources, including moisture, sunshine, and nutrients. A good yield can only be achieved by using the sowing methods, which is one of the most important agricultural activities. Among other agronomic practices, proper method of planting may considerably increase the production of green gram. Ideal planting geometry is precious and important for better and efficient utilization of available plant growth resources in order to get maximum productivity [8]. Among different sowing methods such as flat sowing, ridge sowing and bed sowing methods by using the right techniques can significantly boost mungbean yields. Flat sowing involves sowing seeds or transplanting seedlings on a surface level [9]. The ridge seeding approach improved the conditions for plant root development, increased nutrient absorption, and increased biomass output [10 a]. Bed planting is one of the more successful agronomic interventions when it comes to minimizing the damage caused by water logging, especially during the rainy season, especially in places with significant rainfall and irrigation in the ridge and furrow planting technique decreased the incidence of illness in pulses.

Commented [H2]: 10 a ??

Mungbean is highly responsive to fertilizers and has a considerable response to phosphorus. The conventional approach to treating phosphorus shortage in the soil has been to apply phosphorus fertilizers. Every year, 52.3 billion tons of fertilizers are sprayed on soil and plant systems to maintain constant phosphorus levels [11]. The organic matter composition of the soil decreases under these circumstances, but the immobilization of nutrients like phosphorus increases [12, 13]. Considering the above facts the present study is aimed at following objectives to determine the effects of methods of sowing and phosphorus levels on the growth, yields attributes and yield of summer mungbean.

2. MATERIALS AND METHODS

2.1 Experimental site: A field experiment was laid out during *Rabi* 2023-24 at the Agriculture Research Farm, RIMT University, Mandi Gobindgarh, Punjab. The experimental site (Mandi Gobindgarh) is situated in Punjab at 30.6642° N latitude and 76.2914° E longitude at an altitude of 268 meters. The annual rainfall of Mandi Gobindgarh is 730.2 mm, about three-fourth of which is contributed by the south-west monsoon during July to September. Scanty rainfall is received during winter months of December, January to February. With highs between 40 and 45 °C (104 and 113 °F), summers are quite hot.

Formatted: Font: Not Italic

2.2 Edaphic condition: The composite soil samples from 0-15 cm profile layers were collected before planting from randomly selected sites and analysed for initial soil reaction, electrical conductivity and fertility status. The soil of the experimental field was sandy loam in texture with pH 8.4. It was moderately fertile, being low in available organic carbon (0.38%), available nitrogen (144.6 kg/ha) and medium in available phosphorus (17.3 kg/ha) and medium in available potassium (168 kg/ha). Organic carbon in the soil was 0.38% which was estimated by rapid titration method given by Walkley and Black (1966) [14]. The available Nitrogen in soil was 144.6 kg/ha, which was estimated by the Alkaline potassium permanganate method given by Subbiah and Asija (1965) [15]. The available Phosphorus was 17.3 kg/ha estimated by Olsen's method given by Olsen (1954). The available K was 261.3 kg/ha which was estimated by the Flame photometer Ammonium extractable method given by Jackson (1967) [16].

Commented [H3]: It is missing in reference section

2.3 Treatment details: The experiment was laid out in Randomized Block Design with nine treatments and three replications. The treatments were allocated randomly in each block. The details of the treatment are as follows:

Table 1: List of the Treatment Combinations

Notations	Treatment Combinations
-----------	------------------------

T ₁	Flat sowing + 30 kg/ha P ₂ O ₅
T ₂	Flat sowing + 40 kg/ha P ₂ O ₅
T ₃	Flat sowing + 50 kg/ha P ₂ O ₅
T ₄	Ridge sowing + 30 kg/ha P ₂ O ₅
T ₅	Ridge sowing + 40 kg/ha P ₂ O ₅
T ₆	Ridge sowing + 50 kg/ha P ₂ O ₅
T ₇	Bed sowing + 30 kg/ha P ₂ O ₅
T ₈	Bed sowing + 40 kg/ha P ₂ O ₅
T ₉	Bed sowing + 50 kg/ha P ₂ O ₅

2.4 Preparation of the experimental field and application of fertilizers: The experimental field was ploughed once with disc harrow and twice with cultivators each followed by planking in standing water to bring the field to a good puddle. Plots were prepared according to the layout of the experiment. There are three methods of sowing viz., flat, ridge and bed sowing. Seeds were sown manually in the furrow opened by *kudal* at a row distance of 30 cm and plant distance of 10 cm with the seed rate of 30 kg/ha was used for optimum maintenance of plant population. Immediately after sowing seeds were covered with soil. Gap filling and thinning of the plants were done after one week & 20 days after sowing to get optimum yield. Depending upon the climatic situation and soil factors there was a requirement of three irrigations (13th March, 6th April and 6th May). Water logging in the field should be avoided to ensure healthy seeds. A uniform dose of nitrogen through urea was applied in two splits viz., the half dose of nitrogen applied at the time of sowing and rest of the dose was applied at 45 days after sowing. The Phosphorus was applied as per the treatments at the time of sowing through single super phosphate.

Commented [H4]: Please mention the year

3. RESULTS AND DISCUSSION

3.1 Growth parameters

3.1.1 Plant height as influenced by given treatments: Plant height of summer mungbean recorded at growth stages as affected by methods of sowing and phosphorus levels have been presented in (Table2). Plant height increased successively with age of crop. It is evident from the data that different methods of sowing and phosphorus levels influenced plant height significantly at 50 days after sowing and at harvest. It was quite evident from the data that higher plant height was obtained in T₉ (Bed sowing + 50 kg/ha P₂O₅) at 50 days after sowing (36.30 cm) and at harvest (44.10 cm) while the lowest plant height was recorded in T₁ (Flat sowing + 30 kg/ha P₂O₅) at 50 days after sowing (28.00 cm) and at harvest (38.00 cm). The result obtained from the present study was similar with the findings, Ram et al., 2018 [17 b] reported that bed sowing technique was significantly superior over ridge and flat methods in term of growth.

Commented [H5]: 17 b is not in reference section. Please correct it

Formatted: Font: Not Italic

3.1.2 Number of nodules/plant as influenced by different days: Perusal of the data in (Table 2), revealed that number of nodules/plant of summer mungbean recorded at growth stages and has significant effect on summer mungbean. Among different treatments combination of different methods of sowing and phosphorus levels the maximum treatment with Bed sowing + 50 kg/ha P₂O₅ number of nodules/plant was recorded at 50 days after sowing (27.50) and at harvest (21.50) in summer mungbean and minimum number of nodules/plants was found in treatment with Flat sowing + 30 kg/ha P₂O₅ at 50 days after sowing (27.50) and at harvest (21.50). The result obtained from the present study was similar with the findings by Muthu et al., 2016 [18] who observed significantly higher number of nodules in green gram at wider spacing than at narrow spacing.

Formatted: Font: Not Italic

3.1.3 Number of branches: Data presented in (Table 2), depicted those treatments of number of branches of summer mungbean recorded at growth stages and has significant effect on summer mungbean. The effect of sowing methods and phosphorus levels was observed significantly maximum number of branches on summer mungbean with T₉ (Bed sowing + 50 kg/ha P₂O₅) (25.00 and 23.50) and minimum number of branches was found with T₁ (Flat sowing + 30 kg/ha P₂O₅) at 50

days after sowing and at harvest (17.90 and 14.66). The result indicated that the higher branches/plant with higher phosphorus levels might be due the reason that phosphorus played an important role in better development of plant. Similar results were found by Venkatarao *et al.*, 2017 [19].

Formatted: Font: Not Italic

3.1.4 Dry matter accumulation (g/m²): Dry matter accumulation as influenced by different methods of sowing and phosphorus levels have been presented in (Table 2). It is quite obvious from the data that dry matter accumulation varied significantly due to different methods of sowing and phosphorus levels at all the stages of summer mungbean. Higher dry matter accumulation was recorded on summer mungbean with Bed sowing + 50 kg/ha P₂O₅ (13.33 and 17.93 g/m²) and minimum dry matter accumulation was found with Flat sowing + 30 kg/ha P₂O₅ at 40, 50 days after sowing and at harvest (9.00 and 13.33 g/m²). Rahman *et al.*, 2019 found similar results in dry matter accumulation was significantly influenced by the interaction of sowing methods and variety. Phosphorus application had a significant effect on the dry matter accumulation results reported by Erman *et al.*, 2009 [20].

Formatted: Font: Not Italic

3.2 Yield parameters

3.2.1 Number of pods/plant: The data on the effect of different sowing methods and levels of phosphorus on number of pods/plant have been presented in (Table 3). They showed that the effect of sowing methods and phosphorus levels was observed significantly highest the number of pods/plants on summer mungbean with Bed sowing + 50 kg/ha P₂O₅ (15.45) and lowest number of pods/plants was found in treatment T₁ (13.03). The result obtained from the present study was similar with the findings of Singh, 2007 [21] also reported that varieties had a significant effect on number of pods/plants of mungbean.

3.2.2 Number of seeds/pod: It is evident from the data presented in Table 3, on the effect of different sowing methods and levels of phosphorus on number of seeds/pod. It has been noticed that the maximum number of seeds/pods on summer mungbean treatment combinations with (Bed sowing + 50 kg/ha P₂O₅) (6.02) and found minimum number of seeds/pods in T₁ (Flat sowing + 30 kg/ha P₂O₅) (4.30). Similar findings were made by Tigga *et al.*, 2017 [22] reported that the bed sowing method have maximum seed/pod with narrow spacing. This could be due to different variability and adaptability to the environment of green gram. The result is agreed with the findings of Kumar *et al.*, 2012 [23] in phosphorus levels.

Formatted: Font: Not Italic

Formatted: Font: Not Italic

3.2.3 Test weight (g): Test weight as affected by different sowing methods and levels of phosphorus has been presented in (Table 3). They showed that the effect of sowing methods and phosphorus levels was observed significantly maximum test weight on summer mungbean in T₉ (36.24g) and recorded minimum test weight in T₁ (31.33 g). Hossain and Hamid 2005 [24] also reported that test weight under bed sowing as compare to flat sowing of summer mungbean. Parihar and Tripathi, 1990 [25] also revealed that test weight increased significantly with increasing phosphorus levels up to 50 kg/ha P₂O₅.

3.2.4 Seed and Straw yield (q/ha) as influenced by given treatment: Significant variation was observed on the seed yield and straw yield of mungbean when different methods of sowing and phosphorus levels were applied (Table 3). The highest seed yield and straw yield was recorded in Bed sowing + 50 kg/ha P₂O₅ (12.66 q/ha and 18.76q/ha) and lowest seed yield was recorded in treatment combination with Flat sowing + 30 kg/ha P₂O₅ (6.69 q/ha and 12.69 q/ha). Rahim *et al.*, 2014 [26] was found similar results in sowing methods of seed yield. Goswami and Jat, 2003 [27] also reported that bed planted mungbean gave significantly more straw yield as compare to flat sowing.

Formatted: Font: Not Italic

3.2.5 Harvest index (%): Harvest index exhibited significant effects on different methods of sowing and phosphorus levels have been presented in (Table 3). They observed that the effect of

sowing methods and phosphorus levels was observed significantly maximum on summer mungbean harvest index in T₃- Bed sowing + 50 kg/ha P₂O₅ (40.29 %) and minimum harvest index in T₃- Flat sowing + 50 kg/ha P₂O₅ (34.33 %). Harvest index was significantly influenced by different planting methods. The similar results observed by Chen et al., 2010 [28] and Kumar et al., 2012 [29] also reported that, the increasing rate of phosphorus application significantly increased harvest index over the control plots.

Formatted: Font: Not Italic

Formatted: Font: Not Italic

4. Economics

4.1 Net return and benefit cost ratio (Rs/ha): Economic analysis showed that the combined application of Bed sowing + 50 kg/ha P₂O₅ resulted in significantly maximum net return (Rs 64,650.00) and benefit cost ratio (3.38) than the other methods of sowing and phosphorus levels treatment. The minimum net return (Rs 28,148.00) and benefit-cost ratio (2.38) was recorded under Flat sowing + 30 kg/ha P₂O₅ (Table 3). This might be owing to more seed yield in Bed sowing + 50 kg/ha P₂O₅ levels treatment. These findings are in agreement by Patel et al., 2004 [30].

Formatted: Font: Not Italic

Table: 2 Effect of different sowing methods and levels of phosphorus on plant height, number of nodules/plants and number of branches of the summer mungbean at 50 days after sowing and at harvest.

Treatment combination	Plants height (cm)		Number of nodules/plant		Number of branches		Dry matter accumulation (g/m ²)	
	50 DAS	At harvest	50 DAS	At harvest	50 DAS	At harvest	50 DAS	At harvest
T1: Flat sowing + 30 kg/ha P ₂ O ₅	28.00	38.00	21.66	15.66	17.90	14.66	9.00	13.33
T2: Flat sowing + 40 kg/ha P ₂ O ₅	28.33	38.33	22.00	16.00	19.26	16.00	9.50	13.88
T3: Flat sowing + 50 kg/ha P ₂ O ₅	29.44	39.44	22.33	16.33	23.33	16.33	9.70	14.10
T4: Ridge sowing + 30 kg/ha P ₂ O ₅	30.33	40.33	23.44	17.44	24.00	17.33	10.20	14.59
T5: Ridge sowing + 40 kg/ha P ₂ O ₅	32.66	42.33	23.66	17.66	23.33	18.33	11.20	15.57
T6: Ridge sowing + 50 kg/ha P ₂ O ₅	34.33	42.45	24.33	18.33	21.66	20.66	11.78	16.20
T7: Bed sowing + 30 kg/ha P ₂ O ₅	34.45	43.22	24.66	18.66	24.00	22.33	12.33	16.88
T8: Bed sowing + 40 kg/ha P ₂ O ₅	35.50	43.50	26.00	20.00	25.00	22.33	12.66	16.98
T9: Bed sowing + 50 kg/ha P ₂ O ₅	36.30	44.10	27.50	21.50	25.00	23.50	13.33	17.93
SEm (±)	0.21	0.19	0.14	0.18	0.28	0.211	0.03	0.03
CD at 5%	0.65	0.58	0.41	0.54	0.84	0.63	0.10	0.09




Fig. 1. Effect of different sowing methods and levels of phosphorus on plant height, number of nodules/plants and number of branches of the summer mungbean (*Vigna radiata* L.) at 50 days after sowing and at harvest:

Table: 3 Effect of different sowing methods and levels of phosphorus on number of pods/plant, number of seeds/ pod, test weight (g), seed yield (q/ha), straw yield (q/ha), harvest index (%), net return (Rs/ha) and Benefit-cost ratio (B:C) of the summer mungbean

Treatment combination	Number of pods/plant	Number of seeds/pod	Test weight (g)	Seed yield (q/ha)	Straw yield (q/ha)	Harvest index (%)	Net return (Rs/ha)	Benefit-cost ratio (B:C)
T1: Flat sowing + 30 kg/ha P ₂ O ₅	13.03	4.30	31.33	6.69	12.69	34.52	28,148.00	2.38
T2: Flat sowing + 40 kg/ha P ₂ O ₅	13.60	4.34	31.68	7.19	13.33	35.04	31,609.00	2.54
T3: Flat sowing + 50 kg/ha P ₂ O ₅	13.07	4.49	32.42	7.25	13.86	34.33	31,002.67	2.44
T4: Ridge sowing + 30 kg/ha P ₂ O ₅	14.10	4.56	32.83	8.49	14.66	36.66	37,567.67	2.57
T5: Ridge sowing + 40 kg/ha P ₂ O ₅	14.56	4.77	33.23	9.26	15.44	37.41	41,707.67	2.64
T6: Ridge sowing + 50 kg/ha P ₂ O ₅	15.20	5.01	33.58	10.90	15.88	40.70	52,990.33	3.03
T7: Bed sowing + 30 kg/ha P ₂ O ₅	15.17	5.35	34.10	10.84	16.64	39.45	52,270.33	2.98
T8: Bed sowing + 40 kg/ha P ₂ O ₅	15.25	6.01	34.44	11.41	17.55	39.41	56,195.33	3.11
T9: Bed sowing + 50 kg/ha P ₂ O ₅	15.45	6.02	36.24	12.66	18.76	40.29	64,650.00	3.38
SEm (±)	0.03	0.006	0.007	0.08	0.003	0.21	607.49	0.03
CD at 5%	0.10	0.018	0.021	0.25	0.010	0.63	1,836.95	0.07




Fig. 2. Effect of different sowing methods and levels of phosphorus on number of pods/plant, number of seeds/ pod, test weight (g), seed yield (q/ha), straw yield (q/ha), harvest index (%), net return (Rs/ha) and Benefit-cost ratio (B:C) of the summer mungbean (*Vigna radiata* L.).

5. CONCLUSION

Based on the one-year study on summer mungbean (*Vigna radiata* L.), it may be concluded that, effect of sowing methods and phosphorus levels was significantly better with treatment combination, Bed sowing + 50 kg/ha P₂O₅, in terms of growth attributes, yield attributes, yield and also performed economically well as compared to other treatments.

REFERENCES

- Singh M, Mishra JS, Bhatt BP. Effect of integrated nutrient management on production potential and quality of summer green gram (*Vigna radiata* L.). *J. of Krishi Vigyan* 2018;**5**(2): 39-45.
- Sahaja Deva. Krishi Vigyan Kendra, Darsi, Prakasam, AndhraPradesh, India *Agron. of Pulse Crops*. Paperback. 2019 ISBN: 978-93-5335-679-8).
- Krishna, Deshmukh MK. An economic analysis of production of pigeon pea in Bemetara district of Chhattisgarh. *The Pharma Innovation J.* 2021;SP-10(9):466-469.
- Anonymous, *Ministry of Agriculture and farmer's welfare*. GOI. 2023.
- Choudhary AK, Rana_DS, Bana_RS, Pooniya, Vijay, Dass_A, Rana_KS, Kaur_R. Agronomy of oilseed and pulse crops. *ICAR-Indian Agriculture Research Institute, New Delhi*. 2015; ISBN 978-9383168-21-7.
- Gupta S, Pratap A. Mungbean: Summer Cultivation in India. *Indian Institute of Pulses Research, Kanpur*- 208 024. Extension Bulletin, 2016 pages: 42
- Anonymous. *Ministry of Agriculture and farmer's welfare*. GOI. 2023
- Yadav_S, Singh_BN. Effect of irrigation schedules and planting methods on yield, attributes and economics of green gram (*phaseolus radiata* L.) under rice – wheat – green gram cropping system. *Plant Archives*, 2014;**14**(1): 521- 523
- Rahim, Khan GD, Fazli H, Waheed U. Effect of Deficit Irrigations and Sowing Methods on Mungbean Productivity. *Journal of Biology, Agriculture and Healthcare*. 2014 a; ISSN 2224-3208.

Commented [H6]: Author is suggested to use abbreviated form of the journal name in case of all the reference

Olsen 1954 is missing. Please incorporate in

10. Ram H, Singh G, Aggarwal N, Sekhon HS. Effect of sowing methods, nutrients and seed rate on mungbean (*Vigna radiata* (L.) growth, productivity and water-use efficiency. *J. of Applied and Natural Sci.* 2018 a;**10**(1): 190 –195.
11. FAO. The state of food security and nutrition in the world. Rome: *Food and Agricultural Organization of the United Nations*. 2020; <http://www.fao.org/3/a-l7695e.pdf>. 2017
12. Seng V, Bell RW, Willett IR. Amelioration of growth reduction of lowland rice caused by a temporary loss of soil water saturation. *Plant Soil*. 2004;**265**: 1-16.
13. Zhou W, Fei LT, Chen Y, Westby AP, Ren WJ. Soil physicochemical and biological properties of paddy-upland rotation: a review. *Sci World J*. 2014; DOI: <http://dx.doi.org/10.1155/2014/856352>
14. Walkley AJ, Black, Iriti A. An examination of the dentary method for determining soil organic matter and proposed modification of the chromic acid titration method. *Soil Science*. 1966;**37**: 29-38.
15. Subbiah BV, Asija GL. A Rapid Procedure for the Estimation of Available Nitrogen in Soils. *Current Science*, 1956;**25**, 259-260.
16. Jackson ML. *Soil Chemical Analysis*. Prentice-Hall of India Pvt. Ltd., New Delhi, 1967;498p.
17. Ram H, Singh G, Aggarwal N, Sekhon HS. Effect of sowing methods, nutrients and seed rate on mungbean (*Vigna radiata* (L.) growth, productivity and water-use efficiency. *J. of Applied and Natural Sci.* 2018 b;**10**(1): 190 –195.
18. Muthu MC, Muduli KC, Mohanty SK, Mishra P, Mohapatra U. Effect of inorganics, biofertilizers and spacing on growth and yield parameters of green gram CV. *The Bioscan* 2016;**11**(2): 1251-125
19. Venkatarao CV, Naga SR, Yadav BL, Koli DK, Rao IJ. Effect of phosphorus and biofertilizers on growth and yield of mungbean (*Vigna radiata* L.). *Int. J. of Current Micro. and Applied Sci.* 2017;**6**(7): 3992-3997.
20. Erman M, Yildirim BNF. Effect of phosphorus application and Rhizobium inoculation on the yield, nodulation and nutrient uptake in field pea (*Pisum sativum* sp. *arvense* L.). *J. of Animal and Veterinary Advances*. 2009;**8**(2): 301-304.
21. Singh RS. Effect of organic and inorganic sources of nutrition on productivity of long duration pigeon pea (*Cajanus cajan* L.) *Environment and Ecology*. 2007;**25S** (Special 3A): 768-770.
22. Tigga BDK, Chandraker TRB, Sushil KB, Manoj D. Effect of different genotype and planting geometry on growth and productivity of rabi season pigeon pea (*Cajanus cajan* L.). *Intl. J. Curr. Microbiol. App. Sci.* 2017;**6**(3): 2188-2195.
23. Kumar R, Singh YV, Singh S, Latore AM, Mishra PK, Supriya. Effect of phosphorus and sulphur nutrition on yield attributes, yield of mungbean [*Vigna radiata* (L.) Wilczek]. *Journal of Chemical and Pharmaceutical Research*. 2012;**4**(5): 2,571–2,573.
24. Hossain MA, Hamid A. Influence of N and P fertilizer application on root growth, leaf photosynthesis and yield performance of groundnut. *Bangladesh J. of Agri. Research*. 2005;**32**(3):369-374
25. Parihar SS, Tripathi RS. Yield, water use and nutrients uptake of wheat as influenced by sowing time, irrigation and nitrogen levels. *Madras agric.* 1990;**J77**: 309-316.

26. Rahim, Khan GD, Fazli H, Waheed U. Effect of Deficit Irrigations and Sowing Methods on Mungbean Productivity. *J. of Biology, Agri. and Healthcare*.2014 b;ISSN 2224-3208.
27. Goswami A, and Jat RK. Productivity and soil fertility as effected by organic manures and inorganic fertilizers in green gram (*Vigna radiata*) - wheat (*Triticum aestivum*) system. *Indian Journal of Agronomy*. 2003;**55**(1):16-21.
28. Chen YP, Rekha PD, Arunshen AB, Lai WA, Young CC. Phosphate solubilizing bacteria from subtropical soil and their tricalcium phosphate solubilizing abilities. *Appl Soil Ecol*. 2010;**34**: 33-41.
29. Kumar R, Singh YV, Singh S, Latore AM, Mishra PK, Supriya. Effect of phosphorus and sulphur nutrition on yield attributes, yield of mungbean [*Vigna radiata* (L.) Wilczek]. *Journal of Chemical and Pharmaceutical Research*. 2012;**4**(5): 2,571–2,573.
30. Patel JJ, Mevada KD, Chotaliya R L. Response of summer mungbean to date of sowing and level of fertilizer. *Indian Journal of Pulses Research*. 2004;**17**(2): 143-144.

UNDER PEER REVIEW