

Optimizing sowing windows for wheat in the sub-temperate climate of the North-Western Himalayas under limited irrigation

Comment [A1]: No experimental treatment explains the term limited irrigation.

ABSTRACT

A field investigation was carried out at Research farm of Department of Agronomy, CSKHPKV, Palampur during the winter season to evaluate the suitability of variable sowing windows under limited irrigation regimes with two wheat cultivars. The five sowing windows 15th, 30th October, 15th, 30th November and 15th December along with two wheat cultivars i.e., HPW-349 and HPW-155 were investigated under a factorial randomized block design with three replications for each treatment. The timely sown crop on 15th October resulted in significantly higher values for growth attributes such as plant height, leaf area index, number of tillers per meter square and yield attributes such as spike length, number of grains per spike and test weight. The grain yield consequently was recorded to be the highest for the timely sown crop i.e., 15th October sown crop. The timely sown crop sown on 15th October resulted in 21.6% higher grain yield as compared to 15th December sown crop. Furthermore, delaying the sowing date by one month from 15th October to 15th November, the wheat grain yield declined by 12%, similarly, delaying the sowing date from 15th November to 15th December resulted in yield reduction by 7%. The highest gross, net returns and benefit cost ratio were also recorded to be the higher for the timely sown crop i.e., 15th October sown crop.

Comment [A2]: Shouldn't the yield decline more with more delay in sowing time ?

Keywords: Crop phenology, sowing windows, varieties, and wheat.

1. INTRODUCTION

Bread wheat (*Triticum aestivum* L.) is one of the important cereal crops in the world with wider production and consumption across the globe. Wheat has been cultivated on an area of 219 million hectares and 808.4 million tons globally, whereas in India it is cultivated on an area of 30.4 million hectares with a production of around 107.7 million tons (FAOSTAT, 2024). India produces a vast quantity of wheat grain on global level being second to the China although have a significant potential in improving the its wheat production potential under the influence of modern improved agro-practices (Jat *et al.*, 2024). In Himachal Pradesh, this crop is presently being cultivated on 319.476 thousand hectares with a production of 609.31 thousand tones and productivity of 1.90 metric tonnes per hectare (Department of Agriculture, 2022-23).

Comment [A3]: Rewrite the statement

Changing climatic conditions like increasing temperatures and declining rainfall have been considered significant reasons for declining wheat productivity especially due to shortened grain filling period leading to reduced grain size for wheat (Farhad *et al.*, 2023). The study on climatic trends in HP also revealed decreasing trends of rainfall and on contrary increase in maximum and minimum temperatures in mountain state Himachal Pradesh affecting crop yield (Rana *et al.*, 2012). Altering the planting schedules had proved to be a critical practice to increase resilience of wheat to changing climate (Pathania *et al.*, 2018; Pathania *et al.*, 2019) in North Western Himalayas conditions. The studies carried out in this region also

indicated that application of rainfall and temperature forecasts proved to be beneficial in saving irrigation in crops (Rana *et al.*, 2013)

Exposing the crop to variable environments with optimized ~~changes~~ sowing window can help in understanding the phenological as well as biomass production response to differing weather elements such as temperature, relative humidity, precipitation, sunshine hours etc (Mangani *et al.*, 2023; Pareek *et al.*, 2023). Besides altering sowing windows, considering potential of variable wheat cultivars to photosynthesise efficiency and accumulate significant amount of biomass in times of changing climate can play a vital role in improving wheat productivity (Sattar *et al.*, 2023). Therefore, considering the need to investigate the influence of variable sowing windows as well as wheat cultivars on productivity and profitability of wheat under the conditions of North-Western Himalayas. The objectives of the study were to study the effect of variable sowing windows and wheat cultivars on wheat growth, productivity, agro-meteorological indices and profitability of wheat.

Comment [A4]: Rewrite clearly

2. MATERIAL AND METHODS

Site and soil

The field investigation was conducted during the winter season ~~of~~ at Research farm of Department of Agronomy, CSKHPKV, Palampur. The experimental field was ~~at an average~~ above mean sea level of 1290.8 m with agro-climatic conditions defined as sub-temperate and sub-humid with high rainfall, severe low temperatures during winters and moderate temperatures during mild summers. During the crop season, the weekly maximum and minimum temperature varied from 10 to 34°C and 0.5 to 25.5°C ~~with~~ and total seasonal rainfall of 370.8 mm. The highest average weekly rainfall (78.6 mm) was recorded during the 4th meteorological week whereas in case of evapotranspiration the corresponding highest (50.5 mm) and lowest values (9.3 mm) were recorded during 16th and 4th meteorological week. The maximum average weekly relative humidity (89%) was recorded during the 3rd meteorological week whereas the lowest relative humidity (48%) was recorded in the 19th meteorological week. The soil at the experimental field was silty clay loam in texture, with acidic soil reaction (pH = 5.7), medium content of available nitrogen (316 kg/ha), phosphorus (16.7 kg/ha) and organic carbon (0.8%) whereas available potassium (298 kg/ha) was high in the field soil.

Experimental detail

The field investigation was carried out in a randomized block design with factorial arrangement. The treatments consisted of five wheat sowing dates ~~windows~~ (15th October, 30th October, 15th November, 30th November and 15th December) and 2 wheat cultivars (HPW-349 and HPW-155) which were replicated thrice. The gross plot size for each treatment was 16.8 m² whereas the net plot size was 13 m². The field was prepared using 1-2 ploughings followed by levelling and breaking of clods. The seed rate was 100 kg/ha whereas the row spacing was kept at 22 cm. The primary nutrients i.e., nitrogen, phosphorus and potassium were applied at the rate of 120, 60 and 30 kg/ha. The half of the nitrogen, full dose of phosphorus and potassium were applied as basal dose whereas rest of the nitrogen was applied at crown root initiation stage as top dressing. Hand weeding was practiced along with Isoproturon application at the rate of 1.25 kg/ha to keep weed population below threshold levels. The harvesting was carried out manually using sickles. The crop was threshed to separate the grains from straw.

Comment [A5]: No information regarding the limited irrigation has been given here.

3. RESULTS AND DISCUSSION

Phenological studies

Crop phenological development serves as a critical biological indicator of climate change and the effects of rising temperatures. Studying crop phenology is essential for understanding how crops respond and adapt to changing climate conditions. In the present

field investigation, the **micro-environment** imposed with alteration of sowing dates, significantly impacted the days to complete the crop emergence, crown root initiation, tillering, ear emergence and physiological maturity. The wheat crop sown on 15th October attained the emergence (10 days) and crown root initiation (28 days) in the shortest span whereas the following late sown crops on 15th (12, 30 days, respectively), 30th November (12, 30 days, respectively) and 15th December (14, 32 days, respectively) took comparatively greater days to attain emergence and crown root initiation (Table 1). Contrary to this, moving ahead for the tillering (68 days), ear emergence (131 days) and physiological maturity (167 days), 15th October sown crop took greater number of days whereas the crop sown on 15th (62, 125 and 156 days, respectively), 30th November (61, 122 and 148 days, respectively) and 15th December (58, 119 and 138 days, respectively) completed the tillering, ear emergence and physiological maturity in a comparatively shorter period of time. Wheat cultivars however, did not differed significantly in terms of days required to attain different phenological stages such as emergence, crown root initiation, tillering, ear emergence and physiological maturity. The results observed in the present investigation were consistent with the findings revealed by Sattar and his co-workers in 2023 wherein they reported that delaying the sowing of wheat crop resulted in extended period of vegetative phases whereas shortened period of reproductive stages. Such a response was recorded due to lower temperature in the winter season for the late sown crop whereas exposure to higher temperatures during reproductive phases and therefore resulting shortened reproductive stages (Sattar *et al.*, 2023).

Comment [A6]: What kind of microclimatic modifications have been done?

Table 1. Effect of varieties and date of sowing on number of days taken to different phenological stages

Treatments	Days to Emergence	Days to CRI	Days to tillering	Days to ear emergence	Physiological maturity
Varieties					
HPW-349	11	30	64	126	156
HPW-155	12	29	63	125	154
LSD (P=0.05)	NS	NS	NS	NS	NS
Sowing window					
15 th October	10	28	68	131	167
30 th October	11	28	68	128	165
15 th November	12	30	62	125	156
30 th November	12	30	61	122	148
15 th December	14	32	58	119	138
LSD (P=0.05)	1.4	1.5	3.6	2.8	4.7

Effect of varieties and date of sowing on growth and yield attributes on wheat crop
Growth attributes

Plant height is a vital growth attribute of cereal crops especially wheat. In the present field investigation, wheat cultivars did not respond significantly different in terms of plant height, however a significant response was recorded in terms of plant height for the crop with variable dates of sowing. The crop sown in the month of October whether 15th (114.8 cm) or 30th October (114.2 cm) resulted in significantly higher taller plants whereas crop sown late on 15th (104.9 cm), 30th November (99.5 cm) and 15th December (94 cm) have plants with shortened height (Table 2). Leaf area index was recorded to be responding differently for the crop sown at varying dates of sowing however the wheat cultivars did not show any significant response in terms of leaf area index. The highest leaf area index was recorded for the crop sown on 15th October (3.24), parallely crop sown on 30th October (3.17) resulted in the similar leaf area index for the wheat crop (Table 2). The crop sown late such as on 15th (2.99), 30th November (2.81) and 15th December (2.92) resulted in considerably lower leaf area index for the wheat crop. Number of tillers per meter square was observed to be

Comment [A7]: Observed comparatively shorter plants

significantly responding to the variable dates of sowing with crop sown on 15th (294) and 30th October (291) resulted in the highest number of tillers per meter square for the wheat crop. Contrary to this, crop sown late on dates such as 15th (280), 30th (270) November and 15th (259) December resulted in substantially lower number of tillers per meter square for the wheat crop. However, the wheat cultivars did not respond in different manner in terms of number of tillers per meter square. Reduction in values of growth attributes such as plant height, leaf area index and number tillers per meter square in the present investigation for delayed sowing conditions can be attributed to exposure of crop to lower temperature during winter season which may have slowed down the metabolic activities leading to curtailed growth. However, timely sown crop may have attained the optimum growth before getting exposed to peak winter season (Jha *et al.*, 2023; Sattar *et al.*, 2023). The findings in the present field study reported on reduction in number of tillers per meter square corroborate with the findings reported by Poudel and his co-workers in his field study reported in 2020.

Comment [A8]: resulting

Yield attributes

The variable dates of sowing produced significant effects over wheat spike length. Spike length was recorded to be significantly higher for the sowing dates of 15th (10.5 cm) and 30th October (10.5 cm). However, late sown wheat crop sown on 15th (9.7 cm), 30th November (9.3 cm) and 15th December (9.0 cm) responded with considerable reduction in spike length (Table 2). Wheat cultivars again did not produce any significant effects over spike length of wheat. Number of grains per spike were found to be significantly influenced under the impact of variable dates of sowing. The timely sown crop on 15th (55) and 30th October (52) resulted in considerably higher number of grains per plant whereas delay in sowing date i.e., 15th (47), 30th November (44) and 15th December (42) resulted in considerably loss in terms of number of grains per plant. The wheat cultivars, however, did not produce any significant effect over number of grains per plant. The test weight of wheat crop was significantly influenced under the impact of variable dates of sowing. The timely sown crop especially on 15th October (48.48 g) resulted in significantly higher test weight for the crop whereas further delay in sowing date such as 15th (45.48 g), 30th November (43.72 g) and 15th December (41.6 g) resulted in reduction in test weight of the crop (Table 2).

In the present investigation, optimized growth under the favourable temperature regime for the timely sown crop might have improved the formation of yield attributes at a better rate. However, contrary to this, reduced growth for the crop sown late might have ultimately resulted in poor formation of yield charactering attributes. Significantly higher values for yield attributes such as spike length, number of grains per spike and test weight for the timely sown crop i.e., 15th October can further be attributed to exposure of crop to optimum temperature during period of flowering as well as grain formation and filling whereas in case of late sown crop or delayed sowing of the crop might have exposed to higher temperatures leading to heat stress and ultimately reduced values for yield attributing characters (Jha *et al.*, 2023). Furthermore, for delayed sowing with increasing temperature during reproductive stages there have been grain shrinkage due to production of ROS (reactive oxygen species) as well as grain abortion (Sattar *et al.*, 2023). Jha and his co-workers and Sattar and his co-workers in different studies based on variable dates of sowing for wheat crop observed similar impact over yield attributes (Jha *et al.*, 2023; Sattar *et al.*, 2023).

Comment [A9]: ??

Crop yield

The dates of sowing considerably influenced the grain yield of wheat crop. The timely sown crop sown on 15th October resulted in 21.6% higher grain yield as compared to 15th December sown crop i.e., as the sowing date was delayed from 15th October to 15th December the grain yield declined from 40.1 to 33 q/ha (Table 2). Furthermore, delaying the sowing date by one month from 15th October to 15th November, the wheat grain yield declined by 12%, similarly, delaying the sowing date from 15th November to 15th December resulted in yield reduction by 7%. The cultivars however did not produce significantly different grain yield for the wheat crop. Straw yield was significantly influenced under the impact of various sowing dates. The timely sown crop of 15th October produced 11.7%

Comment [A10]: Howcome more delayed sowing is leading to lesser decline

higher straw yield compared to 15th December sown crop i.e., 15th October sown crop produced 64 q/ha straw whereas 15th December sown crop produced 57.3 q/ha straw yield. Similarly, delaying crop sowing by one month from 15th October to 15th November and 15th November to 15th December resulted in 5% loss in straw yield. The cultivars did not respond significantly in terms of straw yield of wheat crop.

The lower yield levels in the present field investigation for the delayed sowing conditions can be attributed to reduced growth as well as reduction in the sink size. Sink size in the present study can be attributed to lower test weight, spike length and number of grains per spike. Furthermore, due to shortened reproductive period i.e., ear emergence and physiological maturity for the wheat crop may result in shrinkage of sink size and ultimately reduces the wheat yield (Poudel *et al.*, 2021). The findings of the present field investigation were found to be consistent with the study results reported by Dubey and his coworkers in 2019 in terms of reduction in wheat yield due to exposure of crop to elevated temperatures. However, timely sown crop provides the crop an opportunity to accumulate optimum amount of photosynthates for better sink size and ultimately the economic yield (Dar Ejaz *et al.*, 2018). Poudel and his coworkers reported grain yield reduction to even an extent of 47% with exposure of crop to heat stress especially for late sown crop (Poudel *et al.*, 2021). Reduction in grain yield of wheat with exposure of crop to heat stress was also reported by Poudel and his co-workers in 2020. Also, lower aerosol values, higher net radiant energy during the COVID-19 pandemic lockdown period improved the crop productivity as represented by the higher NDVI during the lockdown period in agricultural ecosystems in the Himalayan foothills (Pokharyal *et al.*, 2021)

Parallel to grain yield, straw yield was reduced considerably in the present study owing to reduced values for growth attributes such as plant height, leaf area index and number of tillers per meter square. Sattar and his co-workers in 2020 reported similar impact over wheat straw yield for delayed sowing of wheat crop with exposure to elevated temperatures. The results of the present study corroborate with findings reported by Sattar and Srivastava in 2021 over wheat straw yield with late sown wheat crop.

Table 2. Effect of varieties and date of sowing on growth and yield attributes on wheat crop

Treatment	Plant height (cm)	LAI	No. of tillers/m ²	Spike length (cm)	No. of grains	1000-grain weight	Grain yield (q/ha)	Straw yield (q/ha)
Varieties								
V1 (HPW-349)	106.4	3.06	280	9.9	48	45.65	36.25	61.21
V2 (HPW-155)	104.6	2.99	278	9.7	47	44.98	35.83	60.09
LSD (P=0.05)	NS	NS	NS	NS	NS	NS	NS	NS
Sowing window								
15 th October	114.8	3.24	294	10.5	55	48.48	40.14	64.00
30 th October	114.2	3.17	291	10.5	52	47.32	38.38	63.04
15 th November	104.9	2.99	280	9.7	47	45.48	35.32	60.62

30 th November	99.5	2.81	270	9.3	44	43.72	33.37	58.25
15 th December	94	2.92	259	9.0	42	41.6	33.01	57.32
LSD (P=0.05)	5.0	0.2	10.8	0.8	3.5	2.5	1.4	3.7

Crop profitability

Gross returns were not significantly influenced with different wheat cultivars. However, sowing of the crop at different dates have resulted in significant effects over gross returns for wheat cultivation. Crop sown on 15th October generated 21.6% greater gross returns i.e., crop sown on 15th December was responsible for 53640 INR gross returns whereas crop sown on 15th October resulted in 65222 INR gross returns (Table 3). The highest gross returns for the timely sown crop can be attributed to substantially higher grain yield recorded for the treatment.

Wheat cultivars were not able to exert any significant influence over net returns from wheat cultivation, however, sowing of the crop at different dates resulted in significant impact over the net returns generated from the wheat cultivation. The crop sown on the 15th October resulted in significant higher net returns for wheat cultivation i.e., 40.7% higher than the crop sown on 15th December. As the Crop sown on 15th October was able to generate 40012 INR net returns whereas crop sown on 15th December was only able to generate net returns of 28430 INR. The net returns for the timely sown crop on 15th October can be attributed to significantly higher grain yield as well as considerably higher gross returns for the respective treatment.

Benefit cost ratio was not significantly influenced under the impact of different wheat cultivars. Variable dates of sowing resulted in significant influence over benefit cost ratio. Sowing of the crop at 15th October (1.59) resulted in significantly higher benefit cost ratio whereas the lowest benefit cost ratio was recorded for the crop sown on 15th December (1.13) (Table 3). Significant higher benefit cost ratio for the 15th October sown crop can be attributed to considerably higher gross returns for the corresponding crop.

Table 3. Effect of varieties and date of sowing on growth and yield attributes on wheat crop

Varieties	Gross return	Net return	B:C ratio
V1 (HPW-349)	58914	33704	1.34
V2 (HPW-155)	58228	33018	1.31
LSD (P=0.05)	NS	NS	NS
Sowing window			
15 th October	65222	40012	1.59
30 th October	62367	37157	1.47
15 th November	57402	32192	1.28
30 th November	54223	29013	1.15
15 th December	53640	28430	1.13

Comment [A11]: No discussion has been added to this section

LSD (P=0.05)	3827	3827	0.15
--------------	------	------	------

4. CONCLUSION

In the present investigation crop growth, yield attributes, yield and profitability were not significantly influenced by wheat cultivars whereas a significant influence of sowing dates was observed. The present investigation has concluded sowing of wheat crop in the month of October whether 15th (1st fortnight) or 30th October (2nd fortnight) is optimum for considerably higher grain yield of wheat. Delaying sowing by each month i.e., 15th October to 15th November and from 15th November to 15th December reduced the grain yield by 12 and 7%, respectively. Similarly, substantially higher growth and yield charactering attributes such as plant height, leaf index, number of tillers per meter square and spike length, number of grains per spike, test weight were recorded for the timely sown crop i.e., 15th October sown crop. Consequently, the highest gross, net returns and benefit cost ratio was recorded for the 15th October sown crop.

ETHICAL APPROVAL (WHEREEVER APPLICABLE)

The present investigation did not involve any studies on human or animal participants.

REFERENCES

Dar Ejaz A, Brar AS and Yousuf A. 2018. Growing degree days and heat use efficiency of wheat as influenced by thermal and moisture regimes. *Journal of Agrometeorology* 20 (2): 168–170. doi: 10.54386/jam.v20i2.535.

Comment [A12]: Missing in text

Department of Agriculture. 2022-23. Statement showing crop-wise area, production and productivity of principal crops in Himachal Pradesh. Himachal Pradesh. <https://agriculture.hp.gov.in/en/production-2/>

Dubey R, Pathak H, Singh S, Chakraborty B, Thakur AK and Fagodia RK. 2019. Impact of Sowing Dates on Terminal Heat Tolerance of Different Wheat (*Triticum aestivum* L.) Cultivars. *National Academy Science Letters* 42: 445–449. doi: 10.1007/s40009-019-0786-7.

FAOSTAT. 2024. Food and Agriculture Organization. Rome, Italy. <https://www.fao.org/faostat/en/#data>

Farhad M, Kumar U, Tomar V, Bhati PK, Krishnan JN, Kishowar-E-Mustarin, Barek V, Brestic M and Hossain A. 2023. Heat stress in wheat: a global challenge to feed billions in the current era of the changing climate. *Frontiers in Sustainable Food Systems* 7. 1203721.

Jat SL, Jat ML, Parihar CM, Jat HS, Singh AK, Bijarniya D and Kumar M. 2024. On-farm evidence on breaking yield barriers through optimizing wheat cropping system in Indo Gangetic Plain. *European Journal of Agronomy* 159. 127256.

Jha RK, Sattar A, Singh AK, Kundu MS, Tiwari RK, Singh AK et al. 2023. Managing climatic risks in rice–wheat cropping system for enhanced productivity in middle Gangetic plains of India. *Frontiers in Sustainable Food Systems* 7. 1259528.

Mangani R, Gunn KM and Creux NM. 2023. Projecting the effect of climate change on planting date and cultivar choice for South African dryland maize production. *Agricultural and Forest Meteorology* 341. 109695.

Pareek B, Rana RS, Rana N, Manuja S, Kumar N and Pareek NK. 2023. Wheat productivity and profitability under different sowing window and irrigation scheduling based on weather model and spatial data in north Western Himalayas. *Journal of Soil and Water Conservation* 22(2): 186-192.

Pathania R, Prasad R, Rana RS and Mishra SK. 2019. Heat unit requirement and yield of wheat (*Triticum aestivum* L.) varieties under different growing environment in mid hill conditions of Himachal Pradesh. *Journal of Agrometeorology* 21(3): 282-287.

Pathania R, Prasad R, Rana RS, Mishra S and Sharma S. 2018. Growth and yield of wheat as influenced by dates of sowing and varieties in north western Himalayas. *Journal of Pharmacognosy and Phytochemistry* 7(6): 517-520.

Pokhariyal S, Patel NR, Rana RS and Chauhan P. 2021. Environmental Impact of Lockdown Amid COVID-19 Over Agricultural Sites in Himalayan Foothills. *Journal of the Indian Society of Remote Sensing* 49: 1651–1659. <https://doi.org/10.1007/s12524-021-01343-4>

Poudel MR, Ghimire S, Pandey MP, Dhakal KH, Thapa DB and Paudel HK. 2020. Evaluation of wheat genotypes under irrigated, heat stress and drought conditions. *Journal of Biology and Today's World* 9(1): 1–3.

Poudel PB, Poudel MR and Puri RR. 2021. Evaluation of heat stress tolerance in spring wheat (*Triticum aestivum* L.) genotypes using stress tolerance indices in western region of Nepal. *Journal of Agriculture and Food Research* 5. 100179. doi: 10.1016/j.jafr.2021.100179.

Rana RS, Bhagat RM, Singh MM, Kalia V, Singh S and Prasad R. 2012. Trends in Climate variability over Himachal Pradesh. *Journal of Agrometeorology* 14(1): 35-40.

Rana RS, Sood R and Shekhar J. 2013. Validation of medium range weather forecasts in sub-temperate and sub-humid climate of western Himalayas. *Indian Journal of Agricultural Sciences* 83(12): 1357-1363.

Sattar A and Srivastava RC. 2021. Modelling climate smart rice-wheat production system in the middle Gangetic Plains of India. *Theoretical and Applied Climatology* 144: 77–91. doi: 10.1007/s00704-020-03497-6.

Sattar A, Nanda G, Singh G, Jha RK and Bal SK. 2023. Responses of phenology, yield attributes, and yield of wheat varieties under different sowing times in Indo-Gangetic Plains. *Frontiers in Plant Science* 14. 1224334.