

Original Research Article
Evaluating Particulate Matter (PM₁₀ and PM_{2.5})
Concentrations in Educational Institutions within South City
Corporation, Dhaka, Bangladesh

ABSTRACT

Dhaka is widely recognized as the most polluted city in Bangladesh due to its high levels of dust. In Dhaka South City Corporation, almost all of the educational institutions are built on the side of the main road. The objective of this study is twofold. First, it aims to identify the status of air pollution within educational institutions located in the Dhaka South City Corporation. Second, it seeks to assess the relationships among different zones and various air quality parameters, specifically focusing on particulate matter levels of PM₁₀ and PM_{2.5}. This study was conducted in 37 locations of Dhaka metropolitan, by using Aeroqual s500. It is found that average concentrations of PM₁₀ and PM_{2.5} in 37 places in the study area were 68.82 and 113.95 µg/m³ respectively. The average concentration of PM₁₀ and PM_{2.5} of all the zones was found more than one time higher than NAAQS standard level. It is estimated that the average PM₁₀/PM_{2.5} was 60.41%. It also revealed that the concentrations of PM₁₀ may change significantly between and within the land uses as the P values are less than 0.05. The outcome of this research is arranged in descending order based on the average concentration of PM₁₀ which follows as Zone-5 (Chak Bazar, Bangshal, Kotwali) > Zone-2 (Dhanmondi) > Zone-6 (Sutrapur, Gendaria, Jatra Bari) > Zone-3 (Ramna, Shahbagh) > Zone-1 (Hazaribagh) > Zone-4 (Lalbagh, New Market) > Zone-7 (Kadamtali).

Keywords— Air Pollution, Particulate Matter, Zones, Dhaka South City Corporation, Bangladesh.

1. INTRODUCTION

Air pollution has risen to the status of a major environmental health concern in many nations throughout the globe. Human exposure is the most obvious consequence of pollution, but the hidden cost to society from these contaminants' impacts on people's health and the ecosystem is also significant[1]. Particulate Matter has a harmful effect not only on human health but also on the environment and can raise respiratory problems and global warming, respectively [2].

However, the inventory of air pollutants is declining in Europe and the USA, while in Asia, the situation is worsening in real-time [3]. Bangladesh has the foulest air in South Asian countries and each year 21% of all deaths in the country are caused due to a long-time exposure of high PM contain air [4]. Dhaka, the most populated city in the world and the capital of Bangladesh was identified as the fifth most polluted capital city in the world [5]. The level of air pollution in Dhaka city is now a quite concerning issue, with the AQI concentrations ranging from "Unhealthy" to "Extremely Unhealthy" for two-thirds of the year [6]. The city has grown increasingly crowded during the previous ten years. The municipal infrastructure is being built, locals are building high-rise residential and commercial developments to accommodate the city's growing population [7] which helps to increase air pollution. The massive and uncontrolled increase of emissions from industrial activities and transportation specifically, personal vehicles in Dhaka city deteriorates the air quality [8]. These two parts use a great amount of fuel that resulted in significant increases in emissions as well as economic losses [9]. Vulnerable groups like school going children are the major exposures and sufferers of PM because their immune and whole-body systems are under developing stage at this age [10]. In addition to having an adverse effect on children's physical and mental growth, air pollution aggravates respiratory diseases including asthma and seasonal allergic rhinitis (SAR), better known as hay fever, which is most frequent in schoolchildren [11, 12]. The proximity of educational institutions to major roads and busy traffic areas exacerbates the exposure of students and staff members to harmful pollutants. The levels of $PM_{2.5}$ near schools in Dhaka City exceeded the World Health Organization's (WHO) recommended guidelines, posing a significant health risk to the vulnerable population, especially children [13].

2. OBJECTIVES

The objectives of the Studies are:

1. To identify the status of air pollution in educational institution of Dhaka South City Corporation.
2. To assess the relationship among Zones and parameters ($PM_{2.5}$ and PM_{10}).

3. METHODOLOGY

a. Study Area

The study was conducted in Dhaka South city corporation. Data of Particulate matter concentration ($PM_{2.5}$ & PM_{10}) were recorded continuously 1-min for the average

concentrations. Aeroqual monitors were used for data collection and placed tripods at ~1.5 m elevation above the ground which were positioned horizontally. The monitoring location was nearest to the road with no other primary pollutant sources nearby. We divided the study area into seven zones which include in total 37 educational institution locations of Dhaka South City Corporation area. The zones are divided as following, Hazaribagh area; Zone-(1), Dhanmondi area; Zone-(2), Ramna and Shahbagh area includes in Zone-(3), than Lalbagh and New Market area consisted in Zone-(4), Chak Bazar, Bangshal and Kotwali resided in Zone-(5), after that Sutrapur, Gendaria and Jatra Bari area involves in Zone-(6) and lastly Kadamtali area; Zone-(7).




Fig. 1: Study Area (Dhaka South City Corporation and Data Collection Locations Point)

b. Data collection

Aeroqual portable monitors (Model: S500) were employed to collect air quality data. These monitors were positioned horizontally on tripods at a height of approximately 1.5 meters above ground level. The monitors were deliberately positioned alongside the road near the educational institution, away from other major sources of pollution, in order to obtain precise measurements. The Aeroqual monitors were configured to measure and record the average concentrations of particulate matter (PM) for a period of 1 minute. In order to commence the monitoring process, the device was activated by pressing and holding the power button until the screen became operational. Prior to commencing data collection, the sensors necessitated a 3-minute warm-up period to eradicate any impurities. A distinct Location ID was allocated to each monitoring site for the purpose of identification.

The sensor autonomously gathered and stored air quality data following the initialization period. Data retrieval was expedited by establishing a connection between the device and a personal computer using a USB cable. The data can be readily downloaded and exported to Excel format using the provided software.



Figure 2: Portable Air Quality Monitor

c. Data processing & Interpretation of Results

The collected data were processed using IBM SPSS V20 and MS Excel 2020. These tools were employed for data input, analysis, and visualization. A variety of graphs, tables, diagrams, and box-whisker plots were created in order to comprehend the properties of the data. Descriptive statistics were utilized to evaluate the spread of each land use characteristic, while ANOVA tests were employed to ascertain the statistical significance of the results. The air quality data is displayed in a thorough manner using an array of graphs, charts, and tables

4. RESULT & DISCUSSION

a. Status of Air Pollution Studies in Educational Institutions Area of South Dhaka City Corporation

The status of air pollution in educational institution zone area of South Dhaka City Corporation were shown in figure 3 (a, b, c, d, e, f, g and h). Study found that in zone 1 (fig. 3 (a)) out of four places in Hajaribag area, the three highest average concentration of PM_{10} and $PM_{2.5}$ were in front of Shahid Sheikh rasel Govt. High School, outside of Hajaribag Model

Town High School and Hajaribag Ideal School & College with the concentration of 73 and 146 $\mu\text{g}/\text{m}^3$; 71 and 96 $\mu\text{g}/\text{m}^3$ and 34 and 103 $\mu\text{g}/\text{m}^3$ respectively. On the other hand, the least PM_{10} concentration was found in front of Hajaribag Girl's High School (31 $\mu\text{g}/\text{m}^3$) and $\text{PM}_{2.5}$ concentration was found outside of Hajaribag Model Town High School (96 $\mu\text{g}/\text{m}^3$). It also found that the most polluted place was 1.12fold higher than NAAQS and 5.84 and 2.92 folds higher than WHO standard. Where, the Bangladesh National Ambient Air Quality Standards (NAAQS) is 65 $\mu\text{g}/\text{m}^3$ set by the Department of Environment (DoE) for concentrations of PM_{10} . Also, the Air Quality Standard (24-hour) set by the WHO for PM_{10} and $\text{PM}_{2.5}$ are 25 and 50 $\mu\text{g}/\text{m}^3$ respectively.

Figure 3 (b) illustrated that among four places in Dhanmondi area, the three highest average concentration of PM_{10} was in front of Dhaka City College (108 $\mu\text{g}/\text{m}^3$), outside of Daffodil University (101 $\mu\text{g}/\text{m}^3$) and Stamford College (84 $\mu\text{g}/\text{m}^3$). Comparatively less polluted place was in front of Dhaka College (77 $\mu\text{g}/\text{m}^3$). The average concentration of $\text{PM}_{2.5}$ found higher outside of Daffodil University (206 $\mu\text{g}/\text{m}^3$), Dhaka City College (116 $\mu\text{g}/\text{m}^3$) and Stamford College (105 $\mu\text{g}/\text{m}^3$) respectively and least polluted place was in front of Dhaka College (91 $\mu\text{g}/\text{m}^3$). The study also indicated that the most polluted place were 1.66 and 1.36 times higher than the national standards. The study estimated that PM_{10} was found in $\text{PM}_{2.5}$ in Zone-2 was 76.69%. The comparison of the average concentration of PM_{10} and $\text{PM}_{2.5}$ of zone-3 area was demonstrated through figure 3 (c). Within seven places in Ramna and Shahbagh area, the three highest average concentration of PM_{10} was University of Dhaka (DU) area (80 $\mu\text{g}/\text{m}^3$), outside of BTCL Ideal School and College (73 $\mu\text{g}/\text{m}^3$) and Sher-E-Bangla School & College (56 $\mu\text{g}/\text{m}^3$) and the least contaminated place was in front of Ibrahim Medical College (32 $\mu\text{g}/\text{m}^3$). The average concentration of $\text{PM}_{2.5}$ found higher outside of BTCL Ideal School and College (152 $\mu\text{g}/\text{m}^3$), University of Dhaka (DU) (121 $\mu\text{g}/\text{m}^3$) and in front of Siddeswari Girls College (109 $\mu\text{g}/\text{m}^3$) respectively and comparatively least contaminated place was in front of Ibrahim Medical College (52 $\mu\text{g}/\text{m}^3$). The most polluted place were 1.23 and 1.01 times higher than the national standards (NAAQS). The figure 3 (d) presented that out of four places of zone-4 in Lalbagh and New Market area, the highest average concentration of PM_{10} were in front of Madrasha Rahmania Hafijia (68 $\mu\text{g}/\text{m}^3$) and outside of Govt. laboratory High School (66 $\mu\text{g}/\text{m}^3$) and comparatively less contaminated place was Dhaka Ideal High School (27 $\mu\text{g}/\text{m}^3$). On the other hand, the average concentration of $\text{PM}_{2.5}$ found higher outside of Govt. laboratory High School (133 $\mu\text{g}/\text{m}^3$), Madrasha Rahmania Hafijia (113 $\mu\text{g}/\text{m}^3$) and in front of

Dhaka Ideal High School ($70 \mu\text{g}/\text{m}^3$) respectively. The least contaminated place was Lalbagh Govt. Model School & College ($58 \mu\text{g}/\text{m}^3$). Study observed that the most polluted place was 1.05 times higher than the NAAQS and 2.72 and 2.66 times higher than WHO standard. The study estimated that the ratio of $\text{PM}_{10}/\text{PM}_{2.5}$ was 50.02% in Zone-4. In figure 3 (e) zone-5 indicated that in Chak Bazar, Bangshal and Kotwali area, out of six places the highest average concentration of PM_{10} and $\text{PM}_{2.5}$ were in front of Haybot Nagar Govt. Primary School (232 and $232 \mu\text{g}/\text{m}^3$), outside of Armanitola Govt. Primary School (120 and $186 \mu\text{g}/\text{m}^3$) and Suritola Model Govt. Primary School (117 and $119 \mu\text{g}/\text{m}^3$) respectively. Moreover, the least contaminated place was outside of Darul Huda Madrasa with PM_{10} and $\text{PM}_{2.5}$ concentration of 50.00 and $88.00 \mu\text{g}/\text{m}^3$ respectively.




Figure 3: Comparison of Particulate Matters of Educational Institutions in Dhaka South City Corporation

Area Based on Zones

However, study found that the most polluted place were 3.60 and 1.55 fold higher than the NAAQS level. The study estimated that in Zone-5, 73.45% PM_{10} was found into $PM_{2.5}$. Study found that in Sutrapur, Gendaria and Jatra Bari area Zone-6, the maximum average concentration of PM_{10} was in front of Central Women's College ($91 \mu\text{g}/\text{m}^3$), outside of Narindra Govt. High School ($79 \mu\text{g}/\text{m}^3$) and Dhaka MahanagarMohila College ($74 \mu\text{g}/\text{m}^3$) and the average concentration of $PM_{2.5}$ found higher outside of Narindra Govt. High School ($151 \mu\text{g}/\text{m}^3$), Dhaka MahanagarMohila College ($142 \mu\text{g}/\text{m}^3$) and in front of Central Women's College ($140 \mu\text{g}/\text{m}^3$) respectively.

Furthermore, the least concentration of PM_{10} and $PM_{2.5}$ was found in front of Salimullah College (45 and $73 \mu\text{g}/\text{m}^3$). In addition to the study found that the most polluted place were 3.60 and 1.55 times higher than the NAAQS. Study found in zone-7 (Kadamtali area), the most average concentration of PM_{10} was in front of Muradpur Govt. Primary & High School ($55 \mu\text{g}/\text{m}^3$), outside of Bangladesh bank Adarsha High School ($45 \mu\text{g}/\text{m}^3$) and relatively less contaminated place was Syampur Govt. Primary School and in front of Shyampur Ideal High School ($44 \mu\text{g}/\text{m}^3$). However, the average concentration of $PM_{2.5}$ found higher outside of Bangladesh bank Adarsha High School ($143 \mu\text{g}/\text{m}^3$), in front of Syampur Govt. Primary School ($87 \mu\text{g}/\text{m}^3$) and Muradpur Govt. Primary & High School ($71 \mu\text{g}/\text{m}^3$) respectively. Comparatively less contaminated places less polluted places were in front of Muradpur Govt. Primary & High School ($71 \mu\text{g}/\text{m}^3$). Moreover, the most polluted place were 2.2 and 2.86 times higher than WHO standard. The study estimated that the ratio of $PM_{10}/PM_{2.5}$ was 53.63% in Zone-7. Finally the comparison of the average concentration of PM_{10} and $PM_{2.5}$ of seven zones in Dhaka South City Corporation. The average concentration of PM_{10} was higher in the Chak Bazar, Bangshal and Kotwali area (Zone-5), Dhanmondi area (Zone-2) and Ramna and Shahbagh area (Zone-3) with the values of 116.00, 92.50 and $69.38 \mu\text{g}/\text{m}^3$ respectively. It was also observed that the average concentration of $PM_{2.5}$ was higher in the Chak Bazar, Bangshal and Kotwali area (Zone-5), and Dhanmondi area (Zone-2) with the values of 154.50, and $129.50 \mu\text{g}/\text{m}^3$ respectively with the highest concentration of PM_{10} and $PM_{2.5}$ in the Zone-5. It was also noticed that the concentrations of PM_{10} and $PM_{2.5}$ found in the most polluted Zone were 1.78 and 1.03 times higher than the national standards. The concentration of PM was found relatively lower in industrial area, residential area and sensitive area. Moreover, the average concentration of PM_{10} ($47.00 \mu\text{g}/\text{m}^3$) was found to be least in Kadamtali area (Zone-7) and $PM_{2.5}$ ($93.50 \mu\text{g}/\text{m}^3$) were found to be least in Lalbagh and New Market area (Zone-4).

***b. Dispersion of the Educational Institutions in Dhaka South City Corporation Area
Based on Zones***

Table 1 presents the descriptive statistics for PM₁₀ and PM_{2.5} across seven different zones. For PM₁₀, the highest concentration range was observed in Zone-5 at 182 µg/m³, followed by Zone-2 at 74 µg/m³, while the lowest was in Zone-7 at 11 µg/m³. The minimum concentration of PM₁₀ across all zones was 27 µg/m³ in Zone-4, and the maximum was 232 µg/m³ in Zone-5. The highest mean PM₁₀ concentration was in Zone-5 at 116.00 µg/m³, and the lowest was in Zone-7 at 47.00 µg/m³. Zone-5 also had the highest standard deviation of 62.78 µg/m³, whereas Zone-7 had the lowest at 5.35 µg/m³. The coefficient of variation for PM₁₀ was highest in Zone-5 at 54.12% and lowest in Zone-7 at 11.39%.

For PM_{2.5}, Zone-5 exhibited the highest range at 144 µg/m³, followed by Zone-2 at 115 µg/m³, and the lowest range was in Zone-1 at 50 µg/m³. The minimum PM_{2.5} concentration was 52 µg/m³ in Zone-3, with the maximum again in Zone-5 at 232 µg/m³. The highest mean PM_{2.5} concentration was found in Zone-5 at 154.50 µg/m³, and the lowest was in Zone-4 at 93.50 µg/m³. The standard deviation was highest in Zone-2 at 52.02 µg/m³ and lowest in Zone-1 at 22.20 µg/m³. The coefficient of variation for PM_{2.5} was highest in Zone-2 at 40.17% and lowest in Zone-1 at 19.47%.

Table 1: Descriptive statistics table for PM₁₀ and PM_{2.5}.

S. N.	Land Use	Number of Land use	PM ₁₀				PM _{2.5}			
			Range (µg/m ³) (Min-max)	Mean (µg/m ³)	Std. Deviation (µg/m ³)	Coefficient of Variation (%)	Range (µg/m ³) (Min-max)	Mean (µg/m ³)	Std. Deviation (µg/m ³)	Coefficient of Variation (%)
1.	Zone-1 (Hazaribagh)	4	42	52.2	22.8	43.74	50	114.0	22.20	19.47
2.	Zone-2 (Dhanmondi)	4	31	92.5	14.4	15.60	115	129.5	52.02	40.17
3.	Zone-3 (Ramna, Shahbagh)	7	48	56.8	15.7	27.71	100	97.29	34.59	35.56
4.	Zone-4 (Lalbagh, New Market)	4	41	47.7	22.2	46.65	75	93.50	35.37	37.83
5.	Zone-5 (Chak Bazar, Bangshal, Kotwali)	6	182	116	62.7	54.12	144	154.5	51.68	33.45
6.	Zone-6 (Sutrapur, Gendaria, Jatra Bari)	8	46	69.3	13.6	19.63	78	113.6	29.16	25.66
7.	Zone-7 (Kadamtali)	4	11	47.0	5.35	11.39	72	95.25	32.50	34.12

The whisker box plot shows the average of PM₁₀ and PM_{2.5} concentrations in seven zones in figure 4. Following whisker box plot of PM₁₀ revealed that Zone-5 had the highest dispersion with extremely negative skewed distribution which contradicted with descriptive statistics where Zone-5 showed highest values along with one lower outlier; followed by Zone-4 and Zone-1 with normally skewed values with normally distributed values. This episode was found in front of Dhaka College. Moderate distribution was seen in Zone-2 with normal skewness and Zone-3 with positive skewness. These situations were found upper outlier in front of Haybot Nagar Govt. Primary School and lower outlier found outside of Darul Huda Madrasa in Zone-5. Less moderate distribution was seen in Zone-6 with negatively skewed values. Zone-7 had the most clustered concentration with extremely positive distributed values.

This whisker box plot of PM_{2.5} revealed that the highest concentration that was seen in Zone-5 with positively skewed distribution followed by Zone-2, Zone-4 and Zone-6 with positive to normal skewed values. Moderate distribution was seen in Zone-7 with positively distributed

values. Less Moderate distribution was seen in Zone-1 with extremely positive skewness, which was found in front of Shahid Sheikh Rasel Govt. High School.




Figure 4: Whisker-box plot showing the concentration of PM_{2.5} and PM₁₀ in Different Zones

c. Significance Test

The table 2 of ANOVA is performed to find whether the changes in the concentration of all the parameters between and within land uses are significant. The F values were calculated to be 4.052 for PM_{2.5} and 1.840 for PM₁₀ where P value of PM_{2.5} and PM₁₀ were 0.004 and 0.125. The following table shows that the concentrations of PM_{2.5} change significantly between and within in the land uses as the P values are less than 0.05. However, PM₁₀ do not change significantly as the P values are greater than 0.05.

Table 2: Significance Test

d. Land Use Based Cluster Analysis




Figure 5: Rescaled Distance Cluster Combine for PM_{2.5} and PM₁₀ in Different Zones

Figure 5 illustrates the dendrogram plots resulting from cluster analyses of PM_{2.5} and PM₁₀ concentrations, with Z-score normalization applied. The PM_{2.5} analysis identified four distinct clusters. The first cluster includes Zone-4, Zone-7, Zone-1, and Zone-3. This cluster merges with the second cluster, which comprises Zone-6, at an approximate distance of 10. Subsequently, the combined group joins the third cluster, which includes Zone-2. Finally, the fourth cluster, consisting of Zone-5, merges with the third cluster to form a single cohesive cluster at an approximate distance of 25.

For PM₁₀ concentrations, three clusters were identified. The first cluster comprises Zone-1, Zone-6, and Zone-2. This cluster merges with the second cluster, which includes Zone-4, Zone-7, and Zone-3, at a distance of approximately 13. The merged clusters then join with the third cluster, which includes Zone-5, forming a unified cluster at an approximate distance of 25.

5. CONCLUSION

The study determined that the average concentrations of $PM_{2.5}$ and PM_{10} at 37 educational institutions within the Dhaka South City Corporation were $68.82 \mu\text{g}/\text{m}^3$ and $113.95 \mu\text{g}/\text{m}^3$, respectively. The results show variations in $PM_{2.5}$ levels across different zones, ranked in descending order as follows: Zone-5 ($116.00 \mu\text{g}/\text{m}^3$), Zone-2 ($92.50 \mu\text{g}/\text{m}^3$), Zone-6 ($69.38 \mu\text{g}/\text{m}^3$), Zone-3 ($56.86 \mu\text{g}/\text{m}^3$), Zone-1 ($52.25 \mu\text{g}/\text{m}^3$), Zone-4 ($47.75 \mu\text{g}/\text{m}^3$), and Zone-7 ($47.00 \mu\text{g}/\text{m}^3$). Additionally, $PM_{2.5}$ and PM_{10} concentrations were found to be 1.78 and 1.03 times higher than the National Ambient Air Quality Standards (NAAQS) levels of $65 \mu\text{g}/\text{m}^3$ and $150 \mu\text{g}/\text{m}^3$, respectively. The average ratio of $PM_{2.5}$ to PM_{10} indicated that $PM_{2.5}$ constituted 60.41% of the PM_{10} mass.

The study also found that $PM_{2.5}$ and PM_{10} dispersion varied by land use, with Zone-5 showing the maximum range, followed by Zone-2, while Zone-7 and Zone-1 had the lowest ranges. The coefficient of variation for $PM_{2.5}$ and PM_{10} was highest in Zone-5 and Zone-4, respectively, and lowest in Zone-7 and Zone-1. Whisker-box plots revealed that Zone-4 exhibited the highest dispersion of $PM_{2.5}$ with a normally skewed distribution, whereas Zone-5 had the highest PM_{10} values with a positively skewed distribution. Zone-4 and Zone-3 showed normal to negatively skewed PM_{10} values. Significant variability in $PM_{2.5}$ concentrations was observed between and within land uses, with p-values less than 0.05. Priorities the implementation of consistent air quality monitoring systems at educational institutions, particularly in areas with high levels of $PM_{2.5}$ and PM_{10} concentrations, to effectively monitor pollution patterns and trends. Implementing more stringent restrictions and improving pollution control systems can effectively decrease the release of particulate matter, especially in heavily polluted areas like Zone-5 and Zone-2. In addition, the implementation of more green spaces in the vicinity of educational institutions can contribute to the absorption of pollutants and enhance the overall air quality. It is crucial to increase awareness among students, faculty, and the community on the health hazards linked to air pollution. This will promote the adoption of practices that reduce exposure, especially during periods of high pollution. Moreover, the act of evaluating and revising land use and urban planning regulations can effectively reduce pollutant origins and improve air quality, especially in areas with high population density and industrial activity. Finally, further investigation is required to determine the origins of particulate matter and comprehend the influence of various land uses on air quality.

This will facilitate the creation of focused interventions and policies.

6. REFERENCES

- [1] World Health Organization (WHO) Results Report 2020-2021. <https://www.who.int/.../who-results-report-2020-2021>.
- [2] I. Manisalidis, E. Stavropoulou, A. Stavropoulos, and E. Bezirtzoglou, "Environmental and Health Impacts of Air Pollution: A Review", *Front. Public Health, Sec. Environmental health and Exposome*, 2020; Volume 8 - 2020.
- [3] A. Juginović, M. Vuković, I. Aranza, and V. Biloš., "Health impacts of air pollution exposure from 1990 to 2019 in 43 European countries," *Sci Rep* 2021, 11. 22516.
- [4] The World Bank Annual Report 2018. <https://documents.worldbank.org/en/publication...>
- [5] World Air Quality Index (AQI) Ranking (IQAir) report 2023. <https://www.iqair.com/world-air-quality-ranking>
- [6] M. A. Nahian, T. Ahmad, I. Jahan, N. Chakraborty, Q. Nahar, and P. K. Streatfield, "Poor Air Quality and Pregnancy Outcome in Dhaka, Bangladesh," *The Journal of Climate Change and Health*, 2022, 100187.
- [7] O. B. Dipta, M. M. Hassan, and M. Hasan, "A study of high-rise building construction practice in Bangladesh," *AIP Conference Proceedings*, 2023, 2713(1).
- [8] B. A. Begum, M. Nasirs, S. Randal, B. Sivertsen and P. K. Hopke, "Identification and Apportionment of Sources from Air Particulate Matter at Urban Environments in Bangladesh" *British Journal of Applied Science & Technology*, 2014, 4(27), 3930-3955.
- [9] M. A. Fattah, M. E. U. Zannat, S. R. Morshed, and A. T. Roni, "Modeling the impact of motorized vehicles' activities on emissions and economic losses in a fast-growing developing city, Dhaka, Bangladesh," *Journal of Transport & Health*, 2022, Volume 25, June 2022, 101377.
- [10] J. Currie, J. S. G Zivin, J. Mullins, and M. J. Neidell "What do we know about short and long-term effects of early life exposure to pollution?" *NBER Work. Pap.* 2013, 6, pp. 217-247.
- [11] A. Mizen, J. Lyons, A. Milojevic, R. Doherty, P. Wilkinson, D. Carruthers, A. Akbari, I. Lake, G. A. Davies, M. Al Sallakh, R. Fry, L. Dearden, and S. Rodgers, "Impact of air pollution on educational attainment for respiratory health treated students: A cross sectional data linkage study" *Health & Place*, 2020, 63, [102355].
- [12] D.P. Skoner "Allergic rhinitis: definition, epidemiology, pathophysiology, detection, and diagnosis," *J. Allergy Clin. Immunol.*, 2001, 108, pp. 2-8.
- [13] S. Ahmed, M. S. Islam, and M. M. Rahman, "Assessment of PM₁₀ concentration near schools in Dhaka City, Bangladesh," *Environmental Science and Pollution Research*, 2020, 27(20), 25347-25357.