

Review Article

Agronomic interventions for alleviating stress and optimizing productivity of field crops: A review

Abstract

Water scarcity is the most significant concern facing agriculture today. Agriculture currently utilizes more than 70% of the world's freshwater, with much of it lost through evaporation, leaching and runoff. Drought escape and drought tolerance are the two significant mechanisms for plant growth under water-stressed conditions. Drought, a primary abiotic factor, limits crop productivity. It is a climatic phenomenon characterized by a prolonged lack of rainfall, resulting in moisture loss from the soil and a lack of water potential in plant tissues. It prevents the crop from obtaining its potential yield and substantially reduces crop production. Field crops rely on rainfall and are vulnerable to unpredictable drought stress throughout their vegetative and reproductive growth cycle. Drought stress is typical during the flowering stage in most crops, resulting in lower yield when cultivated with scarce rainfall. It can be reduced through agronomic means, such as mulching, tillage, intercropping and nutrient management, as well as chemical measures, such as the use of soil additives, herbal hydrogel (*Gond katira*), foliar spray of salicylic acid and potassium nitrate. Herbal hydrogel helps in reducing the effect of drought stress while increasing plant growth and performance. Foliar application of salicylic acid boosted bioactive chemical synthesis in the presence of water deficiency. Potassium nitrate promotes water efficiency, pod length and drought tolerance.

Key words: Field crops, drought stress, salicylic acid, herbal hydrogel (*Gond katira*), potassium-nitrate

Introduction

Agriculture is a cause and a victim of water scarcity. The excessive use and degradation of water resources are creating a threat to the sustainability of livelihoods dependent on water and agriculture (Rasul, 2023). There is a need for actions and

strategies that holistically address the interlinkages among water scarcity, agricultural production, food security and climate change. Water stress is considered a worldwide problem and one of the most and major abiotic stresses. Over 25% of the world's agricultural lands is now affected and suffering from water stress (Abido and Zsombik, 2018). Among all abiotic stresses, drought is considered the most damaging natural disaster due to its prolonged and extensive socioeconomic impacts (Ahmadalipouret al., 2019). About 330 million people in 2.5 lakh villages of 11 states in India were affected by drought in 2016 due to deficit rainfall during two consecutive years (Jayan, 2019). Drought stress causes a broad range of physiological changes and impairments of metabolic processes, which result in accumulation of reactive oxygen species (Abid et al., 2018; Lawlor and Cornic, 2002, Qaseem et al., 2019). The effect of water stress is more evident at vegetative stage than at reproductive stage (Ratnasekera and Subhashi, 2015). Water stress during vegetative phase limits root growth and reduces plant size, leaf area, accumulation of dry matter in pods, grain yield and harvest index (Rasul, 2023).

Stress mitigation strategies in field crops

Agronomic practices

The moong bean varieties, *i.e.*, Swarnaprabha and Kattamodan, with less leaf rolling, better drought recovery ability and relative water content, increased membrane stability index, osmolyte accumulation and antioxidant enzyme activities pointed toward their degree of tolerance to drought stress (Swapna and Shylaraj, 2017). Better performance due to higher water retention, photosynthetic performance and osmoregulation capacity was observed in legumes (Zhang et al., 2019).

Significant improvement in water-use efficiency was observed under conservation tillage over conventional tillage (Busari et al., 2015; Johnson et al., 2018). Tillage manages plant residue by incorporating into the soil or retaining on top layer to reduce erosion and to improve physical condition of the soil so that the soil can absorb the rainwater easily and soil erosion can be minimized to permit root elongation and proliferation due to low-density soil (Lundy et al., 2015). Conservation tillage is the method of soil tilling, in which, at least 30% of soil surface is covered with crop residues, which prevents from drought stress (Ali et al., 2016). Conservation tillage practices increased water use efficiency by 19.1-28.4 and 10.1-23.8% in wheat and maize, respectively (Shao et al., 2016). Similarly, under zero tillage, the crops save 20-30% water (Bhan and Behera, 2014).

Rapid and healthy crop establishment leads to strong root development, which minimizes the effects of future drought stress (Vance *et al.*, 2014). Under dry conditions, early seeding time is a key measure to match plant demand with water availability (Bodner *et al.*, 2015). Higher grain yield was observed under early sowing in comparison of late sowing during *Kharif* season (Shrestha *et al.*, 2016; Prasad *et al.*, 2017; Varma *et al.*, 2014).

Mulching enhances water-use efficiency by enhancing infiltration rate (Ahmad *et al.*, 2015), and reducing evaporation loss, runoff and temperature fluctuation increases drought tolerance (Ranjan *et al.*, 2017). Organic mulches act as a barrier to runoff and intercept raindrops, protecting the soil from splashing, particle detachment and clogging of surface soil pores, thereby contributing to the minimization of soil erosion (Prosdocimi *et al.*, 2016). Crops give higher yield with mulching under drought stress (Teameet *et al.*, 2017). Mulching can improve water-use efficiency by 10-20% (Kazemia and Safaria, 2018). Mulches suppress the weeds growth mainly by restricting the light penetration into the soil and thus improve water availability to crop plants under drought conditions (Tyagi *et al.*, 2020).

Intercropping in definite ratio may be followed in rain-fed areas to have minimum risk against total crop failure and also for better moisture utilization (Gautam and Bana, 2014). Intercropping is generally recommended for rain-fed crops to get stable yield (Singh *et al.*, 2014). Intercropping enhances soil water conservation and reduces runoff (Sharma *et al.*, 2017), increases water use efficiency (Hu *et al.*, 2017), and improves crop yield and the yield per unit of water supplied (Chen *et al.*, 2018). Both the wheat cultivars (Yecora E and Elissavet), the pea cultivar Olympos and their intercrops indicated high adaptation capacity to rainfed conditions, whereas, the pea cultivar Isard and its intercrop performed better under irrigation conditions, therefore, the intercropping of wheat with pea uses water resources of the environment more efficiently and could be used under dry land conditions for higher yield (Pankouet *et al.*, 2021).

Application of nitrogen either through soil or through foliar spray is an important strategy to alleviate the adverse effect of drought (Ahmad *et al.*, 2014). Higher yield of crop with foliar spray of potassium chloride 2% + sodium selenite 0.4% under drought conditions (Dewanganet *et al.*, 2017). Higher yield and profit with the application of potassium fertilizer were evident in many crops (Hussain

et al., 2017). Potassium improves many physiological processes by the regulation of turgor pressure, photosynthesis, translocation of assimilates to various organs and enzymes activation and thus improves drought tolerance ability of plants (Raja *et al.*, 2017). Maintenance of adequate potassium nutrition to plants has been found critical to mitigate drought stress (Khan *et al.*, 2018).

Higher yield and water-use efficiency could be obtained with supplemental application of irrigation at critical stages of crop growth (Abbas *et al.*, 2014). In arid crops, life-saving irrigation wherever necessary can be supplied by ground water storage or by storing rain water during heavy rainfall (Yadav *et al.*, 2014). Supplementary irrigation, especially during long dry spell after rainy months, could possibly alleviate moisture stress in growing crops (Praharaj *et al.*, 2016).

Micro-irrigation techniques like sprinkler and drip should be promoted to reduce the risk of yield reduction (Ashoka *et al.*, 2015). Drip irrigation can substantially improve water use efficiency by minimizing loss of water through runoff and evaporation (Jha *et al.*, 2016). In drip irrigation system, water is slowly applied in the form of drop either on soil surface or directly into the root zone, through a network of valves, pipes, tubing and emitters (Kumar *et al.*, 2016). A significant water saving was recorded in chickpea through drip irrigation on account of higher application efficiency and water supply to plant root zone with a discharge rate not more than infiltration rate of soil (Muniyappa *et al.*, 2017). Around 50% saving of irrigation water along with higher productivity was observed under drip irrigation system (Mostafa *et al.*, 2018). Sprinkler irrigation has resulted in 16.22% higher yield and 30.76% higher water productivity (Sharma *et al.*, 2018).

Chemical measures

Hydrogel application results in enhancement of soil physical, chemical and biological properties with positive effects on plant growth and development (Abobatta, 2018). Plants can survive and sustain their life cycle through hydrogel conditioning in areas, where the shortage of water is a serious issue (Tu *et al.*, 2018). Hydrogel (super absorbent) is one of the most popular and commercially available soil additives used to reduce water runoff, infiltration rate in field and irrigation requirement of several crops by improving water holding capacity (Laxmi S., 2019). It is also known to increase plant survival time under drought conditions (Jerszurkiet *al.*,

2017; Ayangbenro and Babalola, 2020). Water loss through evaporation in hydrogel amended soil is lower than the soil with no hydrogel amendment (Saha, 2020).

Under water stress conditions, there was an increase in root development with the rise in K content in mung bean (Nisha *et al.*, 2014). Potassium indirectly promotes the transfer of photosynthates gradually towards the roots for nodule utilization, increasing root nodules under water stress conditions (Suryapani and associates, 2014). The effect of potassium application is evident in enhancing the tolerance of plants under water stress conditions (Zain and Ismail, 2016). The key functions of salicylic acid in drought stress alleviation include promoting seedling growth, enhancing plant antioxidant capacity, regulating plant water balance, promoting the expression of stress-related genes in plants and regulating plant physiological metabolism (Aiman *et al.*, 2018). As a crucial chemical, salicylic acid has been found to alleviate drought stress in plants to some extent (Munsif *et al.*, 2022). Salicylic acid can enhance plant drought tolerance by regulating root growth, improving leaf water use efficiency, stabilizing cell membranes and modulating the antioxidative enzyme system. Meanwhile, relevant studies also indicated that the biosynthesis of salicylic acid also exerted certain regulatory effects on normal plant growth and development, which is beneficial to the enhancement of plant tolerance to various kinds of stress (Rekhteret *et al.*, 2019; Lefevere *et al.*, 2020; Peng *et al.*, 2021; Ullah *et al.*, 2023). In addition, other relevant studies have also demonstrated the important regulatory role of salicylic acid in plant drought tolerance (Alam *et al.*, 2022; Das *et al.*, 2023; Duvnjak *et al.*, 2023; Guo *et al.*, 2023; Moustakas *et al.*, 2023; Tanveer *et al.*, 2023).

Impact of stress mitigation strategies on plant growth and yield parameters

Salicylic acid controls a variety of physiological functions in plants, including photosynthesis, nitrogen metabolism, glycine betaine synthesis and antioxidant synthesis, which ensures plant resilience to abiotic stress management (Khan *et al.*, 2014; Miura and Tada, 2014). Fertilizer containing potassium higher soluble sugar concentration may act as a solute for osmotic regulation and/or a substrate for root-based synthesis of proteins and polysaccharides, which in turn promote the development of entire plants under water stress (Semida *et al.*, 2014). Plants that receive adequate potassium are more resistant to drought stress, water shortage conditions and water use efficiency, all of which encourage crop development and yield (Hassan *et al.*, 2017). Under water deficit, foliar application of salicylic acid

improved the leaf relative water content and photosynthetic pigments in crops (Razmi *et al.*, 2017). Salicylic acid preserves water in plant cells, which stimulates enzyme activity under stressful conditions, improving metabolism and yield. This could be the reason why salicylic acid boosts yield in stressed plants (Ezzo *et al.*, 2018). Application of recommended dose of fertilizer + potassium @ 30 kg ha⁻¹ + two foliar spray of KNO₃ @ 1% at flowering and pod formation stage showed synergistic effects on nutrients (N, P and K) uptake. It was also discovered that providing potassium to cultivars under water stress boosted soil fertility (Bhadane *et al.*, 2019). The application of salicylic acid increased the activity of antioxidant enzymes, such as ascorbate peroxidase and chloramphenicol acetyltransferase, which might have helped in alleviating the negative effect of water stress on plants (Shemiet *et al.*, 2021; Sultana *et al.*, 2024). Foliar application of salicylic acid could be a promising strategy to mitigate water stress in plants (Sultana *et al.*, 2024) since it can manage metabolic pathways and enhance the plants' defense against water deficits (Ghahremani *et al.*, 2023).

Salicylic acid made the plant more resistant to drought, and it might be utilized to boost and stabilize crop output under stressful conditions (Sharma *et al.*, 2017). Under salt and drought stress, the exogenous application of salicylic acid promoted both root and stem growth (Movahhedi-Dehnavi *et al.*, 2019). Plant development is influenced by salicylic acid in a number of ways, including seed germination, induction of flowering, root growth and ion uptake under water stress conditions (Bagautdinova *et al.*, 2022; Liu *et al.*, 2022).

Foliar application of salicylic acid is an efficient technique that can decrease the detrimental effects of drought stress and boost crop output (Kazemi, 2014). Salicylic acid helps in synthesis of osmotic substances, preserving intracellular oxidative equilibrium and enhancing the control of mineral nutrient intake and helps the plants to develop resistance against abiotic stressors (Khan *et al.*, 2015).

Under drought conditions, seed priming by potassium nitrate increased the leaf nitrate reductase activity of the seedlings (Kumar *et al.*, 2014). The application of 2% KNO₃ in fodder legume crop resulted in 26.7 and 27.5% higher pod and grain yield, respectively than the control (Kumar *et al.*, 2014). A foliar spray of KNO₃ at 150 ppm 25 DAS increased total chlorophyll content by 26.5 and 3.3% at 40 and 50 DAS, respectively in legume crops under drought conditions (Mubeen *et al.*, 2014). Greater

plant biomass (1239 kg ha^{-1}) and height (48.1 cm) was recorded with foliar application of salicylic acid as compared to control (Farjam *et al.*, 2015). Seed coating with herbal hydrogel (Gond-Katira) resulted in increased plant height and tillers m^{-2} (Lather *et al.*, 2015). Foliar application of salicylic acid and potassium for reducing drought stress in legumes (Majeed *et al.*, 2016). Foliar spray of potassium nitrate 1% in legumes under water deficiency conditions significantly increased the plant height, leaf area, shoot dry weight and relative leaf water content as compared to control (Rao *et al.*, 2016). Foliar treatment of salicylic acid 75 ppm + 2% urea during flower initiation considerably improved the plant height, number of branches, dry matter accumulation and chlorophyll content under water stress conditions (Sarita *et al.*, 2020). The application of Pusa and herbal hydrogel increased the grain, straw and biological yield by 6.65-13.69, 6.90-14.41 and 6.79-14.09%, respectively as compared to control (Kumar *et al.*, 2019; Kumar *et al.*, 2020). Foliar spraying of potassium nitrate 0.5% resulted in increased seed output (865 kg ha^{-1}) in legume crops as compared to water spray (Kumar *et al.*, 2020a). The application of hydrogel 3 kg ha^{-1} resulted in a greater percentage of Sennoside A (1.78%) and Sennoside B (1.69%) in pods as compared to other treatments under drought conditions (Jnanasha *et al.*, 2021).

Salicylic acid treatment significantly increased growth, photosynthesis, yield and the activity of antioxidant enzymes, *viz.* catalase, peroxidase and superoxide dismutase in chickpea plants exposed to 0, 25, or 50 mg Cd per kg of soil, however, the treatment did not prove to be fruitful in alleviating completely the stress generated by 100 mg Cd per kg of soil (Alyemeni *et al.*, 2014). The maximum response was generated in plants sprayed with $10^{-5} \text{ mol L}^{-1}$ of salicylic acid, showing a statistically significant increase in net photosynthesis (46.92%) over that of the control under drought stress conditions (Hayat *et al.*, 2014). Combined spray of 2% urea and 0.25% multiplex increased plant height, number of branches and dry matter per plant under water stress conditions (Ganga *et al.*, 2014). Foliar application of potassium nitrate @ 2% + boric acid 50 ppm + zinc sulphate 1% at 30 and 60 DAS was found to be superior in increasing seed yield under water stress conditions (Gowthami and Rao, 2014). Hydrogel had a positive effect on photosynthesis by gradually pumping water into the plant (Tango *et al.*, 2014). The treated plants with salicylic acid survive more, exhibit higher relative growth rate and more leaf area and dry mass production under water stress conditions (Boukraâ *et al.*,

2015). Sprays of KNO_3 1.0% twice at 45 and 60 DAS improved all the growth parameters, viz. plant height (cm), number of branches plant^{-1} , nodules plant^{-1} and total dry matter plant^{-1} in crop under water stress (Hiwale, 2015). Application of salicylic acid has a positive effect in ameliorating the oxidative damages in plant and increases the yield (Hossain *et al.*, 2015). Foliar application of KNO_3 2% four times improved all the growth parameters as compared to control (Kumar and Srivastava, 2015). Salicylic acid can reduce the sensitivity of plants to environmental stresses through the regulation of antioxidant defence system, transpiration rate, stomatal movement and photosynthetic rate (Nazar *et al.*, 2015). Application of exogenous salicylic acid could help in reducing the adverse effects of drought stress and might have a key role in providing tolerance to stress by promoting growth and accumulation of proline, soluble sugars and photosynthetic pigments in plant leaves (Vaisnadi *et al.*, 2015). The increase in number of seeds per pod under increasing levels of salicylic acid might be due to increased mobilization of metabolites to the reproductive sinks (Khatun *et al.*, 2016). Application of salicylic acid resulted in a positive effect by protecting plants against oxidative damage caused by drought stress (Najafabadi and Ehsanzadeh, 2017; Wang *et al.*, 2019; Sankari *et al.*, 2019). Hydrogel retains its structural integrity and absorbs water from the environment, helping under drought conditions (Montesano *et al.*, 2015; Domalik-Pyzik *et al.*, 2019; Qu and Luo, 2020). Hydrogel also increased the plants' life span by approximately 9-14 days (Song *et al.*, 2020; Shankarappa *et al.*, 2020). Hydrogel application provided 22-45% more plant growth under water stress conditions (Nassaj-Bokharaei *et al.*, 2021). Addition of hydrogel to soil (in different types of soil and with different hydrogel amounts) increases the water holding capacity of the soil by 56 to 81% under water stress (Saha *et al.*, 2021). Hydrogel application under water stress conditions increased plant height, stem diameter, leaf length and vegetative period along with overall growth of plant (Albalasmeh *et al.*, 2022; Radian *et al.*, 2022; Prisa and Guerrini, 2022; Watcharamule *et al.*, 2022). Hydrogel increased the yield of some crops as compared to the control (Rajanna *et al.*, 2022). The leaf area and calyx yield of *Hibiscus sabdariffa* L. (Besharati *et al.*, 2022) and vegetative growth of plants (Alshallash *et al.*, 2022) increased even under water stress conditions due to hydrogel applications, which increased leaf area, as well as yield of the crops by 35 to 60% under water stress and encouraged the development of *Brassica rapa* ssp. *chinensis* var. *parachinensis* (Zhu *et al.*, 2022a,b).

Salicylic acid is of great potential to improve photosynthesis and chlorophyll content in wheat (Khalvandi *et al.*, 2021). In *Camellia oleifera*, salicylic acid level declines during drought stress, exhibits stronger antioxidant capacity, water regulation ability and drought stress protection (Guo *et al.*, 2023). Substantial reduction of drought influence and enhancement of grain yield and water use efficiency by co-application of K⁺ and salicylic acid and regulation of stress response in wheat (Munsif *et al.*, 2022; Duvnjak *et al.*, 2023). Salicylic acid mitigates the oxidative damage and alleviates water deficiency stress in potato (Acevedo *et al.*, 2023). Salicylic acid helps in regulation of stomatal aperture and adaptation to soil salinity and drought stress in rice (Xu *et al.*, 2023).

Conclusion

The reviewed studies collectively emphasize the crucial role of potassium nitrate, salicylic acid and other nutrients in enhancing crop productivity and quality under drought stress. Potassium nitrate application leads to improved nutrient content in crops and soil fertility, mitigates deficiencies and boosts seed yield and quality. Salicylic acid treatment counteracts drought stress and promotes growth and yield. Additionally, foliar application of various compounds, such as urea and KNO₃, showcased significant effects on growth parameters and yield. These findings underscore the importance of targeted nutrient applications for optimal crop performance, demonstrating the potential for advanced agricultural practices to enhance agricultural productivity sustainably. In conclusion, stress management practices in agriculture are pivotal for enhancing crop productivity and quality. The utilization of herbal hydrogel goud-katira (*Tragacanth katira*) has shown promise in mitigating various stressors. Additionally, incorporating foliar nutrition, such as urea and salicylic acid, as demonstrated by several researchers, can enhance plant resilience to stressors like heat or nutrient deficiencies. The use of innovative technologies like herbal hydrogel systems, offers a sustainable approach to address crop stress under moisture-limited conditions.

References

1. Abbas, S.H., Bhatti, A.A., Asif, M., Islam, Z., Sohail, M., Din, R.U., Qamar, M., Ali, M., Majeed, A. and Munir, M. (2014). Effect of supplemental irrigation on wheat

- water productivity under rainfed ecology of Pothohar, Pakistan. *Innov. J. Agri. Sci.*, **3**(1): 10-13.
2. Abid, M., Ali, S., Qi, L.K., Zahoor, R., Tian, Z., Jiang, D., Snider, J.L. and Dai, T. (2018). Physiological and biochemical changes during drought and recovery periods at tillering and jointing stages in wheat (*Triticum aestivum* L.). *Sci. Rep.*, **8**(1): 4615.
 3. Abido, W.A.E. and Zsombik, L., (2018). Effect of water stress on germination of some Hungarian wheat landraces varieties. *Acta Ecol. Sin.*, **38**: 422–428.
 4. Abobatta, W. (2018). Impact of hydrogel polymer in agricultural sector. *Adv. Agric. Environ. Sci.*, **1**: 59-64.
 5. Acevedo, A.F.G., Lacerda, V.R., Da Silva-Gomes, J.W., Avilez, A.A., Sarria, S.D., Broetto, F., Vieites, R.L. and De Souza-Guimarães, M.L.C. (2023). Foliar salicylic acid application to mitigate the effect of water deficiency on potato (*Solanum tuberosum* L.). *Plant Stress*, **7**: 100135.
 6. Ahmad, R., Waraich, E.A., Ashraf, M.Y., Ahmad, S. and Aziz, T. (2014). Does nitrogen fertilization enhance drought tolerance in sunflower?: A review. *J. Plant Nutr.*, **37**(6): 942-963.
 7. Ahmad, S., Raza, M.A.S., Saleem, M.F., Zahra, S.S., Khan, I.H., Ali, M., Shahid, A.M., Iqbal, R. and Zaheer, M.S. (2015). Mulching strategies for weeds control and water conservation in cotton. *J. Agric. Biol. Sci.*, **10**(8): 299-306.
 8. Ahmadalipour, A., Moradkhani, H., Castelletti, A. and Magliocca, N. (2019). Future drought risk in Africa: integrating vulnerability, climate change and population growth. *Sci. Total Environ.*, **662**: 672-686.
 9. Aiman, A. and Dhriti, K. (2018). Salicylic acid: Its physiological role and interactions. *Res. J. Pharm. Technol.*, **7**: 3171-3177.
 10. Alam, M.U., Fujita, M., Nahar, K., Rahman, A., Anee, T.I., Masud, A.A.C., Amin, A.K.M.R. and Hasanuzzaman, M. (2022). Seed priming upregulates antioxidant defense and glyoxalase systems to conferring simulated drought tolerance in wheat seedlings. *Plant Stress*, **6**: 100120.
 11. Albalasmeh, A.A., Mohawesh, O., Gharaibeh, M.A., Alghamdi, A.G., Alajlouni, M.A. and Alqudah, A.M. (2022). Effect of hydrogel on corn growth, water use efficiency and soil properties in a semi-arid region. *J. Saudi Soc. agric. Sci.*, **21**(8): 518-524.

12. Ali, A.B., Elshaikhb, N.A., Honga, L., Adamc, A.B. and Haofang, Y. (2016). Conservation tillage as an approach to enhance crops water use efficiency. *Acta Agric. Scandinavica, Section B-Soil Plant Sci.*, **67**(3): 252-262.
13. Alyemini, M.N., Hayat, Q., Wijaya, L. and Hayat, S. (2014). Effect of salicylic acid on the growth, photosynthetic efficiency and enzyme activities of leguminous plant under cadmium stress. *Notulae Botanicae Horti. Agrobotanici*, **42**(2): 440-445.
14. Ashoka, P., Kadasiddappa, M.M. and Sanjey, M.T. (2015). Enhancing water productivity through micro irrigation technologies in Indian agriculture. *Ann. Plant Soil Res.*, **17**(Special Issue): 601-605.
15. Ayangbenro, A.S. and Babalola, O.O. (2021). Reclamation of arid and semi-arid soils: The role of plant growth-promoting archaea and bacteria. *Curr. Plant Biol.*, **25**: 100173.
16. Bagautdinova, Z.Z., Omelyanchuk, N., Tyapkin, A.V., Kovrizhnykh, V.V., Lavrekha, V.V., and Zemlyanskaya, E.V. (2022). Salicylic acid in root growth and development. *Int. J. Mol. Sci.*, **23**(4): 2228.
17. Besharati, J., Shirmardi, M., Meftahizadeb, H., Aardakni, M.D. and Ghorbanpour, M. (2022). Changes in growth and quality performance of Roselle (*Hibiscus sabdariffa* L.) in response to soil amendments with hydrogel and compost under drought stress. *South Afr. J. Bot.*, **145**: 334-347.
18. Bhadane, D.T., Mane, S.S. and Sonawane, V.V. (2019). Effect of potassium application on growth yield and quality of chickpea cultivars. *Int. J. Chem. Studies*, **7**(6): 604-606.
19. Bhan, S. and Behera, U.K. (2014). Conservation agriculture in India-problems, prospects and policy issues. *Int. Soil Water Conserv. Res.*, **2**(4): 1-12.
20. Bodner, G., Nakhforoosh, A. and Kaul, H. (2015). Management of crop water under drought: A review. *Agron. Sustain Dev*, **35**: 401-442.
21. Boukraâ, D., Belabid, L., Benabdelli, K. and Bennabi, F. (2015). Implication of salicylic acid in chickpea growth to salt resistance. *Adv. Environ. Biol.*, **9**(27): 270-277.
22. Bukhsh, M.A.A.H.A., Ahmad, R., Iqbal, J., Maqbool, M.M., Ali, A., Ishaque, M. and Hussain, S. (2012). Nutritional and physiological significance of potassium application in maize hybrid crop production (Review Article). *Pak. J. Nutr.*, **11**: 187-202.

23. Busari, M.A., Kukal, S.S., Kaur, A., Bhatt, R. and Dulazi, A.A. (2015). Conservation tillage impacts on soil, crop and the environment. *Int. Soil Water Conserv. Res.*, **3**: 119-129.
24. Chen, G., Kong, X., Gan, Y., Zhang, R., Feng, F., Yu, A., Zhao, C., Wan, S. and Chai, Q. (2018). Enhancing the systems productivity and water use efficiency through coordinated soil water sharing and compensation in strip intercropping. *Sci. Rep.*, **8**: 1-11.
25. Das, D., Bisht, K., Chauhan, A., Gautam, S., Jaiswal, J.P., Salvi, P. and Lohani, P. (2023). Morpho-physiological and biochemical responses in wheat foliar sprayed with zinc-chitosan-salicylic acid nanoparticles during drought stress. *Plant Nano Biol.*, **4**: 100034.
26. Dewangan, S., Singh, R.P., Singh, M.K. and Singh, S. (2017). Effect of integrated nutrient management and drought mitigating practices on performance of rainfed chickpea (*Cicer arietinum*). *Indian J. agric. Sci.*, **87**(3): 301-305.
27. Domalik-Pyzik, P., Chlopek, J. and Pielichowska, K. (2019). Chitosen-based hydrogels: Preparation properties and applications. In: *Cellulose-Based superabsorbent Hydrogels* (Ed. Mondal, I.H.). Springer, New York, USA, pp. 1665-1693.
28. Duvnjak, J., Lončarić, A., Brkljačić, L., Šamec, D., Šarčević, H., Salopek-Sondi, B. and Španić, V. (2023). Morpho-Physiological and Hormonal Response of Winter Wheat Varieties to Drought Stress at Stem Elongation and Anthesis Stages. *Plants*, **12**: 418.
29. Ezzo, M., Ebtihal, M., Elhamid, A., Sadak, M.S. and Abdalla, A.M. (2018). Improving drought tolerance of moringa plants by using trehalose foliar treatments. *Biosci. Res.*, **15**(4): 4203-4214.
30. Farjam, S., Kazemi-Arbat, H., Siosemardeh, A., Yarnia, M. and Rokhzadi, A. (2015). Effects of salicylic and ascorbic acid applications on growth, yield, water use efficiency and some physiological traits of chickpea (*Cicer arietinum* L.) under reduced irrigation. *Leg. Res.*, **38**(1): 66-71.
31. Ganga, N., Singh, R.K., Singh, R.P., Choudhury, S.K. and Upadhyay, P.K. (2014). Effect of potassium level and foliar application of nutrient on growth and yield of late sown chickpea (*Cicer arietinum* L.). *Environ. Ecol.*, **32**(1A): 273-275.
32. Gautam, R.C. and Bana, R.S. (2014). Drought in India: its impact and mitigation strategies-a review. *Indian J. Agron.*, **59**(2): 179-190.

33. Ghahremani, Z., Alizadeh, B., Barzegar, T., Nikbakht, J., Ranjbar, M.E. and Nezamdoost, D. (2023). The mechanism of enhancing drought tolerance threshold of pepper plant treated with putrescine and salicylic acid. *Plant Stress*, **9**: 100199.
34. Gouthami, P. and RamaRao, G. (2014). Effect of foliar application of potassium, boron and zinc on growth analysis and seed yield in soybean. *Int. J. Food Agri. Vet. Sci.*, **4**(3): 73-80.
35. Guo, P.R., Wu, L.L., Wang, Y., Liu, D. and Li, J.A. (2023). Effects of drought stress on the morphological structure and flower organ physiological characteristics of *Camelliaoleifera* flower buds. *Plants*, **12**: 2585.
36. Hassan, M.U., Aamer, M., Chattha, M.U., Ullah, M.A., Sulaman, S., Nawaz, M. and Guoqin, H. (2017). The role of potassium in plants under drought stress: Mini review. *J. Basic Appl. Sci.*, **13**: 268-271.
37. Hayat, S., Hayat, Q., Alyemeni, M.N. and Ahmad, A., (2014). Cumulative effects of proline and salicylic acid on the cadmium-induced changes in *Cicer arietinum* L. *Fresenius Environ. Bull.*, **23**(2): 330-340.
38. Hiwale, R. (2015). Effect of foliar application of potassium nitrate on yield, growth and quality of soybean [*Glycine max* (L.) Merrill]. *Int. J. Agri. Sci.*, **7**(5): 516-519.
39. Hosmath, J.A., Biradar, D.P., Patil, V.C., Palled, Y.B. and Malligawad, L.H. (2014). Nutrient requirement of Bt cotton (*Gossypium hirsutum*). *Indian J. Agron.*, **59**(1): 133-138.
40. Hossain, M.I., Mannan, M.A. and Karim, M.A. (2015). Salicylic acid and gibberelic acid ameliorates the adverse effects of salinity on chickpea. *Bangladesh Agron. J.*, **18**(1): 81-88.
41. Hu, F., Feng, F., Zhao, C., Chai, Q., Yu, A., Yin, W. and Gan, Y. (2017). Integration of wheat-maize intercropping with conservation practices reduces CO₂ emissions and enhances water use in dry areas. *Soil Tillage Res.*, **169**: 44-53.
42. Hussain, S., Maqsood, M., Lal, R., Hussain, M., Sarwar, M.A, Bashair-Ullah, A.M. and Ulhaq, I. (2017). Integrated nutrient management strategies to alleviate drought stress in hybrid maize in Punjab, Pakistan. *Rom. Agric. Res.*, **34**: 233-242.
43. Jayan, T.V. (2019). Drought conditions persist in many states. The Hindu Business line. Accessed, 11.
44. Jerszurki, D., Couvreur, V., Maxwell, T., Silva, L.D.C.R., Matsumoto, N., Shackel, K., de Souza, J.L.M. and Hopmans, J. (2017). Impact of root growth and hydraulic

- conductance on canopy carbon-water relations of young walnut trees (*Juglans regia* L.) under drought. *Sci. Hort.*, **226**: 342-352.
45. Jha, A.K., Malla, R., Sharma, M., Panthi, J., Lakhankar, T., Krakauer, N.Y., Pradhanang, S.M., Dahal, P. and Shrestha, M.L. (2016). Impact of irrigation method on water use efficiency and productivity of fodder crops in Nepal. *Climate*, **4**(4): 1-13.
46. Jha, S.K., Ramatshaba, T.S., Wang, G., Liang, Y., Gao, H.L.Y. and Aiwang, D. (2019). Response of growth, yield and water use efficiency of winter wheat to different irrigation methods and scheduling in North China plain. *agric. Water Mgt.*, **217**: 292-302.
47. Jnanasha, A.C., Kumar, A. and Lal, R.K. (2021). Hydrogel application improved growth and yield in Senna (*Cassia angustifolia* Vahl). *Industrial Crops and Products*, **174**: 114175.
48. Johnson, Y.K., Ayuke, F.O., Kinama, J.M. and Sijali, I.V. (2018). Effects of tillage practices on water use efficiency and yield of different drought tolerant common bean varieties in Machakos county, Eastern Kenya. *Am. Sci. Res. J. Eng. Technol. Sci.*, **40**(1): 217-234.
49. Kazemi, M. (2014). Effect of foliar application with salicylic acid and methyl jasmonate on growth, flowering, yield and fruit quality of tomato. *Bull. Env. Pharmacol. Life Sci.*, **3** (2): 154-158.
50. Kazemia, F. and Safaria, N. (2018). Effect of mulches on some characteristics of a drought tolerant flowering plant for urban landscaping. *Desert*, **23**(1): 75-84.
51. Khalvandi, M., Siosemardeh, A., Roohi, E. and Keramati, S. (2021). Salicylic acid alleviated the effect of drought stress on photosynthetic characteristics and leaf protein pattern in winter wheat. *Heliyon*, **7**(1): e05908.
52. Khan, A., Pan, X., Najeeb, U., Tan, D.K.Y., Fahad, S., Zahoor, R., Luo, H. (2018). Coping with drought: stress and adaptive mechanisms, and management through cultural and molecular alternatives in cotton as vital constituents for plant stress resilience and fitness. *Biol Res.*, **47**: 1-17.
53. Khan, M.I.R., Asgher, M. and Khan, N.A. (2014). Alleviation of salt-induced photosynthesis and growth inhibition by salicylic acid involves glycine betaine and ethylene in mung bean (*Vigna radiata* L.). *Plant Physiol. Biochem.*, **80**: 67-74.
54. Khan, M.I.R., Fatma, M., Per, T.S., Anjum, N.A. and Khan, N.A. (2015). Salicylic acid-induced abiotic stress tolerance and underlying mechanisms in plants. *Front. in Plant Sci.*, **6**: 462.

55. Khatun, S., Roy, T.S., Haque, M.N. and Alamgir, B. (2016). Effect of plant growth regulators and their time of application on yield attributes and quality of soybean. *Int. J. Plant Soil Sci.*, **11**(1): 1-9.
56. Kumar, B., Lamba, J.S., Dhaliwal, S.S., Sarlach, R.S. and Ram, H. (2014). Exogenous application of bio-regulators improves grain yield and nutritional quality of forage cowpea (*Vigna unguiculata*). *Int. J. agric. Biol.*, **16**: 759-765.
57. Kumar, J. and Srivastava, L.H. (2015). Growth and yield attributes, yield, fibre quality and economics of *Hirsutum* cotton as influenced by foliar application of KNO₃. *Plant Arch.*, **15**(2): 1147-1149.
58. Kumar, M. and Singh, B. (2020). Evaluation of irrigation levels and hydrogels on the growth, yield and profitability of wheat. *Int. J. Curr. Sci.*, **8**(4): 1649-1654.
59. Kumar, R., Trivedi, H., Yadav, R., Das, B. and Bist, A.S. (2016). Effect of drip irrigation on yield and water use efficiency on brinjal (*Solanum melongena*) cv. Pant Samrat. *Int. J. Eng. Sci. Res. Tech.*, **5**(10): 7-17.
60. Kumar, S., Bose, B. and Pradhan, N. (2014). Potassium nitrate priming affects the activity of nitrate reductase and chlorophyll content in late sown sesame (*Sesamum indicum* L.). *Trends Biosci.*, **7**(24): 4466-4470.
61. Kumar, S., Sharma, P.K., Yadav, M.R., Sexena, R., Gupta, K.C., Kumar, R. and Yadav, H.L. (2019). Effect of irrigation levels and moisture conserving polymers on growth, productivity and profitability of wheat. *Indian J. agric. Sci.*, **89**(3): 509-14.
62. Kumar, Y.S., Hemalatha, S., Chandrika, V., Latha, P. and Sagar, G.K. (2020)a. Growth and yield of summer black gram (*Vigna mungo* L.) as influenced by moisture stress and foliar nutrition. *A. P. J. agric. Sci.*, **6**: 111-114
63. Lather, V.S., Ashwani, K., Chopra, N.K., Choudhary, D., Yadav, R.N. and Seth, R. (2015). Novel herbal hydrogel *Tragacanth-Katiragel* and farmer's friendly seed priming-hydrogel coating technology for water saving, making agriculture sustainable and resilient to climatic variability. *Int. J. Trop. Agri.*, **33**(2): 1167-1171.
64. Laxmi, S., Chanu, P.H., Rani P., Rai, S., Prasad, S.K. and Singh, R.K. (2019). Effect of hydrogel on soil moisture stress. *J. Pharmacogn. Phytochem.*, **8**(6): 316-320.
65. Lefevre, H., Bauters, L. and Gheysen, G. (2020). Salicylic acid biosynthesis in plants. *Front. Plant Sci.*, **11**: 338.
66. Liu, Z., Ma, C., Hou, L., Wu, X., Wang, D., Zhang, L., Liu, P. (2022). Exogenous SA affects rice seed germination under salt stress by regulating Na⁺/K⁺ balance and endogenous GAs and ABA homeostasis. *Int. J. Mol. Sci.*, **23**(6): 3293.

67. Lundy, M.E., Pittelkow, C.M., Linquist, B.A., Liang, X., van Groenigen, K.J., Lee, J., Six, J., Venterea, R., van Kessel, C. (2015). Nitrogen fertilization reduces yield declines following no-till adoption. *Field Crops Res.*, **183**: 204-210.
68. Majeed, S., Akram, M., Latif, M., Ijaz, M. and Hussain, M. (2016). Mitigation of drought stress by foliar application of salicylic acid and potassium in mungbean (*Vigna radiata* L.). *Leg. Res.*, **39**(2): 208-214.
69. Miura, K. and Tada, Y. (2014). Regulation of water, salinity, and cold stress responses by salicylic acid. *Front. Plant Sci.*, **5**: 4.
70. Montesano, F.F., Prente, A., Santamaria, P., Sannino, A. and Serio, F. (2015). Biodegradable superabsorbent hydrogel increases water retention properties of growing media and plant growth. *Agri. agric. Sci. Procedia*, **4**(10): 451-458.
71. Mostafa, H., El-Nady, R., Awad, M. and El-Ansary, M. (2018). Drip irrigation management for wheat under clay soil in arid conditions. *Ecol. Eng.*, **121**: 35-43.
72. Movahhedi-Dehnavi, M., Behzadi, Y., Niknam, N. and Mohtashami, R. (2019). Salicylic acid mitigates the effects of drought and salinity on nutrient and dry matter accumulation of linseed. *J. Plant Process Func.*, **8**(31): 31-44.
73. Mubeen, B.R., Shamim, M., Khan, A.H., Singh, A.K., Singh, S.P., Yadav, R. K. and Singh, U. (2014). Response of mungbean (*Vigna radiata* L.) to rhizobium inoculation and growth regulators on biochemical changes and yield. *Environ. Ecol.*, **33**: 105-109.
74. Muniyappa, M., Ramachandrappa, B.K., Nagaraju. and Sathish, A. (2017). Studies on the influence of depth and interval of drip irrigation on yield, water use efficiency and economics of chickpea (*Cicer arietinum* L.). *Int. J. Pure. App. Biosci.*, **5**(1): 771-776.
75. Munsif, F., Shah, T., Arif, M., Jehangir, M., Afridi, M.Z., Ahmad, I., Jan, B.L. and Alansi, S. (2022). Combined effect of salicylic acid and potassium mitigates drought stress through the modulation of physio-biochemical attributes and key antioxidants in wheat. *Saudi J. Biol. Sci.* **29**: 103294.
76. Nassaj-Bokharaei, S., Motesharezadeh, B., Etesami, H. and Motamedi, E. (2021). Effect of hydrogel composite reinforced with natural char nanoparticles on improvement of soil biological properties and the growth of water deficit-stressed tomato plant. *Ecotoxicol. Environ. Safety*, **223**: 112576.
77. Nazar, R., Umar, S., Khan, N.A., and Sareer, O. (2015). Salicylic acid supplementation improves photosynthesis and growth in mustard through changes in proline accumulation and ethylene formation under drought stress. *South Af. J. Bot.*, **98**: 84-94.

78. Nisha, K. and Narender, S. (2014). Assessing the effects of applied potassium on selected *Vigna radiata* L. genotypes under water deficit. *Int. J. Adv. Res.*, **2**(10): 369-385.
79. Pankou, C., Lithourgidis, A. and Dordas, C. (2021). Effect of irrigation on intercropping systems of wheat (*Triticum aestivum* L.) with pea (*Pisum sativum* L.). *Agron.*, **11**(2): 283.
80. Peng, Y., Yang, J., Li, X. and Zhang, Y. (2021). Salicylic Acid: Biosynthesis and Signaling. *Annu. Rev. Plant Biol.*, **72**: 761-791.
81. Praharaj, C.S., Singh, U., Singh, S.S., Singh, N.P. and Shivay, Y.S. (2016). Supplementary and life-saving irrigation for enhancing pulses production, productivity and water-use efficiency in India. *Indian J. Agron.*, **61**(4th IAC special issue): S249-S261.
82. Prasad, G., Chand, M., Kumar, P. and Rinwa, R.S. (2017). Performance of maize (*Zea mays* L.) hybrids with respect to growth parameters and phenological stages under different sowing dates in kharif season. *Int. J. Curr. Microbiol. App. Sci.*, **6**(10): 5079-5087.
83. Prisa, D. and Guerrini, G. (2022). Hydrogel application in the germination and growth of *Zea mays* and solanum (lycopersium) seedling. *GSC Adv. Res. Rev.*, **12**(3): 26-32.
84. Prosdocimi, M., Tarolli, P. and Cerdà, A. (2016). Mulching practices for reducing soil water erosion: A review. *Earth-Sci. Rev.*, **161**: 191-203.
85. Qaseem, M.F., Qureshi, R. and Shaheen, H., (2019). Effects of pre-anthesis drought, heat and their combination on the growth, yield and physiology of diverse wheat (*Triticumaestivum* L.) genotypes varying in sensitivity to heat and drought stress. *Sci. Rep.*, **9**(1): 6955.
86. Qu, B. and Luo, Y. (2020). Chitosan-based hydrogel beads: preparations, modifications and applications in food and agriculture sectors- A review. *Int. J. Biol. Macromol.*, **152**: 437-448.
87. Raja, M.A.S., Shahid, A.M., Saleem, M.F., Khan, I.H., Ahmad, S., Ali, M. and Iqbal, R. (2017). Effects and man agement strategies to mitigate drought stress in oilseed rape (*Brassicanapus* L.): A review. *Zemdirbyste Agriculture*, **104**(1): 85-94.
88. Rajanna, G.A., Manna, S., Singh, A., Babu, S., Singh, V.K, Dass, A., Chakraborty, D., Patanjali, N., Chopra, I., Banerjee, T.,Kumar, A., Khandewal, A. and Parmar, B.S. (2022). Biopolymeric superabsorbent hydrogels enhance crop and water

- productivity of soybean- wheat system in Indo-Gangetic plains of India. *Scientific Reports*, **12**(1): 1-18.
89. Ranjan, P., Patle, G.T., Prem, M., Solanke, K.R. (2017). Organic mulching- A water saving technique to increase the production of fruits and vegetables. *Curr. Agric. Res. J.*, **5**(3): 371-380.
90. Rao, D.S.N., Naidu, T. and Rani, Y.A. (2016). Change in Photosynthetic Rate, RWC, SCMR, Dry Matter Production and Yield of Mung Bean Due to Foliar Nutrition Under Receding Soil Moisture Condition. *Adv. Life Sci.*, **5**(19): 8729-8734.
91. Rasul, S. (2023). Effect of foliar application salicylic acid and drought stress on growth and yield of mung bean (*Vigna radiata*). *agric. Plant Sci.*, **34**(5): 102-113.
92. Ratnasekera, D.P. and Subhashi, A.P.T. (2015). Morphophysiological response of selected Sri Lankan mung bean (*Vigna radiata* L.) genotypes to drought Stress. *J. Agri. Res.*, **2**: 62-66.
93. Razmi, N., Ebadi, A., Daneshian, J. and Jahanbakhsh, S. (2017). Salicylic acid induced changes on antioxidant capacity, pigments and grain yield of soybean genotypes in water deficit condition. *J. Plant Interact.*, **12**: 457-464.
94. Rekhter, D., Lüdke, D., Ding, Y., Feussner, K., Zienkiewicz, K., Lipka, V., Wiermer, M., Zhang, Y. and Feussner, I. (2019). Isochorismate-derived biosynthesis of the plant stress hormone salicylic acid. *Sci.*, **365**: 498-502.
95. Saha, A., Sekharan, S. and Manna, U. (2020). Superabsorbent hydrogel (SAH) as a soil amendment for drought management: A review. *Soil Till. Res*, **204**: 104736.
96. Saha, A., Gupt, C.B. and Sekharan, S. (2021). Recycling natural fibre to superabsorbent hydrogel composite for conservation of irrigation water in semi-arid regions. *Waste and Biomass Valorization*, **12**(12): 6433-6448.
97. Salarizdah, M., Baghizadeh, A., Abasi, F., Mozaferi, H. and Salarizdah, S. (2012). Response of *Brassica napus* L. grains to the interactive effect of salinity and salicylic acid. *J. Stress Physiol. Biochem.*, **8**(2): 159-166.
98. Sankari, M., Hridya, H., Sneha, P., Doss, C. G.P., Christopher, J.G., Mathew, J. and Ramamoorthy, S. (2019). Implication of salt stress induces changes in pigment production, antioxidant enzyme activity, and q RT-PCR expression of genes involved in the biosynthetic pathway of *Bixaorellana* L. *Functional and Integrative Genomics*, **19**: 565-574.

99. Sarita, S.O., Singh, I., Shukla, U.N. and Yadav, S.K. (2020). Productivity and economics of mungbean (*Vigna radiata*L.) as influenced by fertility levels and stress mitigating chemicals. *J. Pharmacogn. Phytochem.*, **9**(1): 906-910.
100. Semida, W.M., Abd El-Mageed, T.A. and Howladar, S.M. (2014). A novel organo-mineral fertilizer can alleviate negative effects of salinity stress for eggplant production on reclaimed saline calcareous soil. *Acta Hortic.*, **1034**: 493-499.
101. Shankarappa, S.K., Munayandi, S.J., Chandra shekar, A.B., Singh, A.K., Naganhushanaradhya, P., Shivashanka, B., El-Ansary, D.O., Wani, S.H. and Elansary, H.O. (2020). Standarizing the hydrogel application rates and foliar application rates and foliar nutrition for enhancing yeild of lentil (*Lens culinaris*). *Preocesses*, **8**(4): 420.
102. Shao, Y., Xie, Y., Wang, C., Yue, J., Yao, Y., Li, X., Liu, W., Zhu, Y. and Guo, T. (2016). Effects of different soil conservation tillage approaches on soil nutrients, water use and wheat-maize yield in rainfed dry-land regions of North China. *Eur. J. Agron.*, **81**: 37-45.
103. Sharma, M., Gupta, S.K., Majumder, B., Maurya, V.K., Deeba, F., Alam, A., and Pandey, V. (2017). Salicylic acid mediated growth, physiological and proteomic responses in two wheat varieties under drought stress. *J. Proteomics*, **163**: 28-51.
104. Sharma, N.K., Singh, R.J., Mandal, D., Kumar, A., Alama, N.M. and Keesstra, S. (2017). Increasing farm er's income and reducing soil erosion using intercropping in rainfed maize-wheat rotation of Himalaya, India. *Agric. Ecosyst. Environ.*, **247**: 43-53.
105. Sharma, S.K., Mishra, P.K., Panse, R. and Jamliya, G. (2018). Effect of irrigation methods on yield attributes and water productivity of wheat in vertisol of betwa river basin commands of Vidisha district of M.P. India. *Int. J. Curr. Microbiol. App. Sci.*, **7**(8): 2670-2673.
106. Shemi, R., Wang, R., Gheith, E.S.M., Hussain, H.A., Hussain, S., Irfan, M., and Wang, L. (2021). Effects of salicylic acid, zinc and glycine betaine on morpho-physiological growth and yield of maize under drought stress. *Scientific Reports*, **11**(1): 3195.
107. Shrestha, U., Amgain, P., Karki, T.B., Dahaland, K.M. and Shrestha, J. (2016). Effect of sowing dates and maize cultivars in growth and yield of maize along with their agro-climatic indices in Nawalparasi. *Nepal J. AgriSearch*, **3**(1): 57-62.

108. Singh, L., Beg, M.K.A., Akhter, S., Qayoom, S., Lone, B.A., Singh, P. and Singh, P. (2014). Efficient techniques to increase water use efficiency under rainfed eco-systems. *J. AgriSearch*, **1**(4): 193-200.
109. Song, B., Liang, H., Sun, R., Peng, P., Jiang, Y., She, D. (2020). Hydrogel synthesis based on lignin/sodium alginate and application in agriculture. *International Journal of Biological Macromolecules*, **144**: 219-230.
110. Sultana, S., Rahman, M.M., Das, A.K., Haque, M.A., Rahman, M.A., Islam, S.M.N. and Mostofa, M.G. (2024). Role of salicylic acid in improving the yield of two mung bean genotypes under waterlogging stress through the modulation of antioxidant defense and osmoprotectant levels. *Plant Physiol. Biochem.*, **206**: 108230.
111. Suryapani, S., Malik, A., Sareer., O and Umar, S. (2014). Potassium and Rhizobium application to improve quantitative and qualitative traits of lentil (*Lens culinaris* Medik.). *Int. J. Agron. agric. Res.*, **5**: 7-16.
112. Swapna, S. and Shylaraj, K.S. (2017). Screening for osmotic stress responses in rice varieties under drought condition. *Rice Sci.*, **24**(5): 253-263.
113. Tanveer, S., Akhtar, N., Ilyas, N., Sayyed, R.Z., Fitriatin, B.N., Perveen, K. and Bukhari, N.A (2023). Interactive effects of *Pseudomonas putida* and salicylic acid for mitigating drought tolerance in canola (*Brassica napus* L.). *Heliyon*, **9**: 14193.
114. Teame, G., Tsegay, A., Abrha, B. (2017). Effect of organic mulching on soil moisture, yield, and yield contributing components of sesame (*Sesamum indicum* L.). *Int. J. Agron.*, **2017**(6): 1-6.
115. Tongo, A., Mahdavi, A. and Sayad, E. (2014). Effect of superabsorbent polymer aquasorb on chlorophyll, antioxidant enzymes and some growth characteristics of *Acacia victoriae* seedlings under drought stress. *Ecopersia*, **2**: 571-582.
116. Tu, Y., Wang, R., Zhang, Y. and Wang, J. (2018) .Progress and expectation of atmospheric water harvesting. *Joule*,**2**: 1452-1475
117. Tyagi, V., Nagargade, M. and Singh, R.K. (2020). Agronomic interventions for drought management in crops. In: *New Frontiers in Stress Management for Durable Agriculture* (Eds. Rakshit, A., Singh, H., Singh, A., Singh, U. and Fraceto, L.). Springer, Singapore, pp. 461-476.
118. Ullah, C., Chen, Y.H., Ortega, M.A. and Tsai, C.J. (2023). The diversity of salicylic acid biosynthesis and defence signaling in plants: Knowledge gaps and future opportunities. *Curr. Opin. Plant Biol.*,**72**: 102349.

119. Vaisnad, S. and Talebi, R. (2015). Salicylic acid-enhanced morphological and physiological responses in chickpea (*Cicer arietinum*) under water deficit stress. *Environ. Experl. Biol.* **13**: 109-115.
120. Vance, W.H., Bell, R.W., Johansen, C., Haque, M.E., Musa, A.M., Shahidullah, A.K.M. and Mia, M.N.N. (2014). Optimum time of sowing for rainfed winter chickpea with one-pass mechanised row-sowing: an example for small-holder farms in north-west Bangladesh. *Crop and Pasture Sci.*, **65**(7): 602-613.
121. Wang, Y.Y., Wang, Y., Li, G.Z and Hao, L. (2019). Salicylic acid-altering Arabidopsis plant response to cadmium exposure: Underlying mechanisms affecting antioxidation and photosynthesis-related processes. *Ecotoxicology Environ. Safety*, **169**: 645-653.
122. Watcharamul, S., Lerddamrongchai, S., Siripongpreda, T., Rodtassana, C., Nui sin, R. and Kiatkamjornwong, S. (2022). Effects of carboxymethyl cellulose/nano-calcium carbonate hydrogel amendment of loamy sand soil for maize growth. *ACS agric. Sci. Technol.*, **2**(5): 1071-1080.
123. Xu, L., Zhao, H., Wang, J., Wang, X., Jia, X., Wang, L., Xu, Z., Li, R., Jiang, K. and Chen, Z. (2023). AIM1-dependent high basal salicylic acid accumulation modulates stomatal aperture in rice. *New Phytol.*, **238**: 1420-1430.
124. Yadav, M., Chandola, V.K. and Galkate, R. (2014). Life saving irrigation under drought condition for kharif crops in vindhyan zone (Mirzapur district), UP trends. *Trends Biosci.*, **7**(22): 3700-3704.
125. Yousefzadeh-Najafabadi, M. and Ehsanzadeh, P. (2017). Photosynthetic and antioxidative upregulation in drought-stressed sesame (*Sesamum indicum* L.) subjected to foliar-applied salicylic acid. *Photosynthetica*, **55**: 611-622.
126. Zain, N.A.M and Ismail, M.R. (2016). Effects of potassium rates and types on growth, leaf gas exchange and biochemical changes in rice (*Oryza sativa*) planted under cyclic water stress. *agric. Water Mgt.*, **164**: 83-90.
127. Zhang, C., Shi, S., Liu, Z., Yang, F. and Yin, G. (2019a). Drought tolerance in alfalfa (*Medicago sativa* L.) varieties is associated with enhanced antioxidative protection and declined lipid peroxidation. *J. Plant Physiol.*, **232**: 226-240.
128. Zhu, J., Suhaimi, F., Lim, J.Y., Gao, Z., Swarup, S., Loh, C.S., Li, J., Ong, C. N. and Tan, W.K. (2022a). A field study on using soybean waste-derived superabsorbent hydrogel to enhance growth of vegetables. *Sci. Total Environ.*, **851**: 158141.

UNDER PEER REVIEW