

# Original Research Article

## Optimization and Characterization of Freeze-Dried Pumpkin Powders: Powder Quality and Nutrient Stability

### ABSTRACT

Being a good source of nutrients, viz., ascorbic acid,  $\beta$ -carotene, phenols, and antioxidants, pumpkin pulp (*Cucurbita moschata*) can be utilised as a functional ingredient in many food products by converting them into a more stable product like pumpkin powder. Hence the present study is focused on the production of freeze-dried pumpkin powder by two different methods (drying in petri plates with 5mm sample thickness and in 250 ml flask). The moisture content and water activity were reduced by 92.50% and 74.73%, respectively, compared to the fresh pumpkin pulp, indicating safe storage limits. Freeze-dried pumpkin powders exhibited a decrease of 25% ascorbic acid, 21.58%  $\beta$ -carotene, 39.39% reduction in total polyphenols and 89.58% in antioxidant content. The powder properties of the developed freeze-dried powders exhibited water activity of  $0.4a_w$  to  $0.35a_w$ , bulk density of  $0.15$  to  $0.12 \text{ g cm}^{-3}$ , and flowability of  $18.39^\circ$  to  $22.74^\circ$ .

*Keywords: Pumpkin, freeze-drying, nutritional properties, powder quality*

### 1. INTRODUCTION

*Cucurbita moschata*, commonly known as winter squash or pumpkin, is a highly nutritious crop rich in carotenoids, vitamins, dietary fibre, minerals, phenolic compounds, and bioactive substances [1]. The fruit pulp is a rich source of beta-carotene, vitamin A, tocopherol, and dietary fibre. It aids in regulating insulin, lowering blood sugar, improving glucose tolerance, and offers anti-obesity, anti-diabetic, antibacterial, and anticancer **propertiseffects incorporating** [1] [2]. This versatile crop not only offers a wide array of essential nutrients but also shows promising medicinal value, highlighting its significance in enhancing overall health and nutrition.

Conventional drying methods for pumpkin pulp are hindered by several challenges, like depletion of nutrients and the lengthy drying process. Even though drying process enhances specific nutrients like  $\beta$ -carotene and protein, it also results in decreased levels of zinc, iron, calcium, and energy due to oxidation [3]. Moreover, traditional techniques may not adequately conserve the nutritional quality and visual appeal of pumpkins, giving rise to issues such as nutrient loss, machine sticking, and saccharification during the manufacturing process [4]. These limitations underscore the significance of exploring novel drying methodologies, such as freeze drying, to effectively uphold the nutritional value and excellence of pumpkin powder.

Freeze-drying plays a crucial role in the preservation of pumpkin pulp, as it aids in the retention of its nutritional composition and prolongs its shelf life. Studies have demonstrated that the freeze-drying process of pumpkin pulp leads to superior preservation of essential nutrients such as total carotenoids, proteins, and total phenolic compounds when compared to alternative drying methods [5][6]. Moreover, freeze-drying plays a pivotal role in upholding the intrinsic characteristics of the pumpkin powder, including moisture content, particle size,

Comment [Sk1]: properties

and density, thereby ensuring the secure storage of the final product [7]. Additionally, the efficacy of freeze-drying in maintaining bioactive compounds and antioxidant properties in fruits like carnauba underscores its potential to safeguard the nutritional attributes of pumpkin pulp over an extended period[8]. In conclusion, freeze-drying is a valuable method for the preservation of pumpkin pulp, safeguarding its nutritional integrity, and augmenting its shelf stability.

## 2. MATERIAL AND METHODS

The present study was carried out in the Department of Postharvest Management, College of Agriculture, Vellayani. Fruits of pumpkin (*Cucurbita moschata*) variety 'Ambili', which were discarded as waste at the vegetable seed production unit of the College of Agriculture, Vellayani, were employed for the research. The pumpkin pieces were washed, peeled, cut into small pieces with a vegetable cutter, ground using a mixer grinder, and strained to extract the pulp. The pulp was then pre-frozen at  $-20^{\circ}\text{C}$  for 4-5 hrs. and freeze-dried under vacuum at  $-48 \pm 2^{\circ}\text{C}$  condenser temperature. The frozen pulp thus obtained was freeze-dried in both petri plates and in a 250ml flask. The temperature of the heating plate was maintained constant throughout the process of drying. The dried pulp was converted into powder using a blender. The experiment was done in CRD with two treatments replicated ten times using the statistical web application 'GRAPES' [9]

Biochemical and nutritional quality parameters of the freeze-dried powder were assessed. Moisture content[10], ascorbic acid,  $\beta$  carotene, total polyphenol content [11], crude fibre [12] and total antioxidant activity [13] were analysed.

Physical quality parameters such as water activity were discerned using the NovasinaLabSwift-aw water activity meter (M/s. Novasina..AG, Neuheimstrasse 12, CH-8853 Lachen, Switzerland); bulk density by tapping method [14] and the angle of repose was measured to characterize the flowability of freeze-dried powders through fixed funnel method.

## 3. RESULTS AND DISCUSSION

### 3.1 Biochemical and nutritional quality parameters

The biochemical and nutritional quality parameters of the fresh pumpkin pulp and developed freeze-dried powders are depicted in Figure 1.

Fresh pumpkin pulp exhibited a moisture content of 92.5% which was reduced to 1.98% for FDPP<sub>1</sub> and 2.81% for FDFP<sub>1</sub>. The moisture content of FDFP was greater than FDPP. Similar results were obtained for the moisture content of freeze-dried mango (4%), guava (5%) and strawberry slices (9%) [15]. Freeze-dried powders exhibit a very low moisture content which aids in their storage for a longer period of time [16]. The ascorbic acid content of the developed freeze-dried powders was lower compared to that of the fresh pumpkin pulp ( $8.85 \text{ mg } 100\text{g}^{-1}$ ). An ascorbic acid content of  $6.6 \text{ mg } 100\text{g}^{-1}$  was exhibited by FDPP and  $6.49 \text{ mg } 100\text{g}^{-1}$  by FDFP. A loss of about 25% of ascorbic acid was observed. Similarly, a loss in ascorbic acid content was observed in freeze-dried tomatoes (8.2% loss), guava (37%), mango (26.92%) and pineapple (24.20%) [15] [17].

$\beta$ -carotene content of the fresh pumpkin pulp was found to be  $7.46 \text{ mg } 100\text{g}^{-1}$ . This was reduced to  $5.85 \text{ mg } 100 \text{ g}^{-1}$  for FDPP and  $4.63 \text{ mg } 100 \text{ g}^{-1}$  for FDFP. A reduction of about 21.58 % was observed for the  $\beta$ -carotene content. Freeze-drying retains more  $\beta$ -carotene content compared to other drying methods [18]. A reduction in  $\beta$ -carotene content of 15.75 % in carambola, 8.6% in papaya, 43.1% in watermelon, and 26.19% in freeze-drying were observed [19].


The total polyphenol content of the fresh pumpkin pulp was  $526.81 \text{ mg GAE g}^{-1}$ . The developed FDPP exhibited  $316.62 \text{ mg GAE g}^{-1}$ , and FDFP exhibited  $303.80 \text{ mg GAE g}^{-1}$ , accounting for a loss of about 39.39%. Retention of the majority of the phenolic compounds

Comment [Sk2]: give full abrevation when the code comes for the first time

in freeze-dried strawberry and white zapote were also reported [20] [21]. A crude fibre content of 3.74% was recorded for the fresh pumpkin pulp. FDPP showed a crude fibre content of 1.52% and FDPF showed a crude fibre content of 1.15%. Similar retention in crude fibre content in free-dried watermelon rind was observed [22]. About 89.58% of antioxidant activity was observed for fresh pumpkin pulp, 67.90% for FDPP and 57.11% for FDPF. Antioxidant activity was reduced to only 24.20% after the freeze-drying process. A similar result was observed for Chilean guava, where the antioxidant content was reduced to 33.30% of the fresh fruit [23].

**Figure I & II: Nutritional parameters in fresh pumpkin pulp and freeze-dried pumpkin powder**

Comment [Sk3]: write this sentence separately below the tables as figure 1 and 2



T1- Fresh pumpkin pulp  
T2 -Freeze-dried pumpkin powder in petri plate  
T3- Freeze-dried pumpkin powder in flask

### 3.2 Physical quality parameters

The physical quality parameters of the developed freeze-dried powders are depicted in Table 1.

The water activity of the fresh pumpkin pulp was recorded as 0.95  $a_w$ . FDPP exhibited a water activity of 0.24 $a_w$  and FDPF, 0.35 $a_w$ . A reduction of about 74.73% in water activity of the freeze-dried powders were observed. Water activity values between 0.20 and 0.40 are found to be stable during storage as it reduces the chances of browning and oxidation. Water activity of 0.2 was observed for freeze-dried acerola powder and 0.26 for freeze-dried sour cherry powder [24] [25].

Bulk density is an important parameter in powder production for packaging, storage, and transport. Freeze-dried pumpkin powders exhibited a bulk density of 0.15g cm<sup>-3</sup> for FDPP and 0.12 g cm<sup>-3</sup> for FDPF. A similar bulk density of 0.13 g cm<sup>-3</sup> was observed for freeze-dried pumpkin powder in another study [6]. Flowability (angle of repose) of the developed freeze-dried pumpkin powders done using fixed funnel method recorded a value of 18.39° for FDPP and 22.74° for FDPF. A similar result was reported in freeze-dried jamun powder which recorded an angle of repose of 29° [26].

**Table 1: Physical properties of freeze-dried powder**

#### 4. CONCLUSION

In the present study, freeze-drying of pumpkin pulp was done in petri plates and in 250 ml flasks and their physicochemical characteristics were analysed. Freeze-drying of the pumpkin pulp in petri plates exhibited higher nutrient retention and physical quality parameters compared to the freeze-dried pulp in 250 ml bulb. Physical quality parameter values represented excellent powder quality characteristics for the developed freeze-dried powders.

#### REFERENCES

1. Men X, Choi SI, Han X, Kwon HY, Jang GW, Choi YE, et. al. Physicochemical, nutritional and functional properties of Cucurbita moschata. Food Sci Biotechnol. 2021;30:171-83.
2. Wahyono A, Budiati T, Shahbaz HM. Feasibility of Using Yellow Pumpkin (Cucurbita moschata) in Developing Bakery Products. 2023
3. Kiharason JW, Isutsa DK, Ngoda PN. Effect of drying method on nutrient integrity of selected components of pumpkin (Cucurbita moschata Duch.) fruit flour. ARPN J Agric Biol Sci. 2017;12(3):110-6.
4. Potosi-Calvache, D.C., Vanegas-Mahecha, P. and Martínez-Correa, H.A., 2017.

Treatments	Moisture content (%)	Water activity ( $a_w$ )	Bulk density(g/ml)	Flowability(Angle of repose)(degree)
T1 (Petri plate)	1.98	0.24	0.15	18.39
T2 (flask)	2.81	0.35	0.12	22.74
SE(+ m)	0.041	0.002		1.171
CD (0.05)	0.123	0.006	NS	3.479
Treatments	Moisture content (%)	Water activity ( $a_w$ )	Bulk density(g/ml)	Flowability(Angle of repose)(degree)
T1 (Petri plate)	1.98	0.24	0.15	18.39
T2 (flask)	2.81	0.35	0.12	22.74
SE(+ m)	0.041	0.002		1.171
CD (0.05)	0.123	0.006	NS	3.479

Convective drying of squash (*Cucurbita moschata*): Influence of temperature and air velocity on effective moisture diffusivity, carotenoid content and total phenols. *Dyna*, 84(202), pp.112-119.

5. Umuhozariho MG, Hagenimana T, Nsabimana P, Sirimu C, Uwobasa N, Uwineza AP. Effect of oven and freeze drying on nutritional composition of pumpkin (*Cucurbita maxima*) processed flour. *Rwanda J Agric Sci*. 2020;2(1):33-9.
6. Dirim, S. Nur, and G. Çalışkan. "Determination of the effect of freeze drying process on the production of pumpkin (*Cucurbita moschata*) puree powder and the powder properties." (2012): 203-210.
7. Ibrahim MN, Taip FS, How MS, Lim J, Ab Aziz N. Effects of drying methods on the physicochemical properties of powder made from different parts of pumpkin. *Food Res (Malaysia)*. 2021.
8. Borges JM, Lucarini M, Durazzo A, Rufino Arcanjo DD, Rodrigues Lima SK, da Silva RA. Effect of freezing and freeze-drying on bioactive compounds and antioxidant activity of carnauba pulp (*Copernicia prunifera* (Mill.) HE Moore). *MOJ Food Process Technol*. 2023;11(2):78-82.
9. Gopinath PP, Parsad R, Joseph B, Adarsh VS. GrapesAgri1: collection of shiny apps for data analysis in agriculture. *Journal of Open Source Software*. 2021 Jul 18;6(63):3437.
10. Lane JH, Eynon L. Methods for determination of reducing and non-reducing sugars. *J Sci*. 1923;42:32-7.
11. Ranganna S. Handbook of analysis and quality control for fruit and vegetable products. Tata McGraw-Hill Education; 1986.
12. Sadasivam S. Biochemical methods. New age international; 1996.
13. Sharma OP, Bhat TK. DPPH antioxidant assay revisited. *Food Chem*. 2009 Apr 15;113(4):1202-5.
14. Bhandari BR, Dumoulin ED, Richard HM, Noleau I, Lebert AM. Flavor encapsulation by spray drying: application to citral and linalyl acetate. *J Food Sci*. 1992;57(1):217-21.
15. Marques LG, Prado MM, Freire JT. Vitamin C content of freeze-dried tropical fruits. In *Int Cong Eng Food*. 2011 May; (pp. 22-26).
16. Shishegarha F, Makhlof J, Ratti C. Freeze-drying characteristics of strawberries. *Drying Technol*. 2002;20(1):131-45.
17. Chang CH, Lin HY, Chang CY, Liu YC. Comparisons on the antioxidant properties of fresh, freeze-dried and hot-air-dried tomatoes. *J Food Eng*. 2006;77(3):478-85.
18. Serna-Cock L, Vargas-Muñoz DP, Aponte AA. Structural, physical, functional and nutraceutical changes of freeze-dried fruit. *Afr J Biotechnol*. 2015;14(6):442-50.
19. Shofian NM, Hamid AA, Osman A, Saari N, Anwar F, Dek MS, Hairuddin MR. Effect of freeze-drying on the antioxidant compounds and antioxidant activity of selected tropical fruits. *Int J Mol Sci*. 2011;12(7):4678-92.
20. Yurdugül S. An evaluation of the retention of quality characteristics in fresh and freeze-dried alpine strawberries. *Inter J Food Sci Technol*. 2008;43(5):865-70.
21. Castañeda J, Miñano HA, Jara RS, Rodriguez G. Comparative study of the loss of vitamin C in chalařina (*Casimiroa edulis*) by four methods of dehydration. *Scientia Agropecuaria*. 2010;1(1).
22. Ho LH, Suhaimi MA, Ismail I, Mustafa KA. Effect of different drying conditions on proximate compositions of red-and yellow-fleshed watermelon rind powders. *J Agrobiotechnol*. 2016;7:1-2.
23. Reyes A, Bubnovich V, Bustos R, Vásquez M, Vega R, Scheuermann E. Comparative study of different process conditions of freeze drying of 'Murtilla'berry. *Drying Technol*. 2010;28(12):1416-25.

24. Can Karaca A, Baskaya H, Guzel O, Ak MM. Characterization of some physicochemical properties of spray-dried and freeze-dried sour cherry powders. *J Food Process Preservation*. 2020;44(12):e14975.
25. Marques LG, Ferreira MC, Freire JT. Freeze-drying of acerola (*Malpighia glabra* L.). *Chem Eng Process: Process Intensification*. 2007;46(5):451-7.
26. Reginold Jebitta S, Jeyanth Allwin SI. Effect of drying methods on the physical properties of Jamun pulp powder. *History*. 2016;23(77):50-9.

#### **DEFINITIONS, ACRONYMS, ABBREVIATIONS**

**FDPP:** Freeze-Dried Powder in Petriplate

**FDPF:** Freeze-Dried Powder in Flask

UNDER PEER REVIEW