

Review Article

A Review on status of mushroom pests, pesticides usage and multi-residue analysis methods

Abstract

Mushrooms are valued for their dietary benefits and medicinal properties, being rich in proteins, minerals, vitamins, fiber, and essential amino acids. They offer significant health benefits, including the prevention of hypercholesterolemia and cardiovascular diseases, and possess antitumor, antiviral, antihypertensive, anti-inflammatory, and immunomodulating properties. Global mushroom production has seen substantial growth, particularly in Asia. However, mushroom cultivation is challenged by various insect pests, leading to significant crop losses. Pesticides are commonly used to prevent these pests, but their use can result in pesticide residues in mushrooms, posing food safety risks. Multiresidue analytical methods are essential for detecting low levels of pesticide residues. As QuEChERS method is sensitive, selective and accurate for determining multiple pesticide residues, so it is the most widely used method. These methods can identify various pesticide ingredients in a single run, addressing the unique biological characteristics of mushrooms. The MRL levels for pesticide residues in mushrooms vary, with stringent guidelines set by regulatory bodies. This paper reviews the pests affecting mushroom cultivation, the pesticides used, and the analytical methods developed for pesticide residue analysis in mushrooms, highlighting the importance of accurate detection for food safety and market access.

Key words: Mushroom; Pests; Pesticides; QuEChERS; Multiresidue

1. Introduction

Mushrooms are well known for their dietary benefits and as a significant source of bioactive compounds with medicinal properties (Cheung et al. 2010). Mushrooms are considered a valuable supplement to cereals due to their high content of proteins, minerals, fiber, and essential amino acids. They are rich in protein (2-40%), fat (2-8%), carbohydrates (1-55%), and fiber (3-32%). Additionally, mushrooms provide vitamins (1.4-2.2 mg of thiamine, 6.7-9 mg of riboflavin, 21.1-33.3 mg of pantothenic acid, and 1.2-1.4 mg of folic acid), minerals (such as aluminum, iron, manganese, magnesium, zinc, and selenium), and ash (8-10%) (Kakon et al. 2012). Their low carbohydrate content makes them beneficial for preventing hypercholesterolemia and cardiovascular diseases (Waktola and Temesgen 2018). Moreover, certain mushroom species possess antitumor, antiviral, antihypertensive, anti-inflammatory, and immunomodulating properties (Daba et al. 2008). The popularity of mushrooms is growing, with increased consumption of both fresh and processed varieties in the market. Over the past 20 years, global mushroom production has increased at a compound annual growth rate (CAGR) of 8.26%. Among continents, Asia has experienced the highest CAGR at 8.97%, followed by Africa with a CAGR of 6.31% (Shweta and Sharma 2023). Since 2000, global mushroom production has increased more than fivefold, reaching 44 million tonnes as of 2023 (FAOSTAT 2023). Asia dominates production, contributing 95% of the total, followed by Europe with 3% and the Americas with 1%. Leading countries in fresh mushroom production include China (93%), Japan (0.01%), and Poland (0.01%). India ranks sixth, with around 0.24 million tonnes of production according to data of FAO, 2023. Though domestic updates suggest it has reached 0.31 million tonnes. Despite significant advancements in China, the compound annual growth rates (CAGR) for fresh mushroom production in China (9.23%) and India (8.58%) over the last 20 years are comparable,

reflecting encouraging growth for the Indian mushroom industry relative to the global leader. In regions like Europe and the USA, white button mushrooms are predominantly cultivated on a commercial scale. However, in Asian countries such as China, a variety of mushroom species are also commercially produced. Mushrooms are exported to UAE, Russia, Netherland, Germany, Switzerland, U.K, Denmark, Sweden and other countries from India (Pandey et al. 2018).

Mushrooms are susceptible to a range of insect pests that can cause substantial crop losses (Jiang et al. 2012). In order to prevent this problem, pesticides are commonly used during mushroom cultivation. The indoor cultivation environment, characterized by limited light and high moisture, creates ideal conditions for pests and diseases, which can significantly impact yield and quality (Lewandowski et al. 2004). Key pests include sciarids (*Lycoriella* sp.), phorids (*Megaselia* sp.), springtails (*Lepidocyrtus* sp.), and mites (*Tyrophagus* sp.) (Rijaj et al. 2021). Flies are particularly problematic, contributing to yield losses of up to 60-70% globally (Erler et al. 2015). Their larvae damage the developing fruiting bodies and mycelial front, often rendering mushrooms unmarketable (Grewal 2000). The sciarid fly (*Lycoriella* sp.) is notably detrimental, leading to yield losses of up to 40% (Lewandowski et al. 2004). Additionally, phorid flies act as vectors for the pathogen, *Verticillium fungicola* var. *fungicola*. Improper pasteurization of compost and casing becomes a major source of fungal and bacterial infections. The attraction of mushrooms to decaying fungi also attracts flies, complicating pest management (Maurya et al. 2019). To prevent yield loss, pesticides are widely utilized mainly throughout the growing process as well as after cultivation (FAO 2018). Uncontrolled implementation of these pesticides leads to pesticide residues on agricultural products (Chen et al. 2011). It was found that residues of some of the insecticides such as chlorpyrifos, cypermethrin, methomyl and several pyrethroids occur in the mushrooms, soil and water (Le et al. 2021). Pesticide residue analysis was done in 145

samples of mushroom. Carbendazim was more frequently detected above the MRL (Jess and Kilpatricky 2000). Permethrin (50 g/ kg; MRL - 100 g/kg) and cypermethrin (200 g/kg; MRL - 500 g/kg) were found in shiitake mushroom and malathion (30 g/kg; MRL - 500 g/kg) was found in *Pholiota nameko* (nameko mushroom) samples (Cao et al. 2016). Edible mushrooms are highly preferred and hence presence of detectable residues may pose threat to food safety and entry to international market. Therefore, it is important to monitor residues in mushroom to reduce the use of pesticides and improve the quality.

Highly selective, sensitive and accurate analytical methods are needed to measure low amount of pesticide residues in mushroom. Multiresidue methods are efficiently used as there are large number of pesticides used and should be detected in one single run (Anastasiades et al. 2003). The traditional pesticide residue analysis is typically used to identify a single pesticide ingredient, whereas the multiresidue analysis method can be used to identify not only various pesticide ingredients, but also various ingredients of different pesticide categories. As a result, one of the topics of analytical chemistry that is being actively researched is the analytical technique for pesticide multi-residue (Anastasiades et al. 2007). Vegetables in general have biological characteristics that are extremely different from those of edible fungus. The reverse resistance of fungi is weak due to lack of surface protective structures like wax layers, etc.

The MRL level for pesticide residues in mushroom ranges between 0.001 mg/kg to 3 mg/kg depending on the pesticide (Anastasiades et al. 2007). It may even established higher limits like in Bromide ion compounds has MRL of 30 mg/kg. The EURLs are responsible for promoting new analytical methods by proper guiding and organising proficiency tests, supporting the creation which leads to certification of novel analytical methods. Guidance is provided to laboratories for the validation of procedures to analysing pesticide residues in food and feed by

the Directorate for Health and Food Safety of the European Commission (SANTE 2021). This instruction gives laboratories the freedom to choose their own procedures, which is advantageous for the ongoing improvement of analytical techniques. Laboratories that assess pesticide residues frequently operate under a quality management system like ISO/EC 17025. This paper gives overview of pests, pesticides and different analytical methods which were adopted in analysis of pesticide residues in mushroom upto today.

2. Pests of mushroom

2.1. Insect pests of mushroom

2.1.1. International status

Mushrooms are more prone to insect pests mainly due to their flavor and cultivation practices, which cause tremendous yield loss. The presence of insect pests reduces quality, quantity, marketability and palatability of mushroom. Keil (2002) reported that phorid fly (*Megaselia* sp.) was the major pest in USA and made direct damage by feeding on mushroom and also indirectly by transmission of fungal pathogens. Sciarid fly (*Lycoriella* sp.) considered as one of the dominant insect pest of mushroom crop throughout the world which cause significant yield loss up to 40 per cent (Lewandowski et al. 2004). Compared to other mushroom species, the sciarid fly *Lycoriella mali* (Fitch) prefers the button mushroom, *Agaricus bisporus* (Lange) Imbach. Infestation was more in button mushroom due to high developmental period, size and survival rate (Connor and Keil 2005). Sciarid fly, *Lycoriella ingenua* (Dufour) was reported as dominant pests in North America and both *Lycoriella ingenua* and *Lycoriella castanescens* (Lengersdorf) in U.K. In other regions of the world, *Bradysia* sp. (Sciarid fly) reported as major pest of button mushroom causing serious damage, though it also reproduce on oyster and shiitake mushrooms (Fletcher and Gaze

2008). Pyemotid or pygmy mites, often known as red pepper mites, usually feed on moulds like *Trichoderma*, *Monilia*, and *Humicola* sp. They were predominantly present during commercial button mushroom production, which reduces marketable quality (Castilho et al. 2009). The key pests of mushroom reported from Korea include Sciarid flies namely *Camptomyia corticalis* (Loew), *C. heterobia* (Mamaev), *L. ingenua* (Dufour) (Shin et al. 2012). In Pakistan, the most destructive pest causing serious threat to mushroom cultivation is phorid fly, *Megaselia halterata* (Wood). The excreta of the larvae deteriorate the quality of mushroom. They primarily act as vectors of pathogens especially *Verticillium* sp. (Babar et al. 2012). Outbreak of pomace fly (*Drosophila* sp.) was severely noticed during 4th to 5th month of mushroom cultivation (Thongnaitam et al. 2012). Sciarid flies, *Lycoriella auripila* (Winnertz), *Lycoriella ingenua* (Dufour), *Lycoriella solani* (Winnertz), phorid flies, *Megaselia halterata* (Wood) and black flies - *Scaptopse notata* (Linnaeus) were the pests of mushroom reported from Turkey. *L. solani* was the common and cause more infestation from March to May and September to late November compared to other sciarid species (Erler and Polat 2014). Button mushroom and oyster mushroom were attacked by the cecid larvae, *Mycophila speyeri* (Barnes), *Heteropeza pygmaea* (Winnertz) and *Mycophila barnesi* (Edwards) in USA. The larvae feed on the exterior or at the point where the stipe and gills meet. They may cause a direct yield loss or reduction in the amount of fresh or processed marketable produce (Rinker 2017). *Bradysia impatiens* (Johannsen), *B. ocellaris* (Cosmstock), *Lycoriella agraria* (Felt), and *L. ingenua* were the sciarid flies that were described from Australia. Larvae of sciarid flies mainly feeds on the growing mycelium and through tunneling it destroys sporophore primordia (Broadley et al. 2018). Throughout the world, mushrooms were primarily attacked by three dipteran pests such as sciarid flies, *Lycoriella mali*, *L. auripila*, *L. solani*, phorid fly, *Megaselia halterata* (Wood) and cecid flies, *Mycophila speyeri* (Barnes), *Mycophila barnesi*

(Edwards), *Hetropeza pygmae* (Winnertz). Minor pests of mushroom were pomace fly (*Drosophila* sp.), springtails (*Lepidocyrtus* sp.) and mites, *Tarsonemus mycelophagus* (Hussey), *Pymephorus* sp. and *Tyrophagus* sp. (Rijal et al., 2021). Phorid fly, *M. halterata* has recently turned from minor to major pest and caused mushroom yield reduction from 10-40 per cent in USA, U.K and Turkey. *Megaselia flavinervis* (Malloch), *Megaselia agarici* (Lintner), *Megaselia iriguoiانا* (Felt) and *Megaselia bovista* (Gimmerthal), *Megaselia sandhui* (Disney) and *Megaselia tamilnaduensis* (Disney) were other phorid fly species which were reported in mushroom growing areas worldwide (Navarrao et al., 2021).

2.1.2. National status

Sciarid fly, *L. castanescens* and phorid fly (*Megaselia* sp.) were the threatening insect pests of mushroom in Kashmir valley (Munshi et al. 2010). First report of sciarid fly (*Lycoriella* sp.) on oyster mushroom was recorded during the month of April-May, 2012 in Assam. A single larva can cause damage around 20-35 per cent within 3-4 days of infestation. In severe infestations, 70-80 per cent of the fruit bodies in mushroom were damaged (Das et al., 2014). In India, mushrooms were dominantly attacked by pests such as: Sciarid fly, *Bradysia paupera* (Tuomikoski), *Bradysia. tritici* (Coquillett), *Lycoriella auripilla* (Winnertz), phorid fly (*Megaselia* sp.) and cecid fly, *Heteropezina cathistes* (Pritchard) which cause drastic reduction in mushroom yield. Springtails, *Lepidocyrtus cyaneus* (Tullberg), *Xenylla* sp. *Seria iricolor* (Say), mites (*Tyrophagus* sp., *Pygmephorous* sp.) and beetles, *Staphylinus* sp., *Alphitobius laevigatus* (Fabricius) and *Scaphisoma nigrofasciatum* (Pic) were found to cause 5-10 per cent damage. Minor pests such as spring tails (*Xenylla* sp., *Seria iricolor*) mainly damages mycelium and sporophore in the compost of both oyster and button mushroom. Besides it feeds on fruiting bodies and gills of mushroom. Nematode species such as *Aphelenchoides* sp., *Seinura* sp., *Tylenchids* sp.

and saprophagous nematodes genera like *Rhabditis*, *Caenorhabditis*, *Diplogaster*, *Acrobiloids* and *Panagrolaimus* were also infest mushroom (Singh and Sharma 2016). Duhan et al. (2018) reported that mites (*Tyrophagus* sp.) infest mushrooms starting from spawning to harvest and dipteran flies (sciarids and phorids) act as vectors for mites infestation in cultivated mushroom. Button mushroom is majorly attacked by insect pest belonging to order diptera and coleoptera in the Himalayan region. Spring tails and mites were also frequently encountered. Dipteran flies mainly are sciarid fly and phorid fly. *Lycoriella* sp. (Sciarid fly) has been identified in several countries that grow mushrooms, it has recently been documented in Himachal Pradesh. Additionally, *Megaselia* sp. (Diptera: Phoridae) was only found in two places. *Cyllodes indicus*, *Scaphisoma nigrofasciatum*, *Staphylinus* sp. and *Spondotriplax pallidipes* were four genera of beetles that were commonly found (Sharma et al. 2019). Incidence of sciarid fly (*Lycoriella* sp.) and phorid fly (*Megaselia* sp.) was frequent in button mushroom cultivation when compared to oyster and milky mushroom (Meena et al. 2021). Sciarid flies were notably abundant, with infestation levels of *A. bisporus* ranging from 9.00 to 38.63 percent across different districts. In May 2023, Solan exhibited the lowest infestation at 8.25%, while Mandi recorded the highest at 38.63% in March 2022. The infestation primarily targeted compost and casing materials, leading to substantial yield losses and inferior mushroom quality. A prominent sign of damage was the appearance of brown patches on the sporophores, visible on the fruiting bodies (Ruchika et al. 2021).

2.2. Diseases of mushroom

Apart from pest species, diseases were also a threat in mushroom cultivation. The prevalent bacterial disease affecting white button mushrooms is bacterial blotch, *Pseudomonas tolaasii* (Tolaas). Interstitial necrosis was the bacterial disease caused by bacteria *Ewingella americana*

(Grimont). Fungal diseases were dry bubble, *Verticillium fungicola* (Preuss), wet bubble - *Mycogone pernicioso* (Magn), cob web *Cladobotryum dendroides* (Bull), false truffle *Diehliomyces microsperes* (Deihl and Lambert), white plaster mould *Scopulariopsis fimicola* (Garcha) and brown plaster mould *Papulaspora byssina* (Hots) were the fungal diseases which cause damage to button mushrooms consistently (Grewal 2007). White button mushrooms were frequently affected by the fungi *Verticillium* sp., *Cladobotryum* sp., *Trichoderma harzianum* (Rifai), *Trichoderma viride* (Persoon). Green mould was the most affecting disease over past 20 years, which was due to pathogen *T. aggressivum* (Samuel). Wet bubble, dry bubble and other significant fungal diseases were detrimental to the cultivation of button mushroom in Turkey (Amin et al. 2021). The most prevalent viral disease, dieback was brought by different virus types. These illnesses pose a serious danger to the multimillion-dollar mushroom industries because they could result in losses of up to 100 per cent (Amin et al. 2021).

3. Pesticide usage pattern in mushroom

Due to attack of different pests in mushroom, it has become necessary to use pesticides to maintain yield. Since cultivation practices is having different temperature and humidity that introduces a wide range of pests. In order to solve these issues, growers are turning to use more pesticides recently.

Diflubenzuron showed 90-96 per cent effective control against sciarid larvae but could not avoid reduction of mushroom yield up to 7-8 per cent (Staunton et al., 1999). The quantity of immature sciarids during the first cropping period was reduced by 99 per cent by adding diazonin (25 mg a.i/kg) to mushroom compost, followed by an additional application of diflubenzuron (1 g a.i/m²) to the casing layer. Treatment with diazonin and diflubenzuron together in the first flush, reduced

the appearance of sciarids (Jess and Klipatrick 2000). Stoddart et al. (2004) recommended to use bendiocarb and diflubenzuron either drenched onto casing or incorporated in compost of mushroom which showed effective control around 50 to 70 per cent for flies, spiders, bugs, mites and larvae of insect pests. Malathion 0.05 per cent and dichlorvos 0.5 per cent were recommended as effective control for mushroom pests and applied in composts (DMR 2008). Aldrin, DDT, benzene hexachloride and thionazin was incorporated into the compost at both 100 and 500 µg/g and showed effective control. Bendiocarb was shown to be more effective in the range of 10 to 30 µg/g for sciarids when applied in casing. Diazonin when incorporated into the compost at spawning period showed 96 per cent control of sciarid flies @ 100 mg/kg (Shamshad 2010). Babar et al. (2012) recommended use of deltamethrin, spinosad, trichlorfon, malathion and permethrin for the control of phorid flies in mushroom and conducted study to know effective insecticides against phorid flies in mushroom. The insecticides were deltamethrin 2.5 EC @ 0.4 ml/l, spinosad 240 SC @ 0.4 ml/l, spinoteram 120 SC @ 0.2 ml/l, trichlorfon 80 WP @ 6.8 ml/l, malathion 57 EC @ 7.5 ml/l and permethrin 0.5 per cent @ 0.25 ml/l. The emergence and damage rate were reduced due to use of above insecticides in mushroom cultivation. Compared to other insecticides, spinosad was shown effective by maximum reduction of adult emergence and less damage rate by phorid flies. Shah *et al.* (2013) evaluated five fungicides such as carbendazim, bitertanol, hexaconazole, captan and mancozeb in vitro against greenmould disease and found that the maximum control of green mould disease occurred in carbendazim (90.8%), followed by bitertanol (40.0%), captan (36.6%) and hexaconazole (16.1%). Mancozeb had the least amount of control (11.7%).

Fungicides including prochloraz, benomyl, carbendazim and mancozeb were used for the control of fungal diseases. Prochloraz was the only one that is registered in Turkey for cultivated

mushrooms (Cengiz et al. 2014). Mortality of mushroom pests were observed maximum when treated with imidacloprid (88.88%) followed by dichlorvos (88.09%), thiamethoxam (61.10%) and malathion (35.71%) at 0.01 per cent (DMR 2015-16). Dicofol 50 EC @ 1-2 ml/l was to be sprayed at regular interval of time on the wall of mushroom house and also in compost for the control of mites. Control of springtails can be done by spraying malathion 0.05 per cent as disinfection spray (Raypuriya et al. 2018). Tian et al. (2020a) recommended use of pesticides such as fenprothrin, cyhalothrin, permethrin, cyfluthrin, cypermethrin, flucythrinate, fenvalerate and deltamethrin for effective control of sciarid and cecid flies. Application of permethrin dust (10 g a.i./kg) had controlled mushroom flies (sciarids, phorids and cecids) effectively. Aerosol formulations containing the insecticide active component pyrethrin or a mixture of pyrethrin and resmethrin may be sprayed or fumigated within mushroom manufacturing houses. It should be applied with "cold fogging" tools. A spray application with a maximum concentration of 700 ml of product per 100 litres of water should be used to apply a product containing 10 g/l pyrethrins for the control of mushroom flies (Okigbo and Anuagasi 2021). Permethrin dust (10 g a.i./kg) was used to kill mushroom flies such as sciarids, phorids and cecid flies. Spraying of fenvalerate and deltamethrin (0.05%) on walls, floor and galleries has effectively checked the mushroom adult flies. Applying diazonin 20 EC (1.5-2 ml/10 lt) or dicofol (0.1%) in compost or drenching mushroom houses and their surroundings with diazonin will also helps to control mites. Effective control of cecid fly prior to casing was achieved by applying chlorfenviphos, fenitrothion, and fenthion at 1 g a.i./m² to beds just after spraying (Okigbo and Anuagasi 2021).

4. Common instrumental techniques for analysis

4.1. Sample preparation

About 600 gm - 1 kg of mushroom samples were chopped, mixed and homogenized (Du et al. 2013; Cao et al. 2016; Tian et al. 2020a; Le et al. 2021). In that, only 100 gm is taken for as sub-sample for test purpose. Samples were stored at -18 to 20 °C in freezer for further analysis.

4.2. Sample extraction

The usage of the separation technique depends on type of sample and its any potential matrix interfering during extraction. The most common method used in mushroom for pesticide analysis was LC-MS/MS (Du et al. 2013; Chang et al. 2014; Du et al. 2014; Tian et al. 2015; Lu et al. 2017; Tian et al. 2020a; Le et al. 2021). It is necessary to take into account the physicochemical characteristics of the analyte and also the polarity of the pesticide. The complexity of sample treatment has been reduced while the accuracy and precision of the analysis have grown because to advancements in analytical techniques and extraction procedures. Using acetonitrile and petroleum ether, a multi-residue approach for analyzing organochlorine pesticides in food was created in 1963. A method based on acetone followed by dichloromethane and petroleum ether partitioning and clean-up with Florisil to be able to analyze more polar pesticides than the organochlorine group. The Dutch Food and Consumer Products Safety Authority-Food Inspection Service also created an extraction technique based on acetone in 1983, and it was regularly used for pesticide monitoring for more than 25 years. In 1989, the Swedish National Food Administration created an analytical procedure involving ethyl acetate and a clean-up step utilizing gel permeation chromatography. Since ethyl acetate has a lower polarity index (4.4) than acetone (5.1), polar insecticides partition less readily in ethyl acetate. The water phase is heavily supplemented with anhydrous sodium sulphate (Na_2SO_4) in order to force the polar insecticides into the organic solvent. A novel method based on acetonitrile extraction and clean-up employing dispersive solid phase extraction (dSPE) with primary and secondary amines (PSA) and

octadecylsilyl (C₁₈) was introduced in 2003 by Anastassiades *et al.* This sample treatment method was dubbed QuEChERS (Quick, Easy, Cheap, Effective, Rugged, Safe) by the researchers. This technique gained popularity since it used fewer conventional analytical stages, solvents and glassware. As a result, two reference methods were published: the first was CEN15662 by the European Committee for Standardization, which used acetonitrile and a citrate buffer during the extraction, and the second was "Method 2007.01" by AOAC International, which used acetonitrile and an acetate buffer during the extraction. Many methods have been published for the analysis of pesticides in mushrooms based on QuEChERS methodology. During the year from 2013 to 2022 (Du et al. 2013; Chang et al. 2014; Tian et al. 2015; Lu et al. 2017; Tian et al. 2020a; Le et al. 2021), method standardization was carried out by using QuEChERS method (ACN as main solvent) along with LC-MS/MS mostly. Overview of published literatures are presented in Table 1. In general, common procedure to analyse a large number of pesticide residues in mushroom uses Acetonitrile as organic solvent (Extraction solvent).

Table 1. Overview of published multi-residue methods for the analysis of pesticides in mushroom. Recovery is expressed in percentage. Repeatability is expressed as relative standard deviation percentage

Matrix	Pesticides	Extraction	Clean up	Method	Column	LOQs	LODs	Linear range	Recovery (%)	Reference
Fresh mushroom (Crimini, shiitake and oyster) Substrates	Pyriproxifen, avermectin and diflubenzuron	LPE with ACN	SPE with MgSO ₄ +PSA	UPLC-MS/MS	C-18	0.052 to 5 µg/kg	0.016 –1.5 mg	0.5, 1 or 5 to 5000 µg/l	78.1-112.5	Du et al. 2013

Shiitake mushroom, enoki mushroom and black edible fungus	187 pesticides	LPE with ACN	SPE with GCB	LC-MS/MS	C-18	0.02–170 µg/kg	0.01 to 85 µg/kg	0.05–33 920 µg/kg	70–118	Chang et al. 2014
Edible mushrooms and substrates (oyster, shiitake, eryngii and crimini)	Phoxim, chlorpyrifos and pyridaben	LPE with ACN	MgSO ₄ +PSA	UHPLC-MS/MS	Waters ACQUITY UHPLC BEH C18	1–10 µg/kg	1 to 10 µg/kg	0.01 to 1.0 µg/kg	75.5 to 98.4	Tian et al. 2015
Edible fungus	28 OP compounds	LPE with ACN	SPE MgSO ₄ +PSA	GC-MS	DB-1701 MS fused-silica	1.0 to 10 µg/kg	0.3 to 3.0 µg/kg	3.0 to 1000 µg/kg	85.26 to 100.21	Nan et al. 2015
Mushrooms (<i>Lentinula edodes</i>)	62 Pesticides	LPE with ACN	SPE with PSA + C18 + GCB + MgSO ₄	GC-MS/MS	DB-5MS UI column	0.625 to 20 µg/kg		10, 50, and 100 g/kg	75.4 to 117.1	Cao et al. 2016
Button, enoki, oyster, elm oyster and shiitake mushroom	clothianidin, dinotefuran, imidacloprid, thiacloprid, and thiamethoxam	LPE with ACN	PSA+C18+GCB	UHPLC-MS/MS	Agilent ZORBAX Eclipse Plus C18 column	0.1 to 2 µg/kg	0.03 to 0.7 µg/kg	10, 50 and 500 µg/kg	73.9 – 89.5	Lu et al. 2017
Edible mushrooms	flubendiamide, chlorantraniliprole, cyclaniliprole, tetrachlorantr	LPE with ACN	PSA+C18+GCB+MgSO ₄	HPLC-MS/MS	Agilent Poroshell 120 EC-C18	5 µg/kg	0.05-2 µg/kg	0.005 to 0.5 mg/l	73.5 – 110.2	Tian et al. 2020a

	aniliprole and cyantraniliprole				column					
Oyster mushroom, shiitake mushroom, eryngii mushroom, crimini mushroom, enoki mushroom and bunashimeji mushroom	ten pyrethroid (Bifenthrin, fenpropathrin, cyhalothrin, permethrin, cyfluthrin, cypermethrin, flucythrinate, tau-fuvalinate, fenvalerate and deltamethrin	LPE with ACN	SPE with PSA+C18 + MgSO ₄	GC-MS/MS	Agilent Technologies Capillary Column HP-5MS phenylmethylsiloxane fused-silica capillary analytical column	0.051 to 5.57 µg/kg	0.015 to 1.67 µg/kg	10, 100, and 1000 µg/kg	72.8 to 103.6	Tian et al. 2020b
Enoki, button, oyster and shiitake	180 pesticides	LPE with ACN	SPE with PSA + C18	UPLC-MS/MS	ACQUITY UPLC BEH C18 (1.7 µm, 100 × 2.1 mm) column	0.02–0.17 µg/kg	0.01–0.85 µg/kg	0.05–33 920 µg/kg	70 to 120	Le et al. 2021

4.3. Clean-up extraction method

The preliminary extraction using organic solvents is typically followed by a clean-up step. Various methods for this process are described below.

The most common clean-up agents used in extraction mushroom is PSA, GCB, C₁₈ and MgSO₄ (Tian et al. 2016; Lu et al. 2018; Tian et al. 2020a; Le et al. 2021). Residue behavior of six pesticides in button mushroom was analyzed using UPLC-MS/MS after performing extraction by QuEChERS method. A 10 g of sample was weighed into a 50 ml centrifuge container. Two ml of water and 10 ml of acetonitrile were then added for LC-MS/MS whereas for GC-MS/MS detection 10 ml of ethyl acetate was used followed by addition of 4 g of MgSO₄ (LC-MS), 3 g MgSO₄ (GC-MS) and 1.0 g of NaCl were then added. The containers were immediately sealed, vigorously spun for three minutes, and then centrifuged at 3500 RCF (Relative Centrifugal Force) for five minutes. 1.5 ml of the acetonitrile (upper) layer was decanted in a centrifuge tube containing 100 mg of anhydrous MgSO₄ and 50 mg of PSA. For GC analysis, 20 mg of florisil was used. After being spun for one minute, the materials were centrifuged at 3500 RCF for five minutes. A 0.22 µm nylon syringe filter was used to filter the prepared sample's top layer, which was then moved to an auto-sampler for analysis (Du et al. 2013). For determination of five neonicotinoid residues in mushroom, 10g of homogenized edible mushroom was taken in 50 ml conical beaker after which 1.0 ml of the working solution was added. The sample was left for 20 minutes to enable the solvent to evaporate. The extraction fluid contained 15.0 ml of acetonitrile and 5 millilitres of distilled water. Following that, the beaker was placed on an electric orbital shaking incubator and agitated for 15 minutes at room temperature for 150 rpm. The entire contents of the beaker were then filtered under vacuum into a 100 ml measuring cylinder that contained 5 g of NaCl. The cylinder was then manually shaken for about 2 minutes before being left on the table for 1 hour to enable phase separation. A 5 ml centrifuge tube containing 150 mg of anhydrous MgSO₄ and a specific

quantity of sorbent was then filled with 1.5 ml of the acetonitrile phase and vortexed for 30 seconds before being centrifuged at 5000 rpm for 5 minutes. In order to prepare for UHPLC-MS/MS analysis, 0.6 ml of the supernatant was then removed from the container using an SGE glass syringe. It was then combined with 1.0 ml of the mobile phase (20% methanol + 80% ultrapure water containing 0.1% formic acid) and passed through a 0.22- μ m nylon syringe filter (Lu et al. 2018). Five diamides such as flubendiamide, cyclaniliprole, chlorantraniliprole, tetrachlorantraniliprole and cyantraniliprole residue extraction in edible mushrooms was carried out by QuEChERS method. An aliquot of 10 g homogenized sample was taken in 50 ml of centrifuge tube. Acetonitrile (10 ml) was used as solvent for extraction. Next, salt mixture containing 1 g of NaCl and 4g of MgSO₄ was added followed by vortexing for one minute and centrifugation done at 2077 rpm for 5 minutes. In the cleanup step, for oyster mushroom, enoki mushroom, bunashimeji mushroom and crimini mushroom (20 mg PSA and 30 mg C18), 20 mg PSA and 50 mg C₁₈ for shiitake mushroom and eryngii mushroom was added. To all samples 150 mg of MgSO₄ was too added. The supernatant was decanted (0.9 ml) into an injection vial with a 0.22 μ m organic syringe filter for HPLC-MS/MS analysis (Tian et al. 2020b). Estimation of 180 pesticide residues in mushroom was performed by applying QuEChERS method. A 2 g of homogenized sample was taken in 50 ml of centrifuge tube. To that, 10ml of acetonitrile (ACN) was added then vortexed for 1 minute and below 40°C sonification was done for 15 minutes. Later, 1 g of NaCl, 4 g of MgSO₄, 0.35 g of trisodium citrate dihydrate and 0.2 g of disodium citrate were added. Again vortexed for 1 minute and centrifugation was done at 20,000 rpm. C18 and PSA was used for solid phase extraction. The solid phase extraction device was used to load 5 ml of the extract, which was then passed through the column before being collected. The elution solvent, which combines the sample extract and eluting solution, was 20 ml of the ACN: toluene

(3/1, v/v) solvent system. Using turbo evaporator at 40°C, 10 mbar, and 50 minutes, the solution acquired through SPE extraction was concentrated to 1 ml of enriched solution. After refinement, one millilitre of the extract was dried using nitrogen gas flow at one degree, then redissolved in one millilitre of ACN:H₂O. Finally, the sample filtered by a 0.22 µm PTFE barrier and injected to UPLC-MS/MS (Le et al. 2021).

4.4. Method validation parameters


Multiresidue analysis in mushroom was carried out in many countries such as China, Japan, Turkey, Veitnam, U.S.A, Poland etc. Method validation is done to assess parameters such as sensitivity, precision, linearity and recovery. QuEChERS method is sensitive, selective and accurate for determining multiple pesticide residues, so it is the mostly used method. Du *et al.* (2013) studied detection of pesticide residues (pyriproxyfen, diflubenzuron, and avermectin) in different mushrooms and their substrates using QuEChERS (Quick, Easy, Cheap, Effective, Rugged, and Safe) by Ultra-Performance Liquid Chromatography tandem Mass Spectrometry (UPLC-MS/MS). Relative Standard Deviation (RSD) values for all analytes were below 11.8 per cent, with average recoveries ranging between 78.1 and 112.5 per cent. The Limit of Detection (LOD) and Limit of Quantification (LOQ) were 0.016 to 1.5 g/kg and 0.052 to 5 g/kg, respectively. A quick and accurate Liquid Chromatography-tandem Mass Spectrometry (LC-MS/MS) method was applied for the simultaneous determination of 187 pesticide residues in shiitake mushroom, enoki mushroom and black edible fungus. The linear correlation coefficient r^2 was ≥ 0.990 for each pesticide's linear range. Recovery rates varied from 70 to 118 per cent at both low and high fortification levels. The RSD for 169 pesticides shown below 25 per cent which accounts for 90 per cent of the total. LOD ranged from 0.01 to 85 µg/kg for 165 pesticides (Chang et al. 2014). The determination of prochloraz residues were estimated using QuEChERS method

combined with Gas Chromatography Electron Capture Detector (GC-ECD). This method has shown effective linear relationship with r^2 (0.999) for concentration 52.9 to 1000 $\mu\text{g}/\text{kg}$. The LOD and LOQ were 15.9 $\mu\text{g}/\text{kg}$ and 52.9 $\mu\text{g}/\text{kg}$ respectively. Mushroom samples had shown recovery of 100.6 ± 1.6 per cent for 250 $\mu\text{g}/\text{kg}$ and 105.6 ± 3.9 per cent for 1000 $\mu\text{g}/\text{kg}$ (Cengiz et al. 2014). Tian et al. (2016) developed modified QuEChERS method for detection of phoxim, chlorpyrifos and pyridaben residues in edible mushrooms and substrate using UHPLC-MS/MS and Electro-Spray Ionisation source operating in the positive mode (ESI+) to identify the target molecules. For the three compounds, LOQ were 10 $\mu\text{g}/\text{kg}$, and LOD varied from 1 to 10 $\mu\text{g}/\text{kg}$ with correlation values >0.994 achieved, using calibration curves of standards ranging from 0.01 to 1.0 $\mu\text{g}/\text{kg}$. Recovery of 75.5 to 98.4 per cent for concentration 0.01-0.25 mg/kg was noted. A new analytical method was developed using Gas Purge Micro-Syringe Extraction (GP-MSE) and Gas Chromatography-Mass Spectrometry (GC-MS) for detection of 28 organophosphorus pesticide residues in edible fungus. Organophosphorus insecticide showed recovery ranging from 85.26 to 100.21 per cent and RSD ranged from 1.6 to 6.9 per cent. The LOQ and LOD from 1.0 to 10 $\mu\text{g}/\text{kg}$ and 0.3 to 3.0 $\mu\text{g}/\text{kg}$ respectively (Nan et al. 2015). Based on a modified QuEChERS sample preparation and employing GC-MS/MS, a quick analytical approach for the simultaneous detection of 62 pesticides in mushrooms was developed. The pesticides were quantified using the internal standard curve method after the cleaned extracts were subjected GC-MS/MS analysis in the Multi-Reaction Monitoring (MRM) mode. Recoveries at three concentrations (10, 50, and 100 g/kg) varied for all the chemicals from 75.4 to 117.1 per cent, with RSD between 3.2 and 15.2 per cent and LOQ were 0.625 to 20 $\mu\text{g}/\text{kg}$ (Cao et al. 2016). Five neonicotinoide insecticides (clothianidin, dinotefuran, imidacloprid, thiacloprid, and thiamethoxam) in mushroom (button, enoki, oyster, elm oyster and shiitake mushroom) were detected by UHPLC-MS/MS. The samples

were processed using a modified QuEChERS procedure. For all analytes found in mushroom, LOD ranged from 0.03 to 0.7 µg/kg, whereas LOQ ranged from 0.1 to 2 µg/kg. The ranges of the mean recoveries for clothianidin, dinotefuran, imidacloprid, thiacloprid and thiamethoxam were 100.5-118.0, 73.9-89.5, 88.6-117.8, 72.9-121.8 and 98.9-117.2 per cent respectively with RSD less than 8.1 per cent for 10, 50 and 500 µg/kg concentration (Lu et al. 2018). A modified QuEChERS technique for the simultaneous determination of flubendiamide, chlorantranilprole, cyclanilprole, tetrachlorantranilprole and cyantranilprole in edible mushrooms was developed applying HPLC-MS/MS. By plotting the peak area against the corresponding concentration at six calibration levels, 0.005 to 0.5 mg/l, the linearity for all the target chemicals in acetonitrile and matrix solutions was determined and found satisfactory ($R^2 \geq 0.99$). The LOD was 0.05-2 µg /kg and LOQ was 5 µg/kg. These pesticides had recoveries that were acceptable (73.5–110.2%) for concentration 5, 10 and 100 µg /kg. The RSD was below 12.7 per cent (Tian *et al.*, 2020b). A technique was formed to simultaneously determine the presence of ten pyrethroid (Bifenthrin, fenpropathrin, cyhalothrin, permethrin, cyfluthrin, cypermethrin, flucythrinate, tau-fuvalinate, fenvalerate and deltamethrin) insecticide residues in six different edible mushrooms (oyster, shiitake, eryngii, crimini, enoki and bunashimeji). The samples were prepared using QuEChERS method combined with GC-MS/MS. Six different edible mushroom species were used to extract the 10 pyrethroid pesticides. At three levels (10, 100, and 1000 µg/kg), the average recoveries for the six different edible mushrooms varied from 72.8 to 103.6 per cent. The RSD were less than 13.0 per cent for both intraday and interday data. The LOQ were < 5.57 µg/kg (Tian et al. 2020a). Le et al. (2021) developed a highly sensitive and reliable approach to simultaneously identify 180 pesticides in different mushrooms (Enoki, button, oyster and shiitake) through the optimization of extract purification conditions. By combining QuEChERS extraction with a mixed mode of

SPE (Solid Phase Extraction) cleaned up with various adsorbent materials after sample preparation, matrix effects were minimized. The LOQ varied from 2 to 5 $\mu\text{g}/\text{kg}$, which is significantly less than the MRLs set by the EU (10 - 50 $\mu\text{g}/\text{kg}$). Recovery rates ranged from 70 to 120 per cent and the RSD of repeatability and reproducibility were both less than 20 per cent. The methodology for pesticide residue analysis from sample preparation to analysis was shown in Fig 1.

Fig 1. Methodology for analysis of pesticide residue analysis in mushroom



5. Monitoring of pesticides

Pesticides are widely used in order to prevent and control diseases and insect pests during mushroom cultivation. Monitoring of pesticide residues in mushroom is thus becomes essential in order to prevent health hazards. The use of pesticides was reduced as a result of the application

of integrated pest management concepts and excellent agricultural practices, with a tendency to reduce the most hazardous pesticides. A number of studies related to monitoring were conducted. In one of the study, Jiang et al. (2012) monitored 33 pesticide residues in a total of 4 fruiting body samples such as oyster mushroom, shiitake mushroom, button mushroom and fungus collected from the four provinces of Hubei, Zhejiang, Henan, and Fujian. The study showed that, 23.3 per cent of the samples exceeded the EU standard, 3.27 per cent of the samples above the Japanese standard while no sample exceeded the current level in China. Pesticides such as cymethrin, cyflumethrin, thiophenate methyl, methophos, methyl parathion, malathion and bethion were primarily found in the samples. Bethion was found in seven samples of button mushrooms, oyster mushrooms, and fungus. Cyflumethrin was found in seven samples of oyster mushrooms. Thiophenate methyl was found in six samples of button mushrooms. Individual samples of button mushroom contained highly toxic pesticides like methamidophos, monocrotophos and methylparathion. Galgowska et al. (2012) examined two mushroom species such as *Xerocomus* mushroom (*Xerocomus badius*) and *Boletus* mushroom (*Boletus edulis*) from Poland for the presence of organochlorine compounds using GC-MS. γ -HCH, DDT, and its metabolites (DDE, DDD) were found in all of the samples that were analyzed. *Xerocomus* mushrooms had the highest concentration of γ -HCH (0.125 $\mu\text{g}/\text{kg}$ on average), while *boletus* mushrooms had a lower concentration of 0.11 $\mu\text{g}/\text{kg}$ of mushrooms. Tian et al. (2016) monitored 50 mushroom samples for presence of pesticide residues such as pyridaben, chlorpyrifos and phoxim. Mushroom samples such as oyster, shiitake, button mushroom were collected from Beijing local market of China. In button mushroom, pyridaben was found in the range of 0.08 to 0.015 mg/kg. In Western China, pesticide residue study was conducted to the know amount of pesticide residues in edible mushroom. Around 354 mushroom samples were brought from nearby markets and analyzed 53

pesticides using HPLC/MS/MS. Major pesticides which were found in the edible mushroom samples included carbendazim, acephate, procymidone, prochloraz, aldicarb sulfone etc. The detection rates for carbendazim, acephate and procymidone were 0.0002-2.7316 mg/kg, 0.0248-0.4985 mg/kg and 0.1807-0.3928 mg/kg respectively. Highest detection was noted with carbendazim (70.9%) followed by acephate (13%) and procymidone (7.3%). While the remaining samples of edible mushroom revealed one or more pesticide residues (Li et al. 2022).

6. Future Directions and challenges

Regulation and Monitoring with Stricter regulations and enhanced monitoring are expected to be implemented to ensure that pesticide residues in mushrooms remain within safe limits. This may involve more frequent testing and tighter control measures. Alternative Pest Management, there is a growing interest in developing and adopting alternative pest management strategies, such as biological controls and integrated pest management (IPM), which could reduce reliance on chemical pesticides and thus minimize residues. Organic and Sustainable Farming, the trend towards organic and sustainable farming practices is likely to continue. Organic mushroom cultivation typically uses fewer or no synthetic pesticides, which can reduce the presence of pesticide residues. Advancements in Detection Technology such as more sensitive analytical methods, will enhance the ability to identify and quantify pesticide residues more accurately. This will aid in better compliance and safety assessments. Reducing matrix effect by using suitable clean-up agents and optimizing chromatographic conditions (e.g., adjusting column type, mobile phase composition, and flow rates) to improve separation and reduce matrix interference in techniques like HPLC or GC (Anastassiades et al. 2013). Consumer Awareness and Preferences, As consumers become more aware of pesticide residues and their potential health impacts, there will be an increasing demand for mushrooms grown with minimal pesticide use, potentially

influencing farming practices. Research and Development, ongoing research into the effects of pesticides on mushrooms and their environment will contribute to understanding and managing residues better. This includes studying how pesticides break down and affect mushroom safety. Overall, the future will likely see a combination of regulatory, technological, and consumer-driven changes aimed at minimizing pesticide residues in mushrooms and improving food safety (Stalikas and Konidari 2001).

7. Limitations of multiresidue methods

The most common and efficient way to perform pesticide residue analysis for numerous different compounds involves using multi-residue methods that can measure within the MRL range from 0.01 to 10 mg/kg (Anastassiades et al., 2013). However, these multi-residue methods cannot measure all pesticides with the necessary accuracy in a single run. The significant diversity in the chemical composition of these hundreds of pesticides hinders the application of a single strategy for their simultaneous analysis. Consequently, it is sometimes still essential to develop single-residue methods for the analysis of one pesticide or a few pesticides from the same chemical family. Examples include pesticides with high polarity or ionic characteristics. Another issue might be the low stability of specific pesticides during sample extraction. In the case of compounds with a high polarity or ionic compounds, new approaches based on LC have been proposed. They can be divided in three strategies: (i) the polarity of the analytes is reduced by derivatization of the analytes or by addition of an ion pairing substance to the mobile phase before analysis by RP-LC. This decreased polarity leads to an increased retention and more adequate peak shape (Stalikas and Konidari 2001). (ii) Use of hydrophilic interaction liquid chromatography (HILIC) with carbon or ion exchange phases instead of reversed phases. Also

this leads to an increased retention of the analytes (Bernal et al. 2011). (iii) Elimination of the separation technique and use of direct flow injection (FI) to MS/MS.

Commented [1]: Include more latest references and discussion part

8. Conclusions

Mushrooms are recognized for their significant nutritional value and medicinal properties, making them a valuable addition to the diet and a popular supplement to cereals. Their rich content of proteins, minerals, vitamins, and bioactive compounds contributes to their growing market demand. However, the cultivation of mushrooms faces challenges, particularly from insect pests, which often leads to the extensive use of pesticides. While these pesticides are essential for controlling pests and diseases, their residues on mushrooms pose potential risks to food safety and marketability. The increase in mushroom production and international trade emphasizes the need for stringent monitoring of pesticide residues to ensure compliance with food safety standards. Advances in analytical methods, particularly multiresidue techniques, are crucial for accurately detecting and quantifying pesticide residues despite the complex matrix of mushrooms. These methods help mitigate the risk of pesticide contamination and support the export of safe, high-quality mushrooms. Efforts to reduce pesticide use and improve residue analysis are essential for maintaining food safety and enhancing the quality of mushroom products. By adopting effective pest management practices and utilizing advanced analytical techniques, the mushroom industry can address the challenges of pesticide residues and continue to thrive in both domestic and international markets

References:

Amin, Z., Wani, F. F., Gulzar, H., Dar, W., & Sheikh, P. J. (2021). Diseases of White Button mushroom (*Agaricus bisporus*) - A potential threat to mushroom industry. *International Journal*

of *Current Microbiology and Applied Sciences*, 10(2), 2076-2085.
<https://doi.org/10.20546/ijcmas.2021.1002.247>

Anastassiades, D., Kolberg, M., Mack, D. I., Wildgrube, D., Sigalov, C., & Dork, I. (2013). Quick method for the analysis of residues of numerous highly polar pesticides in foods of plant origin involving simultaneous extraction with methanol and LC-MS/MS determination (QuPPE-Method). *EURL-SRM: 434*. Available from http://www.crl-pesticides.eu/library/docs/srm/meth_QuPPE.pdf.

Anastassiades, M., Lehotay, S. J., Stajnbaher, D., & Schenck, F. J. (2003). Fast and easy multiresidue method employing acetonitrile extraction/partitioning and dispersive solid-phase extraction for the determination of pesticide residues in produce. *Journal of AOAC International*, 86(2), 412-421. <https://doi.org/10.1093/jaoac/86.2.412>

Anastassiades, M., Scherbaum, E., Tasdelen, B., & Stajnbaher, D. (2007). Recent developments in QuEChERS methodology for pesticide multiresidue analysis. In *Pesticide Chemistry: Crop Protection, Public Health, Environmental Safety* (pp. 439-458). <https://doi.org/10.1002/9783527611249>

Babar, M. H., Ashfaq, M., Afzal, M., Bashir, M. H., & Ali, M. A. (2012). Efficacy of different insecticides against mushroom phorid fly, *Megaselia halterata* (Wood) in Punjab, Pakistan. *Pakistan Journal of Nutrition*, 4(1), 183-188. <https://doi.org/10.3923/pjn.2014.50.55>

Bernal, J., Ares, A. M., Pól, J., & Wiedmer, S. K. (2011). Hydrophilic interaction liquid chromatography in food analysis. *Journal of Chromatography A*, 1218(42), 7438-7452.
<https://doi.org/10.1016/j.chroma.2011.05.004>.

Bijla, S. (2023). Status of mushroom production: Global and national scenario. *Mushroom Research*, 32(2). <https://doi.org/10.36036/mr.32.2.2023.146647>

Broadley, A., Kauschke, E., & Mohrig, W. J. (2018). Black fungus gnats (Diptera: Sciaridae) found in association with cultivated plants and mushrooms in Australia, with notes on cosmopolitan pest species and biosecurity interceptions. *Zootaxa*, 4415(2), 201-242. <https://doi.org/10.11646/zootaxa.4415.2.1>

Cao, X., Liu, S., Yang, X., Liu, Z., & Liu, L. J. (2016). A modified QuEChERS sample preparation method for simultaneous determination of 62 pesticide residues in edible fungi using gas chromatography–triple quadrupole mass spectrometry. *Food Analytical Methods*, 9(1), 263-274. <https://doi.org/10.1007/s12161-015-0200-0>

Castilho, R. C., de Moraes, G. J., Silva, E. S., Freire, R. A., & Eira, F. C. (2009). The predatory mite *Stratiolaelaps scimitus* as a control agent of the fungus gnat *Bradysia matogrossensis* in commercial production of the mushroom *Agaricus bisporus*. *International Journal of Pest Management*, 55, 181-185.

Cengiz, M. F., Catal, M., Erler, F., & Bilgin, A. K. (2014). Rapid and sensitive determination of the prochloraz residues in the cultivated mushroom, *Agaricus bisporus* (Lange) Imbach. *Analytical Methods*, 6(6), 1970-1976. <https://doi.org/10.1039/c3ay42293c>

Chang, Q., Fan, C., Chen, H., Kang, J., Wang, M., & Pang, G. (2014). Determination of 187 pesticide residues in edible fungi by liquid chromatography-tandem mass spectrometry. *Analytical Methods*, 6(12), 4288-4304. <https://doi.org/10.1039/C3AY42213E>.

Chen, C., Qian, Y., Chen, Q., Tao, C., Li, C., & Li, Y. (2011). Evaluation of pesticide residues in fruits and vegetables from Xiamen, China. *Food Control*, 22(7), 1114-1120. <https://doi.org/10.1016/j.foodcont.2011.01.007>

Cheung, P. C. K. (2010). The nutritional and health benefits of mushrooms. *Nutrition Bulletin*, 35(4), 292-299. <https://doi.org/10.1111/j.1467-3010.2010.01859.x>

Daba, A. S., Kabeil, S., Botros, W. A., & El-Saadani, M. (2008). Production of mushroom (*Pleurotus ostreatus*) in Egypt as a source of nutritional and medicinal food. *World Journal of Agricultural Sciences*, 4, 630-63. <https://doi.org/10.1016/j.bbrep.2020.100852>

Das, P., Dutta, S. K., Puzari, K. C., Sarmah, G. H., & Dutta, P. (2014). First report of fungus gnat (*Lycoriella* sp.) (Diptera: Sciaridae) on oyster mushroom from North East India. *International Journal of Plant Protection*, 7, 78-81.

Directorate of Mushroom Research. (2009). DMR – 2008-09, Annual Report. Available from https://dmrsolan.icar.gov.in/AR_2008-09.pdf.

Directorate of Mushroom Research. (2016). DMR – 2015-16, Annual Report. Available from <https://dmrsolan.icar.gov.in/DMRAnnualReport2015-16.pdf>.

Du, P., Liu, X., Gu, X., Dong, F., Xu, J., Kong, Z., & Zheng, Y. (2013). Rapid residue analysis of pyriproxyfen, avermectins, and diflubenzuron in mushrooms by ultra-performance liquid chromatography coupled with tandem mass spectrometry. *Analytical Methods*, 5(23), 6741-6747. <https://doi.org/10.1039/C3AY41074A>.

Duhan, K., Gulati, R., & Malik, A. J. (2018). Damage potential of *Tyrophagus putrescentiae* Schrank (Acari: Acaridae) in oyster (*Pleurotus sajor*) mushrooms. *Emerging Life Sciences Research*, 4, 40-44.

Erler, F., & Polat, E. (2015). The flies on mushrooms cultivated in the Antalya-Korkuteli district and their control. *Akdeniz University Journal of the Faculty of Agriculture*, 28, 61-66.

European Union MRL DATABASE. (2023). Retrieved from <https://ec.europa.eu/food/plant/pesticides/eu-pesticides-database/start/screen/products/details/197> (Accessed 20 June 2023).

FAO. (2018). Retrieved October 2021, from <http://www.fao.org/faostat/en/#data/TM>

FAOSTAT. (2023). Retrieved from <http://www.fao.org/faostat/en/#data/TM>

Fletcher, J. T., & Gaze, R. H. (2007). *Mushroom pest and disease control: A color handbook*. Elsevier.

Galgowska, M., & Pietrzak-Fiecko, R. (2017). Pesticide contaminants in selected species of edible wild mushrooms from the north-eastern part of Poland. *Journal of Environmental Science and Health, Part B*, 52(3) 214-217. <https://doi.org/10.1080/03601234.2017.1261553>

Grewal, P. S. (2007). Mushroom pests. In *Field Manual of Techniques in Invertebrate Pathology: Application and Evaluation of Pathogens for Control of Insects and Other Invertebrate Pests* (pp. 457-461). Springer Netherlands Dordrecht. http://dx.doi.org/10.1007/978-1-4020-5933-9_21

Jess, S., & Kilpatrick, M. (2000). An integrated approach to the control of *Lycoriella solani* (Diptera: Sciaridae) during production of the cultivated mushroom (*Agaricus bisporus*). *Pest*

Management Science, 56(5), 477-485. [https://doi.org/10.1002/\(sici\)1526-4998\(200005\)56:5<477::aid-ps161>3.3.co;2-k](https://doi.org/10.1002/(sici)1526-4998(200005)56:5<477::aid-ps161>3.3.co;2-k)

Jiang, X., Liu, D. P., Fan, X., Li, W., & Bian, Y. (2012). Analysis of common pesticide residues in four main producing areas of edible fungi. *Edible Fungi China*, 31, 53-54. <https://doi.org/10.1016/j.bbrep.2020.100852>

Jiang, X., Liu, D., Fan, X., Li, W., & Bian, Y. (2012). Analysis of common pesticide residues in four main producing areas of edible fungi. *Edible Fungi China*, 31, 53-54.

Kakon, A. J., Choudhury, M. B., & Saha, S. (2012). Mushroom is an ideal food supplement. *Journal of Dhaka National Medical College & Hospital*, 18(1), 58-62. <https://doi.org/10.3329/jdnmch.v18i1.12243>

Keil, C. J. (2002). Field trials of the Gnatrol WDG formulation of *Bacillus thuringiensis israelensis* for control of mushroom flies in Pennsylvania and California. *Mushroom News-Kennett Square*, 50, 12-19.

Le, L. H. T., Tran-Lam, T. T., Cam, T. Q., Nguyen, T. N., & Dao, Y. H. (2021). Pesticides in edible mushrooms in Vietnam. *Food Additives & Contaminants: Part B*, 14(2), 139-148. <https://doi.org/10.1080/19393210.2021.1908434>

Lewandowski, M., Szyk, A., & Bednarek, A. (2004). Biology and morphometry of *Lycoriella ingenua* (Diptera: Sciaridae). *Biological Letters*, 41, 41-50. <https://doi.org/10.1016/j.bbrep.2020.100852>

Li, Y., Qin, G., He, F., Zou, K., Zuo, B., Liu, R., Zhang, W., Yang, B., Zhao, G., & Jia, G. (2022). Investigation and analysis of pesticide residues in edible fungi produced in the mid-western region of China. *Food Control*, 136, 108857. <https://doi.org/10.1016/j.foodcont.2022.108857>

Lu, Z., Fang, N., Zhang, Z., Wang, B., Hou, Z., Lu, Z., & Li, Y. (2018). Simultaneous determination of five neonicotinoid insecticides in edible fungi using ultrahigh performance liquid chromatography-tandem mass spectrometry (UHPLC MS/MS). *Food Analytical Methods*, 11(4), 1086-1094. <https://doi.org/10.1007/s12161-017-1080-2>.

Maurya, P. K., Malik, D. S., & Sharma, A. (2019). Impacts of pesticide application on aquatic environments and fish diversity. *Contaminants in Agriculture: Environmental Health Risks and Remediation*, 1, 111. <http://dx.doi.org/10.26832/aesa-2019-cae-0162-09>

Meena, N. K. (2021). *Pest and disease problems in mushroom cultivation*.

Munshi, N. A., Dar, G. H., Ghani, M. Y., Kauser, S., & Mughal, N. (2010). *Button mushroom cultivation*. Communication and Publication Centre SKUAST-K: 1-28.

Nan, J., Wang, J., Piao, X., Yang, C., Wu, X., Quinto, M., & Li, D. J. (2015). Novel and rapid method for determination of organophosphorus pesticide residues in edible fungus using direct gas purge microsyringe extraction coupled on-line with gas chromatography-mass spectrometry. *Talanta*, 142, 64-71. <https://doi.org/10.1016/j.talanta.2015.04.035>

Navarro, M. J., Escudero-Colomar, L. A., Carrasco, J., & Gea, F. J. (2021). Mushroom phorid flies - A review. *International Journal of Pest Management*, 11, 1958.

O'Connor, L., & Keil, C. J. (2005). Mushroom host influence on *Lycoriella mali* (Diptera: Sciaridae) life cycle. *Journal of Economic Entomology*, 98, 342-349. <https://doi.org/10.1093/jee/98.2.342>

Okigbo, R. C., & Anuagasi, C. (2021). Diseases affecting mushrooms in Africa. *Journal of Food Technology and Nutrition Science*, 129, 2-10. [https://doi.org/10.47363/jftns/2021\(3\)129](https://doi.org/10.47363/jftns/2021(3)129)

Pandey, V. V., Kumari, A., Kumar, M., Saxena, J., Kainthola, C., & Pandey, A. J. (2018). Mushroom cultivation: Substantial key to food security. *Journal of Applied and Natural Science*, 10(4), 1325-1331. <https://doi.org/10.31018/jans.v10i4.1941>

Raypuriya, N., Singh, Y., & Prajapati, A. S. (2018). Insect pest and disease management in mushroom. *Rashtriya Krishi*, 23-25.

Rijal, R., Maity, P., & Kumar, A. (2021). Pests of mushroom and their ecological management strategies: A review. *Biological Forum International Journal*, 13, 375-387.

Rinker, D. L. (2017). Insect, mite, and nematode pests of commercial mushroom production. In **Edible and Medicinal Mushrooms: Technology and Applications* (pp. 221-237). <https://doi.org/10.1002/9781119149446.ch11>

Ruchika, K., Upamanyu, S., Sharma, D., Choudhary, K., Yangchan, J., Kumari, P., & Kumar, S. (2021). Incidence and occurrence of insect pest associated with white button mushroom *Agaricus bisporus* (Lange) Imbach in Himachal Pradesh, India. *Mushroom Research*, 19(1) <https://doi.org/10.51220/jmr.v19-i1.10>

SANTE. (2021). Analytical quality control and method validation procedures for pesticide residues analysis in food and feed. *SANTE 11312/2021*. Retrieved from https://www.eurl-pesticides.eu/userfiles/file/EurlALL/SANTE_11312_2021.pdf (Accessed 06 March 2023).

Shah, S. N., & Nasreen, K. (2013). Efficacy of fungicides against *Trichoderma* spp. causing green mould of oyster mushroom (*Pleurotus sajor-caju*). *Research Journal of Microbiology*, 8(1), 13-24. <https://doi.org/10.3923/jm.2013.13.24>

Shamshad, A. (2010). The development of integrated pest management for the control of mushroom sciarid flies, *Lycoriella ingenua* (Dufour) and *Bradysia ocellaris* (Comstock), in cultivated mushrooms. *Pest Management Science*, 66(10), 1063-1074. <https://doi.org/10.1002/ps.1987>

Sharma, A., Khanna, A. S., Raina, R., Kapoor, R., & Thakur, K. S. (2019). Faunistic survey of insect-pests associated with *Agaricus bisporus*. *International Journal of Current Microbiology and Applied Sciences*, 6, 122-125.

Shin, S., Lee, H., & Lee, S. (2011). Two cecidomyiid gall midge (Diptera: Cecidomyiidae) pests of shiitake mushrooms (Agaricales: Marasmiaceae). *Journal of Asia-Pacific Entomology*, 14, 387-391. <https://doi.org/10.1016/j.aspen.2011.05.002>

Singh, A. U., & Sharma, K. (2016). Pests of mushroom. *Advances in Crop Science and Technology*, 4, 213.

Stalikas, C. D., & Konidari, C. N. (2001). Analytical methods to determine phosphonic and amino acid group-containing pesticides. *Journal of Chromatography A*, 907(1-2), 1-19. [https://doi.org/10.1016/s0021-9673\(00\)01009-8](https://doi.org/10.1016/s0021-9673(00)01009-8)

Stoddart, H., Garthwaite, D., Thomas, M. J. D., Department of Environment, Food, and Rural Affairs, & Scottish Executive Environment and Rural Affairs Department. (2004). Pesticide usage survey report 197: Mushroom crops in Great Britain 2003. *Department of Environment, Food and Rural Affairs and Scottish Executive Environment and Rural Affairs Department*.

Thongnaitham, M. (2012). *Organic mushroom cultivation manual*. Freeland Foundation: 591-556.

Tian, F., Liu, X., Xu, J., Dong, F., Zheng, Y., Hu, M., & Wu, Y. (2016). Simultaneous determination of phoxim, chlorpyrifos, and pyridaben residues in edible mushrooms by high-performance liquid chromatography coupled to tandem mass spectrometry. *Food Analytical Methods*, 9(10), 2917-2924. <https://doi.org/10.1007/s12161-016-0490-x>.

Tian, F., Qiao, C., Luo, J., Guo, L., Pang, T., Pang, R., Li, J., Wang, C., Wang, R., & Xie, H. (2020a). Method development and validation of ten pyrethroid insecticides in edible mushrooms by modified QuEChERS and gas chromatography-tandem mass spectrometry. *Food Chemistry*, 10(1), 7042. <https://doi.org/10.1016/j.foodchem.2020.127468>.

Tian, F., Qiao, C., Luo, J., Guo, L., Pang, T., Pang, R., Li, J., Wang, C., Wang, R., & Xie, H. (2020b). Development and validation of a method for the analysis of five diamide insecticides in edible mushrooms using modified QuEChERS and HPLC-MS/MS. *Food Chemistry*, 333, 127468. <https://doi.org/10.1016/j.foodchem.2020.127468>

Waktola, G., & Temesgen, T. (2018). Application of mushroom as food and medicine. *Advances in Biotechnology & Microbiology*, 11(4), 10-19080. <https://doi.org/10.19080/AIBM.2018.11.555817>

Ethical Approval: Not applicable

Consent to Participate: Not applicable

Consent to Publish: Not applicable

Data availability statement

All data generated or analyzed during this study are included in this article.

UNDER PEER REVIEW