

Original Research Article

**Etiologies and Factors associated with Febrile
Consciousness Disorders in People Living with HIV
(PLHIV), hospitalized in the Infectious and Tropical
Diseases Department of the University Hospital Center (CHU)
Donka, Guinea.**

Commented [a1]: Consider title case or any recommendation by the journal

Abstract :

Introduction

Damage to the central nervous system (CNS) is common in ~~HIV~~ the Human Immunodeficiency Virus (HIV) infection and occurs in cases of immunosuppression, which leads to altered consciousness. The objective of this work was to identify the etiologies and factors associated with the death of ~~HIV-positive patients~~ PLHIV hospitalized for febrile consciousness disorders in the infectious and tropical diseases department of Donka University Hospital.

Methodology: This was a five-years retrospective study of descriptive and analytical type lasting 5 years in the department of infectious and tropical diseases of the university hospital center of Conakry. Included in this study were HIV-infected patients, and -on ARV Anti-retroviral therapy (ART) or not, aged over 15 years, hospitalized in the infectious diseases department for febrile consciousness disorders or those who developed it the disorder during our the study period under review.

Commented [a2]: Write in full on first mention before subsequent abbreviation. This also include AIDS, HIV, WHO etc used elsewhere in this document

The parameters studied were sociodemographic characteristics, clinical picture, and clinical staging of HIV/AIDS according to the WHO World Health Organisation (WHO) standard. The Factors associated with deaths were searched from logistic regression in multivariate analysis and all variables with a value of $P < 0.05$ were considered statistically significant.

Results : ~~We collected~~ A total of 205/534 HIV+ patients hospitalized for febrile consciousness disorders were sampled and enrolled for this study, representing a prevalence of 38.39%. The average age of the patients was 42.28 ± 12.54 years with a sex ratio of 0.97. ~~Cerebral toxoplasmosis was the most common etiology. The depth of coma, the CD4 count ≤ 200 cells/mm³ and the absence of antiretroviral treatment were the factors associated with death in this study.~~

Commented [a3]: Avoid personalization. Used third party reporting instead, e.g. A total of 205/534 samples were collected. Again it was not the HIV+ patients that were collected ; report correctly what was collected, or instead say, were enrolled, not collected

Conclusion : Febrile consciousness disorders remain very common, with multiple etiologies in severely immunocompromised HIV+HIV-infected patients. The CD4 count ≤ 200 cells/mm³, the

Commented [a4]: This is repeated in the conclusion where it is most appropriate. Remove it from here

depth of the coma and the absence of antiretroviral treatment were the [most common](#) factors associated with death.

Keywords: Consciousness disorders, Fever, HIV+, Donka, Guinea

Introduction :

Febrile consciousness disorders are defined as any alteration of alertness accompanied by a temperature above 38°C of central origin or not. They can range from a state of simple obtundation or stupor to the stage of coma [1]. [On the other hand, HIV](#) is an infection caused by a retrovirus that attacks the body's immune system by destroying it, thus weakening the effectiveness of the patient's immune system [2]. Damage to the central nervous system (CNS) is common in HIV and [occurs often occur](#) in cases of immunosuppression, which leads to altered consciousness [3,4].

Cerebral toxoplasmosis still remains the first opportunistic infection most frequently encountered in AIDS, followed by neuromeningeal cryptococcosis and bacterial infections which are responsible for impaired consciousness [3,5].

Most studies carried out around the world show that the main factors associated with the death of PLHIV were late diagnosis of HIV, non-[compliance adherence to ART](#), ineffectiveness of [antiviral-ART](#) treatment, absence of [antiviral-ART](#) treatment, a CD4 count ≤ 200 cells / μ L, a Glasgow score less than 12 and the presence of opportunistic infections linked to HIV [6].

[According to UNAIDS, in 2022 the mortality rate among PLHIV was 26% in the USA; 25% in Europe and 48% in Africa \[7\].](#)

[Umgelter et al in their](#) study carried out in Germany found that PLHIV represented 10 to 30% of all adults admitted with disorders of consciousness [8].

[In Nigeria, Obiako et al A study in Nigeria equally](#) found similar results [9].

[Amir Mbondé et al in](#) Uganda, [according to theirs](#) study carried out in 2022, found that the absence of antiviral treatment represented 45% of the factors associated with death and the Glasgow score less than 12 represented 30% of the factors associated with death [10].

[Stepchenkova et al in their study carried out in](#) Ukraine, [a study carried out and published](#) in 2014 showed that late diagnosis of HIV and ineffectiveness of [antiviral-antiretroviral](#) treatment represented respectively 42% and 24% of the factors associated with death [11].

[However, in Guinea, little data exists on this subject, so it is in this spirit that we initiated this study, the objective of which was to:](#) [In Guinea however, little data exists about this subject, hence the justification for this study.](#)

Commented [a5]: Your paragraphs in this section are too short. Combine them and have at least 3-4 sentences per paragraph

Commented [a6]: Do not begin a paragraph with abbreviation.

Commented [a7]: Verify these figures, looks higher especially given the VL suppression reported globally

Objectives

Main Objective

Specific Objective

Research Questions

Identify the etiologies and factors associated with the death of HIV-positive patients hospitalized for febrile disorders of consciousness in the infectious and tropical diseases department of DONKA University Hospital.

Commented [a8]: List the objectives in the manner shown above ; objectives main objectives, specific objectives, and if possible research questions

1-Materials and methods:

Study Design : This was a five-year retrospective study of descriptive and analytical type-nature lasting 05 years from covering the period between July 1st, 2018 and to June 30th, 2024.

Study Site : ~~The study was~~ carried out in the infectious and tropical diseases department of the Conakry University Hospital center. ~~It is~~ Conakry University Hospital is the the reference service facility in the management of HIV infections in among adults in Guinea.

Inclusion Criteria : ~~Included in this work were HIV infected patients on ARV or not~~ This study included participants drawn from among the HIV infected male and female patients regardless of their ART status, aged over 15, ~~regardless of sex, profession and origin, and~~ hospitalized at in the Conakry university hospital infectious diseases department for febrile consciousness disorders as well as those who ~~had~~ developed ~~it: the disorder~~ during ~~our the~~ study period.

Commented [a9]: Even though we know they are patients, for your study they are reported as participants and not patients

Sampling technique : ~~For data collection, we carried out an e~~ Exhaustive recruitment was done ~~of for~~ all patients participants meeting the selection criteria during the period considered under consideration. The parameters studied were: sociodemographic characteristics, clinical picture, clinical staging of HIV/AIDS according to the WHO, etiologies and factors associated with death in among the hospitalised patients hospitalized for febrile consciousness disorders in the infectious diseases department during our period. study.

Commented [a10]: Avoid repetition

Definition of variables:

Age: number of years lived by the person until the day of hospitalization. Patients were grouped by age group of 10 years range.

Commented [a11]: Only list and define the terms used in tis article. I've not come across bachelors for example yet appears here as a variable

Gender: permanent physical character~~istic~~ of a person making it possible to distinguish male and female individuals but also to determine the sex ratio.

Marital status :

Married : two people united by a matrimonial bond.

Bachelor :person who is not married.

Divorce :a person whose marriage has been legally dissolved.

Widower:person whose spouse has died.

Socio-professional layer:

Formal sector:set of official activities, recognized by the state with monthly remuneration.

Informal sector:all activities producing goods and services which escape the gaze or regulation of the state. (Driver, worker, merchant/trader, hairdresser, seamstress, farmer).

Housewife:woman who runs a house, takes care of the household.

Pupil/student:people who receive education in a pre-university school, university or professional establishment.

Unemployed:all people who do not have a job.

School level:it designates the highest level of education attained by the patient: Low: person whose level of education is limited to primary school.

Secondary: person whose level of education is between the end of primary school and the start of university.

Higher: person whose level of education is university.

Etiologies:

Cerebral toxoplasmosis:Opportunistic parasitic infection due to *Toxoplasma gondii* which occurs when the CD4 count is less than 200 cells. Diagnosis is made from medical imaging CT/MRI of the brain or PCR *Toxoplasma gondii* in the CSF / more or less PCR in the plasma or from clinical improvement by a 15-day antitoxoplasmic test treatment.

Tuberculosis:Gen-expert positivity in cerebrospinal fluid.

Gen-Expert: tuberculosis diagnostic tool, rapid, sensitive and specific for *Mycobacterium tuberculosis* with a sensitivity of 89% and a specificity of 91%. It is a real-time molecular biology test.

Cryptococcosis:opportunistic fungal infection appearing at the AIDS stage.

CRAG in CSF: qualitative and semi-quantitative detection of CSF antigens *Cryptococcus neoformans* in the cerebrospinal fluid (CSF)

Neuromalaria:Thick drop positive for *Plasmodium falciparum* + Coma

Bacterial meningoencephalitis:cloudy or purulent cerebrospinal fluid with or without identification of the bacterial germ responsible.

Viral meningoencephalitis: Clear CSF with or without isolation of the virus responsible by PCR

Brain abscess: focal suppuration of infectious origin developed within the brain parenchyma and appears regardless of the CD4 count. The diagnosis is made by brain imaging which highlights multiple foci of abscesses.

CMV encephalitis:occurs if CD4 count is less than 50 cells

Syphilitic encephalitis:non-specific neuropsychological disorders. Syphilitic serology and VDRL positivity in the cerebrospinal fluid allow the diagnosis to be made.

Lymphomas:headaches, vomiting, physical asthenia, mood changes are non-specific. Brain magnetic resonance imaging (MRI) and neurosurgical biopsy help confirm the diagnosis

HIV encephalitis:progressive deterioration of cognitive function in an HIV+ patient not on antiretroviral treatment or on antiretroviral treatment but with failure of cerebral virological control. MRI and cerebral lumbar puncture help eliminate other differential diagnoses

Factors associated with deaths were investigated using logistic regression in multivariate analysis and all variables with a value of $P < 0.05$ were considered statistically significant.

Data entry and analysis were carried out using EPI data software in version 3.1 and SPSS software in version 21, then processed by Microsoft Word and Excel 2013 software.

2-Results:

During the study period, we collected 205/534 HIV+ patients hospitalized for febrile consciousness disorders, representing a prevalence of 38.39%.

The average age of the patients was 42.28 ± 12.54 years [18-85 years] with a slight female predominance, i.e. a sex ratio of 0.97. The informal sector represented nearly 43% (88/205) of patients and nearly 2/3 (69.96) of patients had a low level of education and more than half (62.44%) of patients were married. . (Table 1)

Weight loss (66.34%) and cough (40%) were the most frequently found signs (Table 2).

Almost all (97.07%) of the patients had HIV type 1

More than 94% (180/205) of patients were in the severe immunosuppression stage (Table 3).

Only 6.82% (14/205) of patients were able to perform a brain scan and no magnetic resonance imaging was done (Table 4).

Commented [a12]: In this section, report on the results based on each objective and probably research question. That means the author first has to list the objectives and the research questions above and analyse if the study answered them or not

Cerebral Toxoplasmosis 73/205 (35.61%) and Neuromeningeal Cryptococcosis 45/205 (21.95%) were the most frequently found etiologies (Table 5).

Only 1/3 or 33.66% (69/205) were on antiretroviral treatment at the time of the study.

Death was reported in More than 5/6 (83.90%)(5/6) of our the patients died

The depth of coma, the absence of antiretroviral treatment and a CD4 count \leq 200 cells/mm3 were the factors associated with death in multivariate analysis. (Table 6).

3-Discussion:

This was a retrospective descriptive and analytical study on the etiologies and factors associated with the death of HIV-positive patients hospitalized for febrile consciousness disorders. Despite the retrospective and monocentric nature, this study allowed us to resulted in the identify-identification of the etiologies and factors associated with the death of HIV-positive patients hospitalized for febrile consciousness disorders in the infectious and tropical diseases department of the Donka university hospital center.

We collected 205/534 HIV+ patients hospitalized for febrile consciousness disorders, representing a prevalence of 38.39%.

This high prevalence could be explained by the fact that the majority of patients consult- seek health services in the phase of severe immunosuppression where opportunistic brain infections are very frequent.

The average age of our patients-participants was 42.28 ± 12.37 years.

We noted a slight female predominance, i.e. 50.73%.

This finding is similar to those of Franck et al [14] in Ethiopia in 2019 and Gunda et al [15] in Tanzania in 2016 with 62.1% and 61.85% respectively.

The informal sector was the most affected in this study.

This result was similar to that found by Ossibi I et al[3]who reported a prevalence of 40%.

This could be explained by the low level of education of this professional category who therefore had little information on HIV infection and its cost, little information on means of prevention and the benefits of screening- early and initiation of antiretroviral treatment.

We noted a higher proportion of married people 62.44%. The same observation was reported by Mbula et al [16] in 2020 in the Democratic Republic of Congo with 63%.

Commented [a13]: Elucidate this section based on the results above, you dont have to repeat the results in this section. Do good paragraphs explaining how each objective performed in your analysis

Commented [a14]: Adopt one system of referring to this hospital and use it all through the document soi t doesnt look like you are talking about different hospitals in different sections where it is referred to

Commented [a15]: This still does not explain why the prevalence is exceptionally owere in Morocco (11%) compared to the study site (38.39% and Benin, 38.1%. Does it mea nits only in Benin and Guinea where patients consult at advanced stage and in Morocco they consult uch earlier ?

Commented [a16]: Such statements must be combined with the subsequent statement that explains it soi t doesnt look like a stand alone statement that shold be under the results

Commented [a17]: Same to (a15) above

Commented [a18]: Same to (a15) above

Commented [a19]: What type of education ? It does sound that you have to get formal education to know how to prevent HIV, which may not be true unless backed by references published. Two, remember your study was not on HIV prevalence but on febrile consciousness disorders so focus here in your discussion

Weight loss 136/205 (66.34%) was the most frequently encountered symptom.

This result is similar to those of Zannou et al in Benin in 2004 [17].

Almost all (97.07%) of our patients were infected with HIV type 1. This is similar to those of Traoré et al [18] in 2014 in Mali.

The majority of our patients (94.24%) had a CD4 count below 200 cells/mm³. This result is higher than that of Karfo et al [19] in Burkina in 2018 who found only 78%.

This could be due to the fact that most of our patients consult at a late stage of HIV infection.

Toxoplasmosis (35.61%), neuromeningeal cryptococcosis (21.95%) and tuberculosis (21.47%) were the most frequently found etiologies.

These results are similar to those of Rolland OIB et al in 2016 [3] who also reported that cerebral toxoplasmosis, neuromeningeal cryptococcosis and neuromeningeal tuberculosis were the most common etiologies.

This result is consistent with current literature data which states that toxoplasmosis is the first opportunistic infection of the central nervous system during AIDS [5,20]

Nearly 2/3 of our patients were naive to antiretroviral treatment.

This result is different from that of Fall et al in Dakar in 2017 [21] who found that 91.6% of patients were on antiretroviral treatment.

This observation could be explained by the fact that the majority of our patients were unaware of their serological status and were therefore not on antiretroviral treatment.

Most of our patients (83.80%) had died.

This result is superimposable to those of Rolland et al [3] in 2016 with 83.2% fatality and lower than those of Karfo et al in Burkina in 2018 [19] who reported 44.1% deaths.

This observation could be explained by the fact that the patients were treated at a late stage of HIV infection.

The depth of the coma, the absence of antiretroviral treatment and severe immunosuppression were the factors associated with the death of PLHIV hospitalized for febrile consciousness disorders.

This result is similar to those of Chelli J et al in Tunisia, Mbondé AA et al in 2022 in Uganda, Traoré S in 2021 in Mali and Pang W et al in 2018 in China [6,10,22,23] and different from those of Rolland OIB et al [3] in 2016 in Brazzaville who reported that neuro-meningeal tuberculosis and anemia were significantly associated with death.

This result could be explained by the fact that HIV-positive patients not on antiretroviral treatment can rapidly progress towards severe immunosuppression (CD4 rate \leq 200 cells/mm³) and therefore become very susceptible to opportunistic brain infections with risk of brain disorders, consciousness and subsequent death.

Conclusion :

Febrile consciousness disorders in hospitalized HIV-positive patients remain frequent in the infectious and tropical diseases department of Donka University Hospital and toxoplasmosis was the most common etiology. The depth of the coma, severe immunosuppression and the absence of antiretroviral treatment were the factors associated with death.

Screening and early treatment of HIV infection would help reverse this trend.

Recommendations:

Improvement of the technical platform in order to be able to highlight all opportunistic infections, particularly viral ones, by carrying out molecular biology techniques in the cerebrospinal fluid.

Making brain imaging accessible and at a lower cost would improve guidance and even diagnostic confirmation and could therefore increase the survival rate through the implementation of documented treatment.

Warning (artificial intelligence):

No generative AI technologies such as large language models (ChatGPT, COPILOT, etc.) and text-to-image generators were used in the writing or editing of this manuscript.

Consent :

The free and informed consent of the participant or a trusted person ~~or including relatives~~ was sought and obtained before including the patient in the study.

Commented [a20]: Like stated earlier, adopt one name and use it throughout the document

Commented [a21]: Do this section based on the findings of this study only. For example, if the study found out that febrile consciousness disorders are more likely in those not on ART, your recommendation should be based on factors that would make ART accessible to all

Commented [a22]: This doesn't look authentic and should be excluded from the article. From what is documented here, only review was done on existing records and no patient was physically met

The anonymity and confidentiality of patients were respected.

Ethics approval

The study protocol was approved by the ethics committee of the Faculty of Health Sciences and Technologies of the Gamal Abdel Nasser University of Conakry.

Commented [a23]: Some countries have additional requirement for licensure beyond ethics review committee approval. Peradventure the author may check if licensing from National Ethics Committee for Health is also a requirement

UNDER PEER REVIEW

Tables :

Table 1: Distribution according to socio-demographic characteristics of the 205 HIV-positive patients hospitalized for febrile consciousness disorders from 2018 to 2023 in the infectious and tropical diseases department of Donka University Hospital, Guinea

Variables	Effective (n=205)	Percentage (%)
Age		
41-50 years old	63	30.73
Sex		
Women	104	50.73
Marital status		
Married	128	62.44
Others	77	37.56
Educational level		
Weak	143	69.76
Others	62	30.24
Occupation		
Informal sector	88	42.93
Others	117	57.07
Residence		
Conakry	121	59.02
Others	84	40.98

Average age: 42.28 (\pm 12.54) years with extremes of 18 and 85 years

Table 2: Distribution according to signs of the 205 HIV-positive patients hospitalized for febrile consciousness disorders from 2018 to 2023 in the infectious and tropical diseases department of Donka University Hospital, Guinea

Signs	Frequency	Percentage
Weight loss	136	66.34
Coma stage 1	128	62.44
Cough	82	40.00

Physical asthenia	79	38.54
Vomiting	73	35.61
Diarrhea	60	29.27
Coma stage 2	56	22.44
Hemiparesis	40	19.51
Psychomotor agitation	30	14.63
Coma stage 3	21	10.24
Chest pain	15	7.32
Dysphagia	15	7.32
Abdominal pain	8	3.9
Dyspnea	6	2.93

Table 3: Distribution according to the immuno-virological status of HIV-positive patients hospitalized for febrile consciousness disorders from 2018 to 2023 in the infectious and tropical diseases department of Donka University Hospital, Guinea

Immuno-virological assessment	Frequency	Percentage
Viral Load (n=45)		
Detectable	42	20.48

Undetectable		3	3.41
CD4 rate (n=191)			
Less than 200 cells per mm³	180		94.24
200-350 cells per mm ³		7	3.66
350-500 cells per mm ³		4	2.10

Table 4: Distribution according to diagnostic confirmation examinations of HIV-positive patients hospitalized for febrile consciousness disorders from 2018 to 2023 in the infectious and tropical diseases department of Donka University Hospital, Guinea

Diagnostic confirmation examinations	Frequency	Percentage
GE(DP) /TDR (n=204)		
Negative	191	93.63
Positive	13	6.37

CRAG(n=195)		
Negative	150	76.92
Positive	45	23.08
Gen-expert (n=199)		
Negative	151	77.44
Positive	44	22.56
Brain scan (14)		
Abnormal	11	78.57
Normal	3	21.43

Table 5: Distribution according to the main diagnosis of HIV-positive patients hospitalized for febrile consciousness disorders from 2018 to 2023 in the infectious and tropical diseases department of Donka University Hospital, Guinea

Main diagnosis:	Numbers (N=205)	Percentage
Toxoplasmosis	73	35.61
Neuromeningeal cryptococcosis	45	21.95
Tuberculosis	44	21.47
Bacterial meningoencephalitis	28	13.66
Severe malaria	13	6.34
None	2	0.97

Table 6: Distribution of HIV patients hospitalized for febrile consciousness disorders in the infectious and tropical diseases department of the Donka University Hospital of Guinea according to factors associated with death in multivariate analysis.

Associated factors	Odds Ratio	Confidence interval	P-value
Coma	6	1.77-17.49	0.003
CD4 rate			
[200-500 cells]	3.54	1.06-14.56	0.7
≤ 200 cells/mm ³	6,33	1.72-29.0	0.007
Anemia			
Yes/no	0.91	0.005-1.513	0.095
Creatininemia			
Low	0.273	0.023-3.291	0.306
High	14.15	1,478-5,495	0.072
ARV treatment			
Yes/no	0.02	1.74-19.17	0.004

References:

1. Quaiser S, Alam S, Khan R. Clinical profile and prognostic markers of acute febrile encephalopathy (AFE) in adult patients presenting to a North Indian tertiary care hospital. *International Journal of Nutrition, Pharmacology, Neurological Diseases*, 2015; 5:95-102
2. Key benchmarks on HIV/AIDS [online], 2023 [cited Oct 27, 2023]. Available at: <https://www.who.int/en/news-room/fact-sheets/detail/hiv-aids>
3. Rolland OIB, Obengui, Bébene DB, Macaire ONP, Herdan P, Claude BPJ et al. Neuromeningeal Disorders During HIV in the Infectious Diseases Department of Brazzaville University Hospital: Prevalence and Factors Associated with Death. *European Scientific Journal*, ESJ. 2016;12(33):177-177.
4. HAS.Moulinier. Central nervous system damage and HIV-1 infection, *Rev neurol (Paris)* 2006; 162(1), 22-42
5. HIV Clic: Main opportunistic conditions. [Online]. 2019 [cited 26Oct 2023]. Available at: <https://vihclic.fr/aides-consultation/infections-opportunistes/>
6. Chelli J, Bellazreg F, Aouem A, Hattab Z, Mesmia H, Lasfar NB, et al. Causes of death of HIV-infected patients in Central Tunisia. *The Pan African Medical Journal*, 2016;25(105) 12-19
7. Global HIV & AIDS Statistics — Fact sheet | UNAIDS [Online]. 2023 [cited Oct 26, 2023]. Available at: <https://www.unaids.org/en/resources/fact-sheet>
8. Umgelter A, Huber W, Schmid RM, Reindl W. Coma as a presenting symptom of primary HIV infection. *Sex Transm Infect.* 2007;83(2):85-6.
9. Obiako OR, Oparah S, Ogunniyi A. Causes of medical coma in adult patients at the University College Hospital, Ibadan Nigeria. *Niger Postgrad Med J.* 2011;18(1):1-7.
10. Mbonde AA, Mbatidde L, Demaerschalk BM, Aden AA, Zhang N, Butterfield R, et al. HIV Infection Is an Independent Predictor of Mortality Among Adults with Reduced Level of Consciousness in Uganda. *Am J Trop Med Hyg.* 2022;106(3):909-14.
11. Stepchenkova T, Martynenko O, Yurchenko O. Analysis of causes of death among HIV-infected patients of kyiv Regional AIDS Center during 2013. *J Int AIDS Soc.* 2014;17(4):633-639.
12. Adoukonou T, Dassi-Gbenou J, Glele RA, Dovonou A, Kpangon A, Houinato D. Epidemiology of neurological conditions in a population of people living with HIV in Parakou in 2014. *Revue Neurologique (Paris).* 2015;171:136.
13. El Fane M.

Central neurological manifestations in HIV-infected patients in the infectious diseases department of Casablanca University Hospital. *Bulletin of the Society of Exotic Pathology*. 2018; 111:24-30

14. Frank TD, Carter A, Jahagirdar D, Biehl MH, Douwes-Schultz D, Larson SL, et al. Global, regional, and national incidence, prevalence, and mortality of HIV, 1980–2017, and forecasts to 2030, for 195 countries and territories: a systematic analysis for the Global Burden of Diseases, Injuries, and Risk Factors Study 2017. *The Lancet HIV*. 2019;6(12):831-59.

15. Gunda DW, Godfrey KG, Kilonzo SB, Mpondo BC. Cytopenias among ART-naive patients with advanced HIV disease on enrollment to care and treatment services at a tertiary hospital in Tanzania: A cross-sectional study. *Malawi Med J*. 2017;29(1):43-52.

16. Mbula MMK, Situakibanza HNT, Mananga LG, Longokolo MM, Mandina NM, Mayasi NN, et al. Clinical and biological profile of people living with HIV/AIDS followed in the Infectious Diseases Department of the University Clinics of Kinshasa (Democratic Republic of Congo). *Malian Journal of Infectiology and Microbiology*. 2020;15(2):21-9.

17. Zannou DM, Kindé-Gazard D, Vigan J, Adè G, Sèhonou JJ, Atadokpèdé F, et al. Clinical and immunological profile of HIV-infected patients screened in Cotonou, Benin. *Medicine and Infectious Diseases*. 2004;34(5):225-8.

18. Traoré AM, Minta DK, Fomba M, Cissé H, Diallo K, Coulibaly I, et al. Epidemiological and evolutionary profile of HIV-positive patients referred to the Point G University Hospital, Bamako, Mali. *Bull Soc Pathol Exot*. 2014;107(1):22-6;

19. Karfo R, Kabré E, Coulibaly L, Diatto G, Sakandé J, Sangaré L. Evolution of biochemical and hematological parameters in people living with HIV/AIDS under antiretroviral treatment at the General Camp Aboubacar Sangoule Lamizana Medical Center. *Pan Afr Med J*. 2018;29:159.

20. Cissoko I, Seydi M, Deguenonvo F. Current profile of cerebral toxoplasmosis in hospitals in Dakar. *Med Trop* 2013; 23:197-201.

21. Fall S, Djiba B, Diagne N, Ndiaye M, Dieng F, Kyelem EMC, et al. Profile of hematological manifestations of HIV infection in adults: Single-center study of 262 Senegalese observations. *African Journal of Internal Medicine*. 2017 ;4(1-2):33-6.

22. Traore S. Risk factors associated with deaths of people living with HIV at antiretroviral treatment sites in Bamako, Mali. *Mali Public Health*. 2021; 1(2) 60-6.

23. Pang W, Shang P, Li Q, Xu J, Bi L, Zhong J, et al. Prevalence of Opportunistic Infections and Causes of Death among Hospitalized HIV-Infected Patients in Sichuan, China. *Tohoku J Exp Med*. 2018; 244(3):231-242.