

Case report

A Deadly Cerulean Limb- Phlegmasia Cerulea Dolens complicated with Pulmonary Embolism

ABSTRACT

Aims: Asses the presentation of phlegmasia cerulea dolens and differentiate venous etiologies of ischemic limb with arterial thrombus

Presentation of case: A case reported in a 51 years-old woman with underlying metastatic endometrial carcinoma, presented with sudden onset of pain and swelling of left leg. Initial bedside doppler show patent arterial perfusion over the affected leg. It was later developed into bluish discoloration and bedside doppler show poor perfusion of arteries. Subsequently, she succumbed to illness less than 24 hours from first presentation due to suspected pulmonary embolism.

Discussion and conclusion: Patient with increased risk of developing deep vein thrombosis (DVT) such as those with underlying malignancy has shown to develop a more extreme case of the disease. Phlegmasia cerulea dolens (PCD) is a rare manifestation of DVT over lower extremities that cause critical limb ischemia. Incidence of PCD is unknown however, it is estimated from a case series about 10-20% of patient underwent thrombolysis for extensive DVT, developed into phlegmasia. Common risk factors that contribute to developing PCD are malignancy, surgery and trauma.

This rare condition may be responsible for mortality up 20-40%, mainly due pulmonary embolism. Medical therapy involves administration of anticoagulant such as IV Heparin during acute setting should be initiated as soon as possible, follow by oral factor 10a inhibitors ie; rivaroxaban up to 6 months. Surgical therapy such as catheter-directed thrombolytic however remain a mainstay therapy.

Keywords: Phlegmasia cerulea dolens, Deep vein thrombosis, Limb ischemia, malignancy

1. INTRODUCTION

Phlegmasia cerulea dolens (PCD) is an extreme form of deep venous thrombosis (DVT). It is characterized by an acute history of excruciating pain, profound limb swelling, and skin discoloration[1]. DVT is a common venous thromboembolic (VTE) disorder, reporting an incidence rate of 1.6 per 1000 annually [2], and PCD is a consequence of massive acute venous thrombosis. The incidence is also more common in females than in males and is reported to be higher in the fifth and sixth decades of life[4]. PCD was first described by Gregoire in 1938 to differentiate massive venous thrombosis causing ischemic changes from phlegmasia alba dolens, which characterized venous thrombosis without ischemia [4].

In PCD, it is manifested with ischemia-related extensive venous thrombosis [1]. As venous flow becomes obstructed, venous hypertension increases interstitial oedema and extensive fluid sequestration into the affected limb. High compartment pressure eventually causes the arterial system to collapse, leading to acute ischemia and venous gangrene [3]. Left-sided lower extremities are more common with a ratio of 3:1 or 4:1 [4]. This may be explained by an anatomical

variant in which the right common iliac artery overrides and compresses the left common iliac vein against the lumbar spine, causing thrombosis in the left common iliac vein [5].

Factors that contributed to the development of DVT and subsequently PCD included malignancy (20–40%) [4], with higher case of thrombosis reported in gynaecology malignancy-related patient [6]. Other less common causes are advanced age, with more prominent cases reported in the fifth and sixth decades of life [4]. Use of contraception, post-surgery, and trauma are among other factors. Administration of anticoagulant [7,8] and surgical intervention such as catheter-directed thrombolytic remain a mainstay therapy [8]. The condition, if left untreated, leads to amputation of the gangrenous limb and death, mainly due to the development of pulmonary embolism. In this study, we reported a PCD case of a 51-year-old woman with underlying pelvic malignancy who presented with a one-day history of left lower limb swelling and pain, mimicking acute limb ischemia.

2. PRESENTATION OF CASE

A 51-year-old woman with underlying metastatic cervical adenocarcinoma and receiving palliative chemotherapy presented with a sudden onset of pain and swelling of the left lower limb. She was referred to the surgical team by an emergency physician for acute limb ischemia. Upon review in the emergency department, she appeared breathless and in pain. Her left lower limb was swollen with bluish discoloration, and she was unable to move the toes and ankle. The left distal pulses were well-felt up to the femoral artery. This raised a suspicion of venous ischemia instead of arterial ischemia. The ultrasound doppler revealed a long segment of thrombus from the left common femoral vein up to the external iliac vein. The electrocardiogram showed an S-wave in lead I, a Q-wave, and an inverted T-wave in lead III. Her latest computed tomography (CT) of the thorax and abdomen, which was done one month ago, showed a huge pelvic mass with multiple lung metastases. She was treated with extensive DVT, resulting in PCD and pulmonary embolism secondary to an advanced cervical adenocarcinoma. She deteriorated fast to the point where she became more breathless, and the bluish limb extended to the contralateral lower limb was observed. She succumbed to death within 24 hours of presentation, despite the prompt initiation of anticoagulant therapy.



FIG 1. METASTATIC CERVICAL ADENOCARCINOMA

3. DISCUSSION

PCD is a limb [9] and life-threatening condition [1]. Venous obstruction and intravascular hypertension cause sequestration of fluid, increasing the compartmental pressure. The arterial system will eventually collapse, thus, resulting in tissue ischemia and limb gangrene. Other differential diagnoses, such as acute cellulitis, peripheral artery disease, and acute arterial occlusion might be considered.

Pulmonary embolism, which contributes up to 10%–15% of hospital deaths with relation to the occurrence of DVT [1], is the biggest factor in mortality in PCD as well.

Ultrasound is the main imaging modality that facilitates the diagnosis of PCD. The inability to compress the vein during ultrasonography is pathognomonic for a thrombus in the vein [9]. Catheter venography and arteriography were mentioned as the gold standard in one case reported for their ability to be therapeutic [9]. However, a case reported in a healthy male

patient who presented with a sudden onset of leg swelling shows CT-Angio is superior as it can detect any pelvic mass causing venous compression [10].




Fig 2. CT-Angio report

Early and fast recognition are essential for survival outcomes, as immediate intervention can be initiated. Systemic anticoagulant combined with catheter-directed deep vein thrombolysis has been the treatment for PCD. Fluid resuscitation is essential to replace losses of fluid sequestration in the lower limb [11].

A clinical challenge may arise to differentiate limb ischemia caused by arterial or venous occlusion. PCD of a limb is distinguished by its extensive swelling, out-of-proportion pain, and skin discoloration, while in patients with ischemic limb causes by arterial occlusion, they presented with pulselessness followed by pain, limb paralysis, parasthesia, pallor, and poikilothermia. Venous occlusion caused by thrombosis is detected through ultrasound, while contrast enhanced CT is a recommended imaging method to locate multiple thrombosis that are often seen in patients with peripheral artery disease. However, challenging as it may appear to differentiate both conditions, a patient with a known case of vascular disease indicates the ischemic limb is caused by an arterial thrombus, such as a patient with previous cardiovascular disease who specifies an atherosclerotic or stent- or graft related thrombus. Trauma precipitates an arterial thrombus, and patients with aortic dissection may indicate a similar arterial origin [12]. Nevertheless, both conditions benefit from immediate intervention with both medical and surgical interventions such as intravenous heparin and thromboembolectomy, respectively.

A case reported in a patient with iliac vein compression and deep vein thrombosis showed that open venous thrombectomy promotes fast limb decongestion and contributes to good morbidity and overall mortality for the patient [11]. However, it is worth noting the apparent disadvantages of thrombectomy, such as recurrent thrombosis [11], and the need to perform the procedure under general anaesthesia. In this case, the immediate initiation of anticoagulants is sufficient to prevent venous propagation.

As previously reported, mortality in venous thrombosis is mostly caused by the development of pulmonary embolism. The prognosis remains poor in this case due to the fact that the patient may have already developed massive pulmonary embolism prior to the arrival of the emergency department.

4. CONCLUSION

Though rare and deadly, a high clinical suspicion and prompt diagnosis of PCD are crucial for patients with advanced pelvic malignancy presented with an ischemic limb.

ETHICAL APPROVAL

Consent was obtained by the patient in this study

REFERENCES

1. Chaochankit W, Akaraborworn O. Phlegmasia Cerulea Dolens with Compartment Syndrome. *Ann Vasc Dis.* 2018 Sep 25;11(3):355-357. Accessed 30 June 2024. Available: 10.3400/avd.cr.18-00030. PMID: 30402189; PMCID: PMC6200621.
2. Waheed SM, Kudravalli P, Hotwagner DT. Deep Vein Thrombosis. [Updated 2023 Jan 19]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Accessed 30 June 2024. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK507708/>
3. Ibrahim B, Kattimani R. Phlegmasia Cerulea Dolens, a Deadly Complication of Deep Vein Thrombosis: Case Report and Review of Literature. *Cureus.* 2021 Nov 26;13(11):e19927. Accessed 30 June 2024. Available: 10.7759/cureus.19927. PMID: 34966616; PMCID: PMC8710803.
4. Abootalebi A, Salemi M, Heidari F, Azimi Meibody A. Phlegmasia Cerulea Dolens: A Rare Case Report. *Adv Biomed Res.* 2020 Jul 27;9:31. Accessed 30 June 2024. Available: 10.4103/abr.abr_3_20. PMID: 33072643; PMCID: PMC7532826.
5. Mangla A, Hamad H. May-Thurner Syndrome. [Updated 2022 Nov 30]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Accessed 30 June 2024. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK554377>
6. Pookcharoen, N., Insin, P., & Asavapiriyanon, S. (2018). Risk Factor of Deep Vein Thrombosis in Gynecologic Cancer Patients at Rajavithi Hospital. *Asian Pacific Journal of Cancer Care*, 3(1), 5. Accessed 30 June 2024. Available: <https://doi.org/10.31557/apjcc.2018.3.1.5>
7. Said A, Sahlieh A, Sayed L. A comparative analysis of the efficacy and safety of therapeutic interventions in phlegmasia cerulea dolens. *Phlebology.* 2021 Jun;36(5):392-400. Accessed 30 June 2024. Available: 10.1177/0268355520975581. Epub 2020 Nov 25. PMID: 33236674.
8. Saydam O, Serefli D. Outcomes of rheolytic thrombectomy in phlegmasia cerulea dolens. *Vascular.* 2021 Apr;29(2):280-289. Accessed 30 June 2024. Available: 10.1177/1708538120943320. Epub 2020 Jul 26. PMID: 32715971.
9. Chaochankit W, Akaraborworn O. Phlegmasia Cerulea Dolens with Compartment Syndrome. *Ann Vasc Dis.* 2018 Sep 25;11(3):355-357. Accessed 30 June 2024. Available: 10.3400/avd.cr.18-00030. PMID: 30402189; PMCID: PMC6200621.
10. Said A, Sahlieh A, Sayed L. A comparative analysis of the efficacy and safety of therapeutic interventions in phlegmasia cerulea dolens. *Phlebology.* 2021 Jun;36(5):392-400. Accessed 30 June 2024. Available: 10.1177/0268355520975581. Epub 2020 Nov 25. PMID: 33236674.
11. Mwita, JC; Mwandri, MB1; Cox, M2. Phlegmasia cerulea dolens in a long distance driver. *Nigerian Journal of Clinical Practice* 19(2):p 290-292, Mar-Apr 2016. Accessed 30 June 2024. Available: 10.4103/1119-3077.175966
12. Obara H, Matsubara K, Kitagawa Y. Acute Limb Ischemia. *Ann Vasc Dis.* 2018 Dec 25;11(4):443-448. Accessed 30 June 2024. Available: 10.3400/avd.ra.18-00074. PMID: 30636997; PMCID: PMC6326052.