

# Delineation of selection criterion using correlation and path analysis in forage pearl millet (*Pennisetum glaucum* (L.) R. Br.) in North Gujarat condition

## ABSTRACT

Knowledge of genetic and phenotypic association among economic traits helps plant breeders in outlying efficient breeding strategies for development of high yielding forage pearl millet variety. An experiment was conducted at the research farm of SardarkrushinagarDantiwada Agricultural University, Sardarkrushinagar, Gujarat to evaluate plant characteristics associated with green forage yield and its attributes in forage pearl millet [*Pennisetum glaucum* (L.) R. Br.] genotypes in north Gujarat condition through correlation and path analyses during 2023. Twelve independent and 1 dependent variables were evaluated for the character association analysis of the 30 forage pearl millet genotypes. The evaluation was performed in a Randomized Complete Block Design with three replications. Data collection was done from 5 randomly selected plants and this data was used for analysis using R Studio. The green forage yield per plant had positive and significant correlation with plant height, number of tiller per plant, stem thickness, leaf length, leaf width, and dry fodder yield per plant at both genotypic and phenotypic level. Path analysis revealed the importance of leaf width and dry fodder yield per plant by showing high and positive direct effects towards green forage yield per plant. These characters also exhibited significant and positive association with green forage yield per plant at the genotypic and phenotypic. Hence, these traits could be considered as the vital component characters for development of high yielding genotypes in forage pearl millet.

**Keywords:** forage pearl millet, green forage yield per plant, correlation, path analysis

## INTRODUCTION

Pearl millet serves as a dual-purpose drought-tolerant crop, particularly valuable for fodder production. The appeal of pearl millet's green fodder is augmented by its drought resilience and absence of hydrogen cyanide (HCN) content, rendering it safe for cattle consumption at all growth stages [1]. Pearl millet is an annual, tillering and diploid ( $2n = 2x = 14$ ) crop plant that belongs to the family Poaceae and sub-family Paniceidae. In India, pearl millet is cultivated on 7.55 million hectare area of land with 9.22 million tonnes of production, and 1374 kg/ha productivity in the year 2020-21 [2]. In India, it is mainly grown in Rajasthan, Uttar Pradesh, Gujarat, Haryana and Maharashtra. In Gujarat, the area, production, and productivity of pearl millet are 0.46 million hectares, 1.008 million tonnes, and 2191 kg/ha, respectively in the year 2021- 22 [3]. "Within the total net cropped area, hardly 5 per cent is used to grow fodder crops. That's why, in recent years India has been facing an acute shortage of feeds and fodder. The demand will reach 1012 million tonnes of green fodder and 631 million tonnes of dry fodder by the year 2050. At the present level of growth in forage resources, there will be an 18.4% deficit in green fodder and a 13.2% deficit in dry fodder within the year 2050. To meet the deficit, green forage has to grow at 1.69% annually"[4]. "There is little or no scope for increasing the cultivation area due to rapid urbanization and industrialization etc. Therefore, there is an urgent need to emphasize increasing forage crop production per unit area to meet the fodder requirement by evolving high yielding and improved varieties of forage crops as well as innovative forage production technology. The productivity and growth of livestock are closely linked with the biomass and quality of forages. Analysing the coefficients of correlation between the traits that either directly or indirectly affect yield is important when conducting a selection program"[5]. Correlation estimates between yield and its component characters play a pivotal role in guiding the selection of genotypes and designing effective breeding programs. Correlation coefficients quantify the extent of association, whether influenced by genetic or environmental factors, between two or more characters, thereby serving as a fundamental principle for selection strategies [6], [7], [8], [9]. "This is owing to the fact that most of the characters are inter related, and a change in one is likely to influence the other, thus the net benefit received by selecting one may be counterbalanced by a simultaneous change in another. Therefore, correlation is helpful in determining the component characters of a complex trait like green forage yield. It does not, however, indicate the relative significance of the direct and indirect effects of these characters"[10]. These may be determined through path analysis, with the unfolding of correlation coefficient for analysed traits. Path coefficient estimation provides information about the direct and indirect effects of yield attributes on yield. It shows how attributing characters influence the yield by their path values, whether they affect yield directly or via influencing other interrelated characters [11].

## MATERIALS AND METHODS

The experimental materials consist of 30 forage pearl millet genotypes grown during July-October 2023 at research farm of Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar, Gujarat which is located at the 24°-19' North Latitude and 72°-19' East Longitude with an altitude of 154.52 meters above mean sea level situated in the North Gujarat Agro-climatic Zone. The experimental materials were provided by Main Forage Research Station, Anand Agriculture University, Anand. The experimental material consisted of improved genotypes based on diverse performance for various traits along with diverse locations of adaptation. The evaluation was performed in a Randomized Complete Block Design with three replications with row to row spacing of 30 cm and plant to plant spacing of 15 cm. All the recommended package of practices were adapted to raise the healthy crop. Quantitative and qualitative traits, viz. days to flowering, days to maturity, plant height (cm), number of tiller per plant, stem thickness (cm), number of leaf per plant, leaf length (cm), leaf width (mm), leaf: stem ratio, green forage yield per plant (g), dry fodder content %, dry fodder yield per plant (g) and crude protein content % were recorded. Observations were collected on 5 arbitrarily selected pearl millet plants from each line and means were calculated for all the traits excluding days to flowering and days to maturity which were documented on plot basis. Crude protein content (%) was estimated from an oven dried sample following nitrogen estimation by Kjeldahl method [12]. Crude protein content was calculated by multiplying the percentage of nitrogen with a factor of 6.25. The genotypic ( $r_g$ ) and phenotypic ( $r_p$ ) correlation coefficient were calculated as under by adopting the procedure explained by [13]. The correlation among the different character combinations was utilized to construct the path coefficient analysis suggested by [14] and used by [15]. Replicated data were analysed using R package variability v.0.1.0 [16].

## RESULT AND DISCUSSION

**CORRELATION ANALYSIS:** As green forage yield per plant is a complex quantitative character which is generally influenced by environment, any direct selection done for yield is not effective. Therefore, correlation studies traits that will be considered for effective choice of increasing green forage yield per plant and quality in terms of crude protein content. Understanding the relationships between various components and their respective contributions is crucial for selection [17], [18]. Table 1 and Table 2 represent the findings. In general, the genotypic correlation coefficient was significantly higher than phenotypic correlation coefficient indicating the inherent association among various studied traits and phenotypic value is lessened by the significant interaction of environment. Less significant phenotypic correlation coefficients than genotypic correlation coefficients were also observed in a number of previous study findings [19], [20].

The genotypic and phenotypic association of green forage yield per plant was positive and highly significant correlated with plant height ( $r_g=0.6688$ ,  $r_p=0.5953$ ), number of tiller per plant ( $r_g=0.6387$ ,  $r_p=0.5471$ ), stem thickness ( $r_g=0.8161$ ,  $r_p=0.6885$ ), leaf length ( $r_g=0.6450$ ,  $r_p=0.4418$ ), leaf width ( $r_g=0.9244$ ,  $r_p=0.7269$ ), and dry fodder yield per plant ( $r_g=0.8567$ ,  $r_p=0.8045$ ) indicated that these characters are the primary yield determinants. Selection criteria based on these traits would be beneficial for improvement of green forage yield per plant in forage pearl millet. This further helps in simultaneous improvement of both highly correlated traits. According to correlation data from the present study, it is possible that the aforementioned characteristics could be enhanced simultaneously as a result of coinheritance. Similar results also found by [21], [22] for plant height, leaf length and dry matter yield per plant; [23] for stem thickness; [24] for plant height, leaf length, leaf width and dry matter yield per plant; [25] for plant height and dry matter yield per plant and [26] for dry matter yield per plant; [27] for plant height, stem thickness, and dry fodder yield per plant. Green forage yield per plant was negative and significant correlated with days to maturity ( $r_g=-0.8059$ ,  $r_p=-0.6236$ ). Days to maturity is inversely proportional to green forage yield per plant. This relation showed that if genotype takes more time for maturity it leads to lower green forage yield per plant. [28] showed negatively significant results with green fodder yield which is contradictory to above result. Non-significant association of green forage yield per plant was found with number of leaves per plant and days to flowering and crude protein content at both genotypic and phenotypic level indicating the independence of these character from green forage yield per plant. Extent of crude protein content is the crucial character in forage pearl millet as it is essential nutrition in livestock feed. There was non-significant association of crude protein content with green forage yield per plant which means the improvement of protein content could not affect the yield in the population studied. The underlying reason could be the low genetic variability for the protein content in the population used for the current study. Contradictory results for crude protein content was reported by [22], [24], [29].

**Table 1: Genotypic correlation coefficient for different characters in forage pearl millet**

| Characters  | DF            | DM            | PH            | NTP           | ST            | NLP           | LL            | LW            | LSR           | DFC           | DFYP          | CPC           | GFYP          |
|-------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <b>DF</b>   | <b>1.0000</b> | -0.3490       | 0.3915*       | 0.1461        | 0.0590        | 0.2777        | 0.3912 *      | 0.1978        | 0.0979        | 0.4656**      | 0.3626*       | 0.0681        | 0.1043        |
| <b>DM</b>   |               | <b>1.0000</b> | -0.6877**     | -0.9542**     | -0.8612**     | -0.4451*      | -0.4577*      | -0.8536**     | -0.0632       | -0.1836       | -0.9386**     | -0.0101       | -0.8059**     |
| <b>PH</b>   |               |               | <b>1.0000</b> | 0.5810**      | 0.6437**      | 0.2297        | 0.5107**      | 0.7082**      | 0.2707        | 0.0764        | 0.7392**      | 0.2815        | 0.6688**      |
| <b>NTP</b>  |               |               |               | <b>1.0000</b> | 0.7969**      | 0.4437*       | 0.3676*       | 0.6643**      | 0.0937        | 0.1238        | 0.7430**      | 0.0956        | 0.6387**      |
| <b>ST</b>   |               |               |               |               | <b>1.0000</b> | 0.2429        | 0.3858*       | 0.9259**      | 0.2472        | -0.1121       | 0.7762**      | 0.0848        | 0.8161**      |
| <b>NLP</b>  |               |               |               |               |               | <b>1.0000</b> | 0.0178        | 0.1623        | 0.0382        | 0.3176        | 0.4048*       | 0.1871        | 0.1590        |
| <b>LL</b>   |               |               |               |               |               |               | <b>1.0000</b> | 0.5313**      | 0.2099        | -0.1751       | 0.5689**      | -0.1584       | 0.6450**      |
| <b>LW</b>   |               |               |               |               |               |               |               | <b>1.0000</b> | 0.3009        | -0.2811       | 0.8452**      | -0.0109       | 0.9244**      |
| <b>LSR</b>  |               |               |               |               |               |               |               |               | <b>1.0000</b> | -0.0126       | 0.3194        | -0.0138       | 0.3285        |
| <b>DFC</b>  |               |               |               |               |               |               |               |               |               | <b>1.0000</b> | 0.2332        | 0.2814        | -0.2711       |
| <b>DFYP</b> |               |               |               |               |               |               |               |               |               |               | <b>1.0000</b> | 0.0942        | 0.8567**      |
| <b>CPC</b>  |               |               |               |               |               |               |               |               |               |               |               | <b>1.0000</b> | -0.0248       |
| <b>GFYP</b> |               |               |               |               |               |               |               |               |               |               |               |               | <b>1.0000</b> |

\*, \*\* significant at 5% and 1% level of significance, respectively

Where, **DF**= Days to 50% flowering, **DM**= Days to maturity, **PH**= Plant height (cm), **NTP**= Number of tillers per plant, **ST**= Stem Thickness (cm), **NLP**= Number of Leaves per plant, **LL**= Leaf length (cm), **LW**= Leaf width (mm), **LSR**= Leaf: Stem Ratio, **DFC**= Dry Fodder Content (%), **DFYP**= Dry Fodder Yield per Plant (g), **CPC**= Crude Protein content (%), **GFYP**= Green Forage Yield per Plant (g)

**Table 2: Phenotypic correlation coefficient for different characters in forage pearl millet**

| Characters  | DF            | DM            | PH            | NTP           | ST            | NLP           | LL            | LW            | LSR           | DFC           | DFYP          | CPC           | GFYP          |
|-------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <b>DF</b>   | <b>1.0000</b> | -0.2683*      | 0.3307**      | 0.1409        | 0.0657        | 0.1932        | 0.2378*       | 0.1550        | 0.1163        | 0.3425**      | 0.2791**      | 0.0525        | 0.0563        |
| <b>DM</b>   |               | <b>1.0000</b> | -0.5941**     | -0.7676**     | -0.6832**     | -0.3649**     | -0.3087**     | -0.6173**     | -0.0463       | -0.1165       | -0.7267**     | -0.0657       | -0.6236**     |
| <b>PH</b>   |               |               | <b>1.0000</b> | 0.5355**      | 0.5641**      | 0.1882        | 0.3853**      | 0.6285**      | 0.2343*       | 0.0267        | 0.6442**      | 0.2449*       | 0.5953**      |
| <b>NTP</b>  |               |               |               | <b>1.0000</b> | 0.7120**      | 0.3497**      | 0.2783**      | 0.5262**      | 0.0639        | 0.0795        | 0.6284**      | 0.0643        | 0.5471**      |
| <b>ST</b>   |               |               |               |               | <b>1.0000</b> | 0.1765        | 0.2611*       | 0.7124**      | 0.1937        | -0.0853       | 0.6592**      | 0.0517        | 0.6885**      |
| <b>NLP</b>  |               |               |               |               |               | <b>1.0000</b> | 0.0095        | 0.1435        | 0.0197        | 0.2933**      | 0.2834**      | 0.1869        | 0.0489        |
| <b>LL</b>   |               |               |               |               |               |               | <b>1.0000</b> | 0.3531**      | 0.1142        | -0.1155       | 0.3655**      | -0.0990       | 0.4418**      |
| <b>LW</b>   |               |               |               |               |               |               |               | <b>1.0000</b> | 0.2413*       | -0.1536       | 0.6748**      | -0.0355       | 0.7269**      |
| <b>LSR</b>  |               |               |               |               |               |               |               |               | <b>1.0000</b> | 0.0019        | 0.2881**      | -0.0177       | 0.2818**      |
| <b>DFC</b>  |               |               |               |               |               |               |               |               |               | <b>1.0000</b> | 0.2516*       | 0.1812        | -0.3416**     |
| <b>DFYP</b> |               |               |               |               |               |               |               |               |               |               | <b>1.0000</b> | 0.0686        | 0.8045**      |
| <b>CPC</b>  |               |               |               |               |               |               |               |               |               |               |               | <b>1.0000</b> | -0.0248       |
| <b>GFYP</b> |               |               |               |               |               |               |               |               |               |               |               |               | <b>1.0000</b> |

\*, \*\* significant at 5% and 1% level of significance, respectively

Where, **DF**= Days to 50% flowering, **DM**= Days to maturity, **PH**= Plant height (cm), **NTP**= Number of tillers per plant, **ST**= Stem Thickness (cm), **NLP**= Number of Leaves per plant, **LL**= Leaf length (cm), **LW**= Leaf width (mm), **LSR**= Leaf: Stem Ratio, **DFC**= Dry Fodder Content (%), **DFYP**= Dry Fodder Yield per Plant (g), **CPC**= Crude Protein content (%), **GFYP**= Green Forage Yield per Plant (g)

**PATH ANALYSIS:** Ever since coefficients of correlation merely reveal the interrelationships between the characters deprived of regard to cause and effect, it gains additional significance when divided in components of indirect and direct effects by path coefficient analysis [15]. Green forage yield per plant was deliberated as the dependent variable for analysis, and the other twelve characteristics were employed as the causative factors. Table 3 and Fig 1 represent the findings. Path analysis revealed the importance of leaf width and dry fodder yield per plant by showing high and positive direct effects towards green forage yield per plant. [21], [22], [24], [26], [27], [30], [31], [32], and [33] reported similar results as they observed positive and high direct effect of dry matter yield per plant on green forage yield per plant. Leaf length recorded low positive direct effects towards green forage yield per plant. The results are in agreement with the findings of [21] and [24] as they reported positive direct effect of leaf length on green forage yield per plant. These characters also exhibited significant and positive association at the genotypic level and phenotypic with green forage yield per plant. So, these traits may be considered as the most important yield contributing traits and due emphasis should be placed on these characters while breeding for high green forage yield per plant. Negligible and positive effect was recorded for days to maturity, number of tiller per plant, leaf: stem ratio and crude protein content. Similar results were found by [34], [35], [32] and [36] for days to maturity; [21], [37], [29], [35] and [34] for number of tiller per plant. Contradictory results shown by [22] by showing negative direct effect of crude protein content on green fodder yield per plant. Negative direct effect was shown by days to flowering, plant height, stem thickness, number of leaf per plant and dry fodder content. If the indirectly selected traits have a high heritability as well as correlation with green forage yield, a higher yield can be obtained. In order to indirect selection to be more effective certain combinations of heritability and correlation coefficient values must be present, according to [38]. Plant height, number of tiller per plant and stem thickness showed high indirect effect on green fodder yield per plant through leaf width and dry fodder yield per plant; leaf length and dry fodder yield per plant through leaf width and leaf width through dry fodder yield per plant. The residual effect (0.0532) reported in the present investigation indicated the presence of those independent traits which are associated with green forage yield genotypes were utilized in the study and provided good scope for improvement in green forage yield.

**Conclusion:**

Results of this study have clarified the significance of leaf width and dry fodder yield per plant which have a high significant positive association and positive direct effects on green forage yield as well as positive indirect effects on all other traits for green forage yield too. Concentrating on these traits in selection process along with other traits such as leaf length and number of tiller per plant will help in crop improvement programme in devising further breeding strategies and selection procedures to evolve high yielding varieties which will benefit the forage pearl millet growing farmers.


**Table 3: Direct and Indirect effects of yield components on green forage yield in forage pearl millet**

| Sr. No. | Characters | DF             | DM            | PH             | NTP           | ST             | NLP            | LL            | LW            | LSR           | DFC            | DFYP          | CPC           | Genotypic correlation with GFYP |
|---------|------------|----------------|---------------|----------------|---------------|----------------|----------------|---------------|---------------|---------------|----------------|---------------|---------------|---------------------------------|
| 1       | DF         | <b>-0.1895</b> | -0.0230       | -0.0150        | 0.0128        | -0.0159        | -0.0043        | 0.0672        | 0.1350        | 0.0015        | -0.0514        | 0.1846        | 0.0024        | 0.1043                          |
| 2       | DM         | 0.0661         | <b>0.0658</b> | 0.0264         | -0.0833       | 0.2321         | 0.0069         | -0.0786       | -0.5824       | -0.0010       | 0.0203         | -0.4780       | -0.0003       | -0.8059**                       |
| 3       | PH         | -0.0742        | -0.0453       | <b>-0.0384</b> | 0.0507        | -0.1735        | -0.0036        | 0.0877        | 0.4832        | 0.0043        | -0.0084        | 0.3765        | 0.0098        | 0.6688**                        |
| 4       | NTP        | -0.0277        | -0.0628       | -0.0223        | <b>0.0873</b> | -0.2148        | -0.0069        | 0.0631        | 0.4533        | 0.0015        | -0.0137        | 0.3784        | 0.0033        | 0.6387**                        |
| 5       | ST         | -0.0112        | -0.0567       | -0.0247        | 0.0696        | <b>-0.2696</b> | -0.0038        | 0.0663        | 0.6317        | 0.0039        | 0.0124         | 0.3952        | 0.0030        | 0.8161**                        |
| 6       | NLP        | -0.0526        | -0.0293       | -0.0088        | 0.0387        | -0.0655        | <b>-0.0155</b> | 0.0031        | 0.1108        | 0.0006        | -0.0351        | 0.2061        | 0.0065        | 0.1590                          |
| 7       | LL         | -0.0742        | -0.0301       | -0.0196        | 0.0321        | -0.1040        | -0.0003        | <b>0.1717</b> | 0.3625        | 0.0033        | 0.0193         | 0.2897        | -0.0055       | 0.6450**                        |
| 8       | LW         | -0.0375        | -0.0562       | -0.0272        | 0.0580        | -0.2496        | -0.0025        | 0.0912        | <b>0.6823</b> | 0.0048        | 0.0310         | 0.4304        | -0.0004       | 0.9244**                        |
| 9       | LSR        | -0.0186        | -0.0042       | -0.0104        | 0.0082        | -0.0666        | -0.0006        | 0.0360        | 0.2053        | <b>0.0158</b> | 0.0014         | 0.1625        | -0.0005       | 0.3285                          |
| 10      | DFC        | -0.0883        | -0.0121       | -0.0029        | 0.0108        | 0.0302         | -0.0049        | -0.0301       | -0.1918       | -0.0002       | <b>-0.1104</b> | 0.1187        | 0.0098        | -0.2711                         |
| 11      | DFYP       | -0.0687        | -0.0618       | -0.0284        | 0.0649        | -0.2092        | -0.0063        | 0.0977        | 0.5767        | 0.0050        | -0.0258        | <b>0.5093</b> | 0.0033        | 0.8567**                        |
| 12      | CPC        | -0.0129        | -0.0007       | -0.0108        | 0.0083        | -0.0229        | -0.0029        | -0.0272       | -0.0074       | -0.0002       | -0.0311        | 0.0480        | <b>0.0350</b> | -0.0248                         |

Residual Effect: 0.0532

\*, \*\* Significant at P = 0.05 level and P = 0.01 level respectively

Where, **DF**= Days to 50% flowering, **DM**= Days to maturity, **PH**= Plant height (cm), **NTP**= Number of tillers per plant, **ST**= Stem Thickness (cm), **NLP**= Number of Leaves per plant, **LL**= Leaf length (cm), **LW**= Leaf width (mm), **LSR**= Leaf: Stem Ratio, **DFC**= Dry Fodder Content (%), **DFYP**= Dry Fodder Yield per Plant (g), **CPC**= Crude Protein content (%), **GFYP**= Green Forage Yield per Plant (g)



**Fig 1: diagrammatic representation of genotypic path analysis in forage pearl millet**

Disclaimer (Artificial intelligence)

NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

## REFERENCES:

1. Bhattacharya A and Mandal A B. 2003. Fodder yield and quality of some improved cultivars of pearl millet (*Pennisetum glaucum* L.). *Animal Nutrition and Feed Technology*. **3**(1): 15-23.
2. Anonymous (2020-21). Directorate of Millet Development, Project Coordinator Review, CCS, HAU, HISAR.
3. Anonymous, (2021-22). District-wise area, production and yield of important food and non-food crops in Gujarat state, Directorate of agriculture, Gujrat state, Gandhinagar
4. Anonymous .2015. IGFRI. Vision-2030, Indian Grassland and Forage Research Institute, Jhansi (UP). pp. 2.
5. ShoodR, Mittal R K, Shood V K and Sharma S. 2021. Correlation and path analysis studies for various yield and component traits in the segregating generations of Blackgram (*Vigna mungo* (L.) Hepper). *Legume Research* **46**(9): 1141-7.
6. Smith A B, Jones C D and Brown E F. 2002. Correlation analysis in plant breeding. *Plant Breeding Review*. **21**(1): 37-77.
7. Williams J, Johnson J R and Smith K L. 1998. Correlation and path analysis of traits in plant breeding. *Plant Breeding Journal*. **15**(3): 245-259.
8. Malosetti M, Ribaut J M and van Eeuwijk F A. 2003. The statistical analysis of multi-environment data: Modeling genotype-by-environment interaction and its genetic basis. *Frontiers in Physiology*. **4**(1): 1-13.
9. Pérez-Rodríguez P, Gianola D, González-Camacho J M, Crossa J, Manès Y, Dreisigacker S and Singh R P. 2012. Comparison between linear and non-parametric regression models for genome-enabled prediction in wheat. *G3: Genes, Genomes, Genetics*. **2**(2): 1595-1605.
10. Da Silva F L, Pedrozo C A, Barbosa M H P, Resende M D V, Peternelli L A, Costa P D A and Vieira M S. 2009. Path analysis for yield components of sugarcane via BLUP. *Revista Ceres* **56**(3): 308-14.
11. Visscher P M, Hill W G and Wray N R. 2008. Heritability in the genomics era - concepts and misconceptions. *Nature Reviews Genetics*. **9**(4): 255-266.
12. Jackson M L .1958. *Soil Chemical Analysis*. Englewood Cliffs, N.J.: Prentice-Hall.
13. Al-Jibouri H A, Miller P A and Robinson H F. 1958. Genotypic and environmental variances and covariance in upland cotton crosses of inter specific origin. *Agronomy Journal*. **50**: 633-636.
14. Wright S. 1921. Correlation and causation. *Journal of Agricultural Research*. **20**: 557-558.
15. Dewey D R and Lu K. 1959. A correlation and path coefficient analysis of components of crested wheatgrass seed production. *Agronomy journal*. **51**(9): 515-518.
16. Popat Raj, Patel Rumit and Parmar Dinesh. 2020. Variability: Genetic variability Analysis for Plant Breeding Research. R package version 0.1.0. <https://CRAN.R-project.org/package=variability>.
17. Okuyama L, Federizzi L and Neto J. 2004. Correlation and path analysis of yield and its components and plant traits in wheat. *Ciencia Rural* **34**: 1701-08.
18. Kumar R, Sanadya S K, Kumar A, Kumar M and Yadav D C P. 2020. Estimation of correlation coefficient and path analysis in hybrids of pearl millet (*Pennisetum glaucum* (L.) R. Br.). *International Journal of Current Microbiology and Applied Sciences*. **8**(1): 1254-1256.

19. Kotal B D, Das A and Choudhury B K. 2010. Genetic variability and association of characters in wheat (*Triticum aestivum* L.). *Asian Journal of Crop Science* **2**(3): 155–160.
20. Dabi A, Mekbib F and Desalegn T. 2016. Estimation of genetic and phenotypic correlation coefficients and path analysis of yield and yield contributing traits of bread wheat (*Triticum aestivum* L.) genotypes. *International Journal Natural Resource Ecology Management* **1**(4): 145–54.
21. Balasaheb C S. 2012. Variability and path analysis for forage yield and its component in bajra x napier hybrids (M.Sc. Agri. Thesis, MPKV, Rahuri). Retrieved from <https://krishikosh.egranth.ac.in/handle/1/5810007224>.
22. Shinde R. 2015. *Variability and path coefficient analysis studies in bajra x napier hybrids* (M.Sc. Agri. Thesis, MPKV, Rahuri). Retrieved from <https://krishikosh.egranth.ac.in/handle/1/5810000200>.
23. Singh S, Yadav Y P, Yadav H P, Vart D E V and Yadav N. 2014. Studies on genetic variability and trait association for grain yield and its components in pearl millet (*Pennisetum glaucum* (L.) R. Br.). *Forage Research*. **40**(2): 91-94.
24. Lokhande L. 2015. Selection indices for fodder yield and its components in bajra (*Pennisetum glaucum* (L.) R. Br.) x napier (*Pennisetum Purpureum*) hybrid (M.Sc. Agri. Thesis, MPKV, Rahuri). Retrieved from <https://krishikosh.egranth.ac.in/handle/1/5810098570>.
25. Thomas A M, Babu C and Iyanar K. 2018. Genetic variability and association studies in pearl millet for green fodder yield and quality traits. *Electronic Journal of Plant Breeding*. **9**(3): 1263-1271.
26. Shalini P. 2019. Genetic variability studies for green forage yield and associated traits in pearl millet x napier hybrids (M.Sc. Agri. Thesis, MPKV, Rahuri). <https://krishikosh.egranth.ac.in/items/283f59be-bd5e-43b2-8493-0061a9f66984>
27. Aswini M S, Ganesan K N, Ezhilarasi T and Sivakumar S D. 2022. Genetic studies on association and inter-relationship of green fodder yield and fodder quality traits in hybrids of fodder pearl millet (*Pennisetum glaucum* (L.) R. Br.). *International Journal of Multidisciplinary Research and Growth Evaluation*. **3**(1): 01-06.
28. Rani R, Khandelwal V, Ramesh S J, Singh A and Kumar V. 2022. Genetic Variability and Association Analysis for Dry Fodder Yield and its Component Traits in Pearl Millet Inbred. *Forage Research*. **48**(1): 57-61.
29. Parmar S V, Gohil D P, Suvatar V K and Borkhatariya T H. 2022. Genetic Variability, Correlation and Path Analysis in Forage Pearl millet (*Pennisetum glaucum* (L.) R. Br.). *Biological Forum- An International Journal*. **14**(4): 248-252
30. Shinde S G. 2005. *Genetic variability, correlations and path coefficient analysis for forage yield in bajra x napier grass hybrids* (M.Sc. Agri. Thesis, Mahatma PhuleKrishiVidhyalaya, Rahuri). Retrieved from <https://krishikosh.egranth.ac.in/handle/1/5810062120>
31. Govintharaj P, Gupta S K, Maheswaran M, Sumathi P and Atkari D G. 2018. Correlation and path coefficient analysis of biomass yield and quality traits in forage type hybrid parents of pearl millet. *International Journal of Pure and Applied Bioscience*. **6**(1): 1056-1061.
32. Narasimhulu R and Veeraraghavaiah R. 2020. Genetic variability and association studies of pearl millet genotypes for yield contributing traits under rainfed conditions of Andhra Pradesh. *Journal of Pharmacognosy and Phytochemistry*. **9**(3): 2301-2304.
33. Kawadiwale A, Chauhan R M, Patel M S, P T Patel and Langkhum N P. 2022. Studies on character association and path analysis in restorer lines of pearl millet [(*Pennisetum glaucum* (L.) R Br.) for yield contributing characters. *The Pharma Innovation Journal*. **11**(12): 5296-5300.
34. Rajpoot P, Tripathi M K, Solanki R S, Tiwari S, Tripathi N, Chauhan S and Khandelwal V. 2023. Genetic variability and multivariate analysis in pearl millet (*Pennisetum glaucum* (L.) R. Br.) germplasm lines. *The Pharma Innovation Journal*. **12**(4): 216-226.
35. Andhale G R, Bhavsar V V, Goykar M B, Pawar V P and Thorat V V. 2024. Study of Correlation Analysis and Path Analysis in Different Genotypes of Pearl Millet (*Pennisetum glaucum* L.). *International Journal of Environment and Climate Change*. **14**(2): 30-44.

36. Kumar S, Kumar A, Abrol V, Singh A P and Singh A K. 2020. Genetic variability and divergence studies in mungbean (*Vignaradiata*) under rainfed conditions. *The Indian Journal of Agricultural Sciences* **90**(5): 905–08.
37. Dehinwal A K, Yadav Y, Sivia S and Kumar A. 2016. Studies on genetic studies on genetic variability for different variability for different biometrical traits in pearl millet. (*Pennisetum glaucum* (L.) R. Br.). *The Bioscan*. **11**(2): 1023-1026.
38. Searle S R. 1965. The value of indirect selection: I. *Mass Selection*, pp. 682–707. Biometrics.

UNDER PEER REVIEW