

First report of *Carpophilus dimidiatus* (Fabricius, 1792) (Coleoptera: Nitidulidae: Coleoptera) in stored groundnut and field maize from West Bengal

Comment [o1]: Order comes first, then Family

Abstract

Carpophilus genus belongs to the family nitidulidae under the order coleoptera. Majority of the members under this genus are either sap feeder or fungus feeder. A small number of these, however, is economically categorised as stored grain pests. The commonly known species under this genus are *C. dimidiatus*, *C. hemipterus*, *C. humeralis*, *C. marginellus*, *C. mutilatus*, *C. obsoletus*, *C. davidsoni*, *C. lugubris* etc. Among these, *Carpophilus dimidiatus* (Fabricius) has been reported worldwide as a pest causing significant post-harvest loss. In India, different cereals like bajra, rice products, sago flour as well as oil seeds have been found to be infested by this pest. This pest was reported earlier from our country. But the information on identification as well as the name of the crop(s) infested by the pest were lacking. It was recorded with proper identification in stored groundnut from Bangalore and Chennai for the first time from our country in the year 2022. This report is going to be the first ever report of *Carpophilus dimidiatus* on groundnut in storage and on maize in field from West Bengal.

Keywords: ~~First report, *Carpophilus dimidiatus*, groundnut, nitidulid beetle, groundnut, maize,~~
nitidulid beetle, West Bengal

Comment [o2]: I consider this is not a keyword

Formatted: Strikethrough

Comment [o3]: I sorted the keywords alphabetically

UNDER PEER REVIEW

1. Introduction

Nitidulids are the most diverse group of beetles distributed widely in all zoogeographical regions of the world. This family is represented by 350 genera with more than 4500 species [1]. They show tremendous diversity in their feeding habitat. Many species are associated with decaying materials, some are sap feeders found on sap-flows or wound of trees [2], some are pollinators[3], whereas many species are recorded as stored product insects [4]. These beetles also display great diversity in their morphological adaptations and general body form. As majority of the Nitidulids are saprophagous and mycetophagous, they receive less importance and are thought to be without any economic importance. However, a small group of nitidulids is regarded as economically important, particularly the genus *Carpophilus*, infesting different stored products like grain product, cracked rough rice, wheat bran, nuts and nut products, spoiled commodities and vegetable products including the dry fruits. Some of these are *Carpophilus dimidiatus* (Fabricius), *C. hemipterus* (Linnaeus), *C. humeralis* (Fabricius), *C. marginellus* (Motschulsky), *C. mutilatus* (Erichson) and *C. obsoletus* (Erichson), *C. davidsoni* (Dobson), *C. lugubris* (Murray) etc. Among these, *Carpophilus dimidiatus* (Fabricius) has been reported from several countries, throughout the globe, to cause significant post-harvest loss [5].

2. History of Nomenclature

Carpophilus dimidiatus was described originally as *Nitidula dimidiata* by Fabricius in the year 1792 [6]. Afterwards, this species strived a long path to arrive at *Carpophilus dimidiatus*. *Nitidula dimidiata* (Fabricius) was changed to *Carpophilus auripilosus* Wollaston, *Carpophilus lewisi* Reitter, *Carpophilus pusillus* Stephens, *Carpophilus vittiger* Murray, *Carpophilus biguttatus* Gemminger and Harold and finally *Carpophilus dimidiatus*, Grouvelle [7]. The reason behind this was the close similarity between the species. Several misidentification throughout the globe had happened earlier among the several species of *Carpophilus*.

3. Background of this work

This beetle was found to infest corn fields around Uttar Banga Krishi Viswavidyalaya, Pundibari (26° 31' N, 89° 06' E) region (Block: Coochbehar II) from 2020 onwards at irregular basis. Afterwards the infestation was decreased. Again the infestation was started from 2022 onwards. This time the infestation was not restricted only within Pundibari region, but also in the other areas under Coochbehar – II block. Add to this, the pest was also found to infest groundnut in storage. The stored groundnut samples infested by this pest were also found in the several parts of different districts under Sub-Himalayan region of West Bengal; i.e. Alipurduar, Darjeeling and Coochbehar. Afterwards, a study was initiated to record the presence of *C. dimidiatus* in the said region covering different areas.

Comment [o4]: Please put the country of the area of this finding

Formatted: Font: Italic

Formatted: Font: Italic

4. Materials and methods:

Surveys were conducted in the different districts mentioned above. Groundnut samples of 500g were collected from each of the grocery shops. Collected samples from storage were kept separately (within glass jars) in the laboratory. After a certain period, small beetles were found to emerge from the said samples.

Under field condition, the infestation of the said pest was spotted only on maize. Cob samples with specific damage symptoms were collected from fields of Coochbehar district and put into plastic zip covers and were taken to the laboratory for further observation.

All the images and photographs were captured by Magnus TZM6 stereozoom microscope and Samsung Galaxy S21 FE handset. The specimens were stored at the laboratory of Department of Agril. Entomology, Uttar Banga Krishi Viswavidyalaya (26° 31' N, 89° 06' E), Pundibari, Coochbehar (West Bengal) for further identification. Regarding taxonomic identification, keys of different scientists were utilised [5, 7, 8]. Identification of this beetle was confirmed on the basis of molecular characterisation, too.

5. Results and Discussion

5.1 Survey report

Table 1 and Table 2 describes the occurrence of *C. dimidiatus* from both storage and field condition –

Table 1. Survey report on *C. dimidiatus* infesting stored groundnut from Terai region of West Bengal

Name of District	Location	GPS Code	Damage (%)*
Coochbehar	South Khagrabari	26°33'N89°44'E	65-70%
	Malgudam Road	26°31'N89°45'E	
Darjeeling	Pradhan Nagar, Siliguri	26°72'N88°41'E	20-25%
	Bidhan Nagar, Siliguri	26°71'N88°42'E	
Alipurduar	Sobhaganj	26°49'N89°52'E	55-60%
	Samuktala Road	26°48'N89°52'E	

* Damage (%) was calculated on the basis of how many seeds were damaged out of 100 after keeping it in laboratory for three months

Table 2. Survey report on *C. dimidiatus* infesting Maize under field condition in Coochbehar

Name of District	Location	GPS Code	Cob infestation (%)
Coochbehar	Kaminirghat	26°33'N89°43'E	25-30%
	Tengnamari	26°34'N89°42'E	
	Platoon Bridge	26°36'N89°37'E	
	Singimari Paschimpar	26°46'N89°36'E	

* Cob infestation (%) was calculated on the basis of how many cobs were infested out of 100

5.2 Identification of the pest

The identification of this pest was done with the help of both taxonomic and molecular characterisation.

Formatted: Font: Not Italic

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

5.2.1 Taxonomic identification

The key taxonomic identifying characters of this beetle are as follows –

- i) Dorsum dark red-brown to black
- ii) Each elytron contains large dull-orange yellow spot.
- iii) Elytra short. It exposes two abdominal tergites and pygidium thereby. The tergite VII is visible dorsally, whereas the VIIIth one is visible from ventral surface. In case of male, this VIIIth tergite is button like.
- iv) Males are smaller than the females.
- v) Cuticle is clothed with light yellow hairs.
- vi) Punctures on pronotum are very deep.
- vii) Labrum is somehow visible and free.




Fig. 1. Images of *Carophilus dimidiatus* under stereozoom microscope

5.2.2 Molecular characterisation

The taxonomic identification of this pest was confirmed through molecular characterisation. Fragment of mitochondrial 16S rRNA (ribosomal RNA) gene was amplified and then sequencing process was completed. The submission of the said sequences to National Center for Biotechnology Information (NCBI) database resulted in generation of GenBank accession number. The sequence showed more than 99% match with species recorded or reported elsewhere and already submitted in NCBI database. The submission of amplified 16S rRNA resulted in generation of GenBank accession number **OP346785**.

5.2.3 Damage Symptom

In maize field, it was observed that a good number of cobs remained partially opened (due to some unknown reason). That cobs were found harbouring adult *C. dimidiatus* beetle. The adults were found to damage the seeds which, ultimately, turned brown and dried. The partial opening of the immature cob has been reported as a common symptom from several maize growers in this region.

Further investigation is needed to ascertain whether *C. dimidiatus* is the sole causal agent for this immature cob opening or some other factors (like bird attack) are also present. But this was a confirmed observation that *C. dimidiatus* was found to be present in each and every partially opened cob.

In storage, both the grubs and adults damage the groundnut seeds. These bore the seeds and the bore holes are prominently visible. In case of excessive damage, only shells are left and internal content of the seed is converted into powdery mass.



Fig. 2. Damage caused by *Carpophilus dimidiatus*

6. Summary and Conclusion

Naveena *et al.* (2022) [9] reported this pest in stored groundnut (from Bangalore and Chennai) for the first time from our country with proper identification. In the said article, the author clearly mentioned that the earlier authors like Sengupta *et al.* (1984) [10] and Dasgupta *et al.* (2013) [11] from West Bengal had reported *C. dimidiatus*. But there was lack of specific information regarding the grain infested by the pest as well as taxonomic key of the pest. That is why, the present report of *Carpophilus dimidiatus* in both field (corn) and store (groundnut) is going to be the first report from West Bengal with taxonomic and molecular identification.

References

1. Jelinek J, Carlton C, Cline AR, Leschen RAB. Nitidulidae. In: Handbook of Zoology, Volume 2. De Gruyter, Berlin, Germany. 2011: 390–407.
2. Martin JEH. Collecting, preparing and preserving insects, mites and spiders. In: The insects and arachnids of Canada (Part 1). Agriculture Canada Publication. 1977; 1643:182.
3. Nagulapalli SL, Kunal G, Rakesh Das R, Jha, S. Insect pollinators and their abundance in West Bengal. Indian J. Entomol. 2020; 82 (4): 849-853. DoI: 10.5958/0974-8172.2020.00110.8.
4. Bousquet Y, Bouchard P, Davies A, Sikes DS. Checklist of beetles (Coleoptera) of Canada and Alaska. Second edition. Zookeys. 2013; 360: 1-44. DoI: 10.3897/zookeys.360.4742
5. Connell WA. Hosts of *Carpophilus dimidiatus*. Journal of Economic Entomology. 1975; 68(2): 279–280.
6. Fabricius JC. Entomologia systematica emendata et aucta. Secundum classes, ordines, genera, species adjectis synonymis [sic], locis, observationibus, descriptionibus. Tom. I. [Pars I]. - pp. I-XX [= 1-20], 1-330. Hafniae. (Proft).

Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Formatted: Font: Bold

7. Grouvelle AH. Byturidae, Nitidulidae pars 56. In: Junk W, Schenkling S (Eds.) *Coloeopterorum Catalogus*. Springer, Dordrecht. 1913; 1-223.
8. Reales N, Rocamundi N, Marvaldi AE, Gorgolas MCF, Stadler T. Morphological and molecular identification of *Carpophilus dimidiatus* (Coleoptera: Nitidulidae) associated with stored walnut in Northwestern Argentina. *Journal of Stored Products Research*. 2018; 76: 37-42.
9. Naveena NL, Shashank PR, Aidbhavi R. Occurrence of *Carpophilus dimidiatus* Fabricius on Stored Groundnut from India. *Proc. Entomol. Soc. Wash.* 2022; 124(1): 184-186. doi: 10.4289/0013-8797.124.1.184.
10. Sengupta T, P. Mukhopadhyay P, Sengupta R. Major beetle pests of stored food products in India. *Zoological Survey of India*. 1984; 65.
11. Dasgupta J, Pal TK, Hegde VD. On the occurrence of *Carpophilus maculatus* Murray from Kolkata, India (Coleoptera: Nitidulidae). *Entomon*. 2013; 38(1): 39-46.

Formatted: Font: (Default) Times New Roman

UNDER PEER REVIEW