

# BER PRODUCTION IN ARID ZONES: CHALLENGES, MANAGEMENT AND OPPORTUNITIES FOR THE FUTURE

## ABSTRACT

Ber (*Ziziphus mauritiana*) is a drought-resistant fruit tree that shows promise for arid and semi-arid regions due to its ability to thrive in low-water conditions and poor soils. This review explores the challenges and opportunities associated with ber production in arid zones. Key challenges include water scarcity, soil degradation, and limited agricultural infrastructure. Conversely, opportunities exist in advancing irrigation techniques, utilizing drought-tolerant varieties, and improving soil management practices. Recent innovations in sustainable agriculture, such as rainwater harvesting and soil moisture conservation, could enhance ber cultivation efficiency. Additionally, increased research into genetic improvement and pest management could lead to higher yields and better-quality fruits. The integration of modern technologies with traditional practices holds potential for expanding ber production, offering economic and environmental benefits to arid regions. Addressing these challenges while leveraging opportunities could significantly improve food security and livelihoods in these areas.

**Keywords:** Ber, *Ziziphus mauritiana*, arid zones, drought resistance, irrigation, soil management, sustainable agriculture, genetic improvement etc.

## 1. INTRODUCTION

The cultivation of Ber (*Ziziphus mauritiana*), commonly known as Indian jujube, has gained considerable attention in recent years, particularly in arid and semi-arid regions. This resilient fruit crop, native to South Asia, is celebrated for its adaptability to harsh climatic conditions, making it an ideal candidate for cultivation in regions where water scarcity, poor soil fertility, and extreme temperatures limit the productivity of other crops. As climate change continues to exacerbate these environmental challenges, the expansion of Ber cultivation presents a promising solution for sustaining agricultural productivity, enhancing food security, and supporting rural livelihoods in arid zones (Ahmed *et al.* 2020).

Arid regions, characterized by low and erratic rainfall, high temperatures, and limited vegetation, pose significant challenges for agricultural production. Traditional crops often struggle

to survive under such conditions, leading to low yields and increased vulnerability to food insecurity. However, Ber's inherent drought tolerance, low water requirement, and ability to thrive in nutrient-poor soils make it a valuable crop for these regions. Its deep root system allows it to access water from deeper soil layers, while its hardy nature enables it to withstand high temperatures and poor soil conditions. Moreover, Ber trees are known for their long productive life, providing a reliable source of income for farmers over many years. Despite its potential, Ber cultivation in arid zones is not without challenges (Alexander, 1990). The primary obstacle is the lack of awareness and knowledge among farmers about the best agronomic practices for Ber cultivation. This includes understanding the optimal planting time, irrigation management, pest and disease control, and post-harvest handling. Additionally, the lack of access to quality planting material and inputs, such as fertilizers and pesticides, further hampers the productivity of Ber orchards. Furthermore, the marketing of Ber fruits poses another challenge. In many arid regions, the absence of well-established market channels and infrastructure limits farmers' ability to sell their produce at competitive prices, reducing their overall profitability (Allen *et al.* 2005).

Addressing these challenges requires a multi-faceted approach that involves research, extension services, and policy support. Research institutions need to focus on developing improved Ber cultivars that are not only high-yielding but also resistant to pests and diseases common in arid zones. Extension services play a crucial role in disseminating knowledge about improved cultivation practices and providing technical support to farmers (Arumugam *et al.* 2021). Additionally, governments and policymakers must create an enabling environment by investing in infrastructure, such as roads and storage facilities, and by facilitating access to credit and markets for Ber producers. On the other hand, the opportunities for expanding Ber production in arid zones are significant. The increasing demand for Ber fruits, both in local and international markets, presents a lucrative opportunity for farmers. Ber fruits are not only consumed fresh but are also processed into various products such as dried fruits, candies, and beverages, adding value to the crop and opening up new revenue streams. Furthermore, the cultivation of Ber can contribute to environmental sustainability by preventing desertification and soil erosion, as the trees help in stabilizing the soil and improving its organic matter content (Aslam *et al.* 2006).

The development of agroforestry systems that integrate Ber trees with other crops or livestock can further enhance the sustainability and productivity of farming systems in arid zones. Such systems can improve soil fertility, increase biodiversity, and provide additional sources of

income for farmers. Moreover, the cultivation of Ber in arid regions can help in the conservation of water resources, as the crop requires significantly less water compared to other fruit crops, thus contributing to the efficient use of limited water resources in these areas (Athawale *et al.* 2020). Ber production in arid zones presents both challenges and opportunities for the future. While the crop's adaptability to harsh conditions makes it a promising option for improving food security and livelihoods in arid regions, overcoming the challenges related to cultivation practices, market access, and policy support is crucial for realizing its full potential. With concerted efforts from researchers, extension agents, and policymakers, Ber cultivation can become a key component of sustainable agricultural systems in arid zones, offering a viable solution to the pressing issues of climate change, desertification, and rural poverty (Ayars *et al.* 2006).

## 2. Ecophysiological Adaptations of Ber to Arid Environments

Ber (*Ziziphus mauritiana*), known for its resilience in challenging climates, exhibits a range of ecophysiological adaptations that make it exceptionally suited for cultivation in arid environments. These adaptations are crucial for its survival and productivity under conditions of extreme water scarcity, high temperatures, and poor soil fertility, common characteristics of arid zones. One of the most notable adaptations of Ber is its deep root system. The tree's roots can penetrate deep into the soil, allowing it to access water reserves far below the surface. This deep-rooting ability is particularly advantageous in arid environments where surface water is limited and sporadic. Additionally, Ber roots have a high level of hydraulic conductivity, enabling efficient water uptake and transport even under drought conditions. Another key adaptation is the tree's ability to minimize water loss through transpiration. Ber leaves are small, thick, and covered with a waxy cuticle, which reduces the surface area exposed to sunlight and limits water loss. The stomata, or pores on the leaf surface, are also strategically adapted to close during the hottest parts of the day, further conserving water. This stomatal regulation helps Ber maintain a positive water balance, even during prolonged dry periods (Azhane Ahmad *et al.* 2019).

Ber also exhibits remarkable tolerance to high temperatures. The tree's leaves contain heat shock proteins and other protective compounds that stabilize cellular structures and enzymes, preventing heat-induced damage. Additionally, the plant's photosynthetic machinery is adapted to function efficiently under high light intensities and temperatures, ensuring sustained growth and productivity even during the peak of summer. In terms of nutrient acquisition, Ber trees are highly efficient in utilizing the scarce nutrients available in arid soils. They have a symbiotic relationship

with mycorrhizal fungi, which enhance nutrient uptake, particularly phosphorus, a limiting nutrient in many arid soils. This symbiosis allows Ber to thrive in nutrient-poor conditions where other crops would fail. Moreover, Ber's ability to produce fruit under these harsh conditions is supported by its flexible reproductive strategies. The tree can flower and fruit multiple times a year, ensuring that even if one reproductive cycle fails due to adverse conditions, others may succeed (Balendres *et al.* 2021). This phenological plasticity is a critical survival mechanism in unpredictable arid environments. Overall, the ecophysiological adaptations of Ber underscore its potential as a key crop for arid regions. By harnessing these natural traits, Ber production can be optimized to provide sustainable livelihoods in areas where traditional agriculture struggles to succeed, thereby addressing both environmental and socioeconomic challenges in arid zones (Beltran, 1999).

### 3. Current Status of Ber Cultivation in Global Arid Zones

Ber (*Ziziphus mauritiana*) cultivation has become increasingly significant in arid and semi-arid regions globally due to its resilience and adaptability to harsh environmental conditions. In countries such as India, Pakistan, and China, Ber has long been an essential crop, providing food, fodder, and economic stability to farming communities in water-scarce regions. India, in particular, is a leading producer of Ber, with extensive cultivation in the arid regions of Rajasthan, Gujarat, and Maharashtra. The crop has also gained traction in other parts of Asia and Africa, including countries like Thailand, Sudan, and Kenya, where it is valued for its drought resistance and minimal water requirements (Beltran *et al.* 2006). In the Middle East, particularly in the arid regions of Saudi Arabia and the United Arab Emirates, Ber is cultivated on a smaller scale, often as part of agroforestry systems that aim to combat desertification. The cultivation practices in these regions are primarily traditional, with limited adoption of modern agronomic techniques, leading to variable yields and quality. Despite its potential, the global expansion of Ber cultivation is constrained by challenges such as limited access to improved cultivars, inadequate irrigation infrastructure, and lack of awareness among farmers about best practices. Overall, while Ber cultivation is well-established in some parts of the world, its full potential in global arid zones remains untapped. Addressing these challenges through research, extension services, and policy support could significantly enhance Ber production and contribute to sustainable agriculture in these vulnerable regions (Ben-Gal *et al.* 2004).

### 4. Drought Tolerance Mechanisms in Ber: A Comparative Analysis

Ber (*Ziziphus mauritiana*) is a remarkable fruit crop known for its ability to thrive in arid and semi-arid regions, where water scarcity poses significant challenges to agriculture. The plant's resilience to drought is attributed to several physiological and morphological adaptations that enable it to survive and produce under harsh conditions. A key drought tolerance mechanism in Ber is its deep root system, which allows the plant to access water from deeper soil layers, well beyond the reach of many other crops. This extensive root network also helps in stabilizing the soil and preventing erosion, contributing to environmental sustainability in arid zones (Ben-Gal *et al.* 2006). Another critical adaptation is Ber's ability to regulate stomatal closure, thereby reducing water loss through transpiration. This physiological response is particularly important during prolonged dry spells, as it minimizes water expenditure while maintaining essential metabolic processes. Additionally, Ber exhibits osmotic adjustment by accumulating solutes such as proline and sugars, which helps the plant retain water and maintain cell turgor under drought stress. Comparatively, Ber's drought tolerance mechanisms are more effective than those of many other fruit crops, making it an ideal candidate for cultivation in water-limited environments. The plant's ability to withstand high temperatures and poor soil conditions further enhances its suitability for arid regions. Understanding and harnessing these drought tolerance mechanisms are crucial for optimizing Ber production, improving food security, and supporting sustainable agriculture in areas increasingly affected by climate change and desertification (Benzioni, 1995).

## 5. Agronomic Practices for Optimizing Ber Yield in Arid Regions

Optimizing Ber (*Ziziphus mauritiana*) yield in arid regions requires a strategic approach to agronomic practices tailored to the unique challenges of these environments. Given Ber's resilience to drought and poor soils, it is essential to enhance these inherent traits through careful management.

**Site Selection and Soil Preparation:** The selection of well-drained, sandy loam soils with a pH of 6.5 to 8.5 is crucial for successful Ber cultivation. Deep plowing and the incorporation of organic matter improve soil structure, water retention, and nutrient availability, fostering healthier tree growth. Additionally, raised bed planting can enhance root aeration and reduce soil salinity, a common issue in arid zones (Benzioni *et al.* 1992).

**Water Management:** Efficient irrigation is vital in arid regions. Drip irrigation is the preferred method, delivering water directly to the root zone and minimizing evaporation losses. Scheduling irrigation based on critical growth stages—such as flowering and fruiting—ensures optimal water

use efficiency. Mulching with organic materials, such as straw or leaves, helps conserve soil moisture and regulate soil temperature, further supporting tree growth in water-scarce conditions (Benzioni et al. 1999).

**Nutrient Management:** The application of balanced fertilizers, particularly nitrogen, phosphorus, and potassium, is essential for maximizing Ber yield. Regular soil testing guides precise fertilizer application, preventing nutrient deficiencies or excesses. Additionally, the incorporation of organic fertilizers, such as compost or farmyard manure, improves soil fertility and microbial activity, enhancing nutrient uptake by the trees (Bernstein, 1975).

**Pruning and Canopy Management:** Regular pruning helps maintain an open canopy structure, facilitating light penetration and air circulation. This practice not only enhances fruit quality but also reduces the incidence of pests and diseases, which can be exacerbated in dense, poorly ventilated canopies. By implementing these agronomic practices, farmers can significantly improve Ber productivity in arid regions, contributing to sustainable agricultural systems and resilient livelihoods in these challenging environments (Bonfil *et al.* 2007).

## 6. Soil Management Strategies for Ber Orchards in Nutrient Poor Soils

The cultivation of Ber (*Ziziphus mauritiana*) in arid zones often faces the challenge of nutrient-poor soils, which can significantly impact tree health and fruit yield. Effective soil management strategies are crucial for optimizing the growth and productivity of Ber orchards in these challenging environments.

**Organic Matter Incorporation:** One of the primary strategies for improving soil fertility in nutrient-poor soils is the incorporation of organic matter. Adding compost, farmyard manure, or green manure can enhance soil structure, increase water retention, and boost microbial activity. These organic amendments provide essential nutrients, such as nitrogen, phosphorus, and potassium, which are critical for the healthy growth of Ber trees (Brown, 2004).

**Use of Mulching:** Mulching with organic materials, such as straw, leaves, or grass clippings, helps conserve soil moisture, reduce soil erosion, and maintain a stable soil temperature. Mulching also contributes to the gradual decomposition of organic material, releasing nutrients over time and improving soil fertility (Cai *et al.* 2001).

**Soil pH Management:** Maintaining an optimal soil pH is essential for nutrient availability to Ber trees. In arid regions, soils tend to be alkaline, which can limit the availability of certain nutrients.

The application of soil amendments like gypsum or sulfur can help lower the soil pH, making nutrients more accessible to the plants (Carr *et al.* 2004).

**Microbial Inoculants and Biofertilizers:** The use of microbial inoculants, such as nitrogen-fixing bacteria and mycorrhizal fungi, can enhance nutrient uptake in Ber trees. These beneficial microbes form symbiotic relationships with the tree roots, increasing the availability of essential nutrients like phosphorus and nitrogen (Chaoiun *et al.* 2021).

**Balanced Fertilization:** Applying balanced fertilizers based on soil tests is essential to address specific nutrient deficiencies in Ber orchards. This targeted approach ensures that trees receive the right amounts of macro- and micronutrients, promoting vigorous growth and high fruit yields. Implementing these soil management strategies can significantly improve the productivity of Ber orchards in nutrient-poor soils, contributing to the sustainable cultivation of Ber in arid zones (Chovatia *et al.* 1993).

## 7. Water Use Efficiency in Ber Cultivation: Irrigation Techniques and Strategies

In arid zones, where water scarcity is a critical constraint to agricultural productivity, enhancing water use efficiency (WUE) is essential for sustainable crop production. Ber (*Ziziphus mauritiana*) stands out as a highly adaptable fruit crop, capable of thriving under minimal water inputs, making it an ideal candidate for cultivation in water-limited environments. The crop's inherent drought tolerance, coupled with strategic irrigation practices, can significantly improve WUE in Ber orchards. One of the most effective irrigation techniques for Ber cultivation in arid zones is drip irrigation ((Christensen, 1998). This method delivers water directly to the root zone, minimizing losses due to evaporation and deep percolation, and ensuring that the trees receive a consistent supply of moisture. Drip irrigation not only enhances WUE but also allows for the precise application of fertilizers through fertigation, further optimizing resource use. Another strategy to improve WUE in Ber cultivation is deficit irrigation, where water is applied at critical growth stages, such as flowering and fruit development, while being reduced or withheld during less sensitive periods. This approach encourages deeper root growth, enabling the trees to access moisture from deeper soil layers, thereby improving drought resilience (Clake *et al.* 2006).

Mulching is also a valuable practice in Ber orchards, as it helps to conserve soil moisture, reduce evaporation, and suppress weed growth, which competes with the trees for water. Organic mulches, such as straw or compost, additionally contribute to soil fertility, promoting healthy root development. The integration of these irrigation techniques and strategies not only maximizes

WUE in Ber cultivation but also contributes to the sustainability of agricultural systems in arid zones. By optimizing water resources, Ber production can be expanded, providing a reliable source of income for farmers while conserving precious water in these challenging environments (Concibido *et al.* 2003).

## **8. Pests and Diseases Affecting Ber in Arid Zones: Challenges and Control Measures**

In arid zones, Ber (*Ziziphus mauritiana*) is increasingly recognized for its ability to thrive under extreme environmental conditions. However, the production of Ber in these regions is significantly challenged by various pests and diseases, which can lead to substantial yield losses and reduced fruit quality. Among the most prevalent pests are fruit flies (*Carpomya vesuviana*) and leaf-eating caterpillars (*Anarsia ephippias*), which cause direct damage to the fruits and leaves, leading to reduced photosynthetic activity and premature fruit drop. Additionally, mealybugs (*Maconellicoccus hirsutus*) and scale insects (*Aspidiotus destructor*) are common sap-sucking pests that weaken the trees by extracting vital nutrients, making them more susceptible to environmental stress (Croser *et al.* 2003).

Fungal diseases, such as powdery mildew (*Oidium erysiphoides*) and leaf spot (*Cercospora spp.*), are also significant concerns in Ber cultivation. These diseases thrive in the fluctuating humidity levels typical of arid zones, causing defoliation, stunted growth, and a decline in fruit production. Moreover, the limited availability of water exacerbates the vulnerability of Ber trees to these diseases, as water-stressed plants are less capable of mounting an effective defense response (Daily and Ellison, 2002).

Control measures for these pests and diseases in arid zones require an integrated approach. Cultural practices, such as timely pruning, proper spacing, and maintaining orchard hygiene, are essential for reducing pest and disease incidence. Biological control, including the use of natural predators and parasitoids, can effectively manage pest populations without harming the environment. Additionally, the application of neem-based biopesticides and other organic treatments offers an eco-friendly alternative to chemical pesticides, which may be less effective in arid conditions due to rapid degradation under high temperatures. Overall, a holistic pest and disease management strategy, tailored to the unique conditions of arid zones, is critical for ensuring sustainable Ber production and improving farmers' resilience against biotic stressors (Datta *et al.* 2000).

## 9. Post-Harvest Management and Value Addition in Ber Production

Post-harvest management is crucial for maximizing the profitability and sustainability of Ber (*Ziziphus mauritiana*) production, especially in arid zones where resources are scarce and market access is limited. After harvesting, Ber fruits are highly perishable, making proper handling, storage, and processing essential to prevent significant post-harvest losses, which can exceed 30% in some cases. Key practices include timely harvesting, gentle handling to minimize bruising, and rapid cooling to preserve fruit quality. In arid zones, where infrastructure for cold storage is often lacking, traditional methods like sun drying are commonly used to extend the shelf life of Ber fruits. However, modern techniques such as controlled atmosphere storage and the use of natural preservatives can significantly enhance fruit preservation. Additionally, the development of value-added products like dried Ber, candies, jams, and beverages offer substantial opportunities for increasing farmers' income and expanding market reach (De Villiers, 2000).

Value addition not only enhances the economic viability of Ber production but also provides an avenue for reducing waste and utilizing surplus production. By converting fresh Ber into processed products, farmers can tap into new markets, including urban consumers and international buyers, where there is growing demand for healthy, natural foods. To fully capitalize on these opportunities, investments in processing technologies, training for farmers, and the development of robust market linkages are necessary. This will ensure that Ber production in arid zones becomes a sustainable and profitable enterprise, contributing to rural development and food security (Deen Dayal Giri *et al.* 2022).

## 10. Market Opportunities and Challenges for Ber Fruits in Arid Regions

The market for Ber (*Ziziphus mauritiana*) fruits in arid regions presents a mix of opportunities and challenges that are pivotal for the success of its cultivation. On the opportunity side, the growing demand for Ber fruits in both domestic and international markets offers significant income potential for farmers. The fruit's nutritional benefits, including high vitamin C content and antioxidant properties, have increased its appeal among health-conscious consumers, further driving demand. Additionally, the versatility of Ber fruits, which can be consumed fresh or processed into products such as dried fruits, candies, and beverages, opens up diverse revenue streams, adding value to the crop (Deng *et al.* 2004).

However, several challenges hinder the full exploitation of these market opportunities. One of the primary challenges is the lack of established marketing channels and infrastructure in many

arid regions. Farmers often face difficulties in accessing markets, leading to lower profitability. The absence of efficient storage facilities exacerbates post-harvest losses, reducing the quantity of marketable produce. Moreover, price volatility and competition from other fruit crops in the market can also impact the income stability of Ber producers. To overcome these challenges, there is a need for investment in market infrastructure, including roads, storage, and processing facilities, as well as the development of reliable market linkages. Additionally, promoting awareness of Ber's health benefits can help in creating a niche market, potentially leading to higher returns for farmers in arid regions (Diamond, 2005).

### **11. Role of Ber in Agroforestry Systems for Sustainable Agriculture in Arid Zones**

Ber (*Ziziphus mauritiana*) offers a promising role in agroforestry systems within arid zones, where traditional agriculture faces significant challenges. Its integration into agroforestry systems can enhance sustainability and resilience in these regions, contributing to both environmental and economic benefits. In arid environments, where water scarcity and soil degradation are prevalent, Ber trees provide valuable ecological functions (Diamond, 2002). Their deep root systems access moisture from deeper soil layers, reducing competition for water with shallow-rooted crops. This adaptation not only sustains Ber trees during dry periods but also helps in maintaining soil moisture levels, benefiting associated crops. Additionally, Ber trees contribute to soil fertility through the accumulation of organic matter from leaf litter, which improves soil structure and nutrient content (Dudley *et al.* 2006).

The inclusion of Ber in agroforestry systems can also help mitigate soil erosion. The dense canopy of Ber trees reduces wind and water erosion, stabilizing soil and preventing further land degradation. This protective role is crucial in arid zones, where erosion can rapidly degrade land quality and reduce agricultural productivity. Economically, Ber's integration into agroforestry systems provides diversified income streams for farmers. In addition to the fruit, which has market potential both locally and internationally, Ber trees offer timber and fodder, enhancing the economic resilience of farming households (Estan *et al.* 2005). The ability to cultivate Ber alongside other crops or livestock maximizes land use efficiency and reduces the risk of total crop failure due to adverse conditions. Moreover, Ber-based agroforestry systems promote biodiversity by creating habitats for various species, thus supporting ecological balance. This biodiversity, in turn, can contribute to natural pest control and pollination, further enhancing agricultural productivity. Ber plays a vital role in agroforestry systems in arid zones by improving soil health,

reducing erosion, and providing economic benefits. Its integration into these systems supports sustainable agriculture by enhancing both environmental stability and farmer livelihoods (Fleischer and Tsur, 2000).

## **12. Environmental Benefits of Ber Cultivation: Soil Stabilization and Desertification Control**

The cultivation of Ber (*Ziziphus mauritiana*) in arid zones offers significant environmental benefits, particularly in soil stabilization and desertification control. As these regions grapple with severe soil erosion and land degradation, Ber's unique characteristics make it an effective tool for environmental restoration and sustainability. Ber trees are well-adapted to the harsh conditions of arid environments, including low rainfall and poor soil fertility. Their deep and extensive root systems play a crucial role in soil stabilization. By anchoring the soil and preventing erosion, Ber trees help maintain soil structure and reduce surface runoff, which is vital in regions prone to wind and water erosion (Fleskens et al. 2007). The root systems of Ber trees also improve soil aeration and enhance the infiltration of water, contributing to the overall health of the soil. In addition to soil stabilization, Ber cultivation contributes to desertification control. Desertification is a major concern in arid and semi-arid regions, where the loss of vegetation cover leads to the expansion of desert areas. Ber trees, with their ability to thrive in nutrient-poor soils and withstand extreme conditions, help combat desertification by providing a vegetative cover that protects the soil from wind and water erosion (Eshed and Zamir, 1995).

The presence of Ber trees increases organic matter content in the soil, which improves soil fertility and promotes the growth of other plants. Furthermore, Ber orchards contribute to the creation of microclimates that can enhance local biodiversity. The shade provided by Ber trees can create a more favorable environment for other plant species, leading to increased plant diversity and ecological resilience. This, in turn, supports a variety of wildlife and contributes to the overall health of the ecosystem. Ber cultivation offers valuable environmental benefits in arid zones by stabilizing soil, controlling desertification, and enhancing biodiversity. Its ability to improve soil health and mitigate land degradation makes it a vital component of sustainable agricultural practices in these challenging environments (Frey *et al.* 1975).

## **13. Future Prospects and Research Needs for Enhancing Ber Production in Arid Zones**

The future of Ber (*Ziziphus mauritiana*) production in arid zones presents substantial opportunities for agricultural innovation and development. As climate change intensifies the challenges faced by traditional crops, Ber's resilience and adaptability to harsh conditions position it as a key candidate for expanding agricultural sustainability in these regions. To realize its full potential, targeted research and strategic interventions are necessary (Gal *et al.* 2021).

#### **Future Prospects:**

**1. Cultivar Improvement:** Advances in breeding and genetic research can yield new Ber cultivars with enhanced traits such as increased drought tolerance, disease resistance, and improved fruit quality. This can lead to higher yields and better adaptability to varying arid conditions (Galnoor, 1980).

**2. Agroforestry Integration:** Ber can be integrated into agroforestry systems, where it can complement other crops and livestock. This approach can enhance soil fertility, improve water use efficiency, and increase overall farm productivity (Gardner, 1991).

**3. Market Expansion:** Expanding market opportunities through value-added products, such as processed Ber fruits and derivatives, can increase profitability and create new revenue streams for farmers (Garrido *et al.* 2006).

#### **Research Needs:**

**1. Water Management Innovations:** Research into efficient irrigation techniques, such as drip irrigation and soil moisture management, is critical for optimizing water use and improving Ber productivity (Gelburd, 1985).

**2. Pest and Disease Management:** Developing integrated pest management strategies and disease-resistant varieties will be essential to mitigate losses and ensure the sustainability of Ber production (Gibson and Noble, 1986).

**3. Soil Health:** Investigating soil amendments and management practices that enhance soil fertility and structure can support better growth and yield of Ber in nutrient-poor soils (Gill *et al.* 2005).

**4. Economic Analysis:** Comprehensive economic assessments are needed to evaluate the cost-effectiveness of various Ber cultivation practices and to identify strategies for improving market access and profitability for farmers (Glennon, 2002).

By focusing on these research areas and leveraging future technological advancements, Ber production can be significantly enhanced, contributing to the sustainable development of agriculture in arid zones.

## **CONCLUSION**

Ber (*Ziziphus mauritiana*) holds significant promise for enhancing agricultural sustainability and food security in arid zones, given its adaptability to harsh environmental conditions. The crop's drought tolerance, minimal water requirements, and ability to thrive in nutrient-poor soils offer a viable solution to the challenges posed by arid climates. However, maximizing its potential requires addressing key challenges, including improving agronomic practices, pest and disease management, and market access. Research and extension services must focus on developing high-yielding, pest-resistant cultivars and disseminating knowledge to farmers. Additionally, investments in infrastructure and policy support are crucial for facilitating market access and ensuring profitability. By overcoming these obstacles, Ber production can contribute to environmental sustainability, boost rural economies, and offer a resilient agricultural model for arid regions facing the impacts of climate change. Future efforts should prioritize these areas to fully realize the benefits of Ber cultivation.

## **CONSENT**

It is not applicable.

## **ETHICAL APPROVAL**

It is not applicable.

## **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

## **REFERENCES**

1. Bar-Yosef, B. 1999. Advances in fertigation. *Advances in Agronomy* 65: 1-77.
2. Ahmed K Rashwan, Naymul K, Mohammad Rezaul Islam Shishir, Tao Bao, Yang Lu, Wei Chen. Jujube fruit: A potential nutritious fruit for the development of functional food products, *Journal of Functional Foods*, 2020, 75.
3. Alexander, DJ. 1990. Managing resources of the Murray-Darling Basin of Australia. *Water Resources Journal* 164: 87-99.
4. Allen, V.G., Brown, C.P., Kellison, R, Segarra, E., Wheeler, T., Dotray, P.A., Conkwright, I C., Green, CJ. and Acosta-Martinez, V. 2005.
5. Arumugam J, Thambidurai S, Suresh S, Selvapandiyam M, Kandasamy M, Pugazhenthiran N, et al. Green synthesis of zinc oxide nanoparticles using *Ficus carica* leaf extract and

their bactericidal and photocatalytic performance evaluation, *Chemical Physics Letters*, 2021, 783.

6. Aslam, M. and Prathapar, S.A. 2006. Strategies to Mitigate Secondary Salinization in the Indus Basin of Pakistan: A Selective Review. Research Report 97. International Water Management Institute. Colombo, Sri Lanka. 31 pp.
7. Athawale, Gauri, Sanjay, Akbari. Aonla and its processing. *International Journal for Green and Herbal chemistry*, 2020, 62-73.
8. Ayars, I.E., Christen, E.W. and Hornbuckle, I.W. 2006. Controlled drainage for improved water
9. Ayars, R.S. and Westcot, D.W. 1985. Water quality for agriculture. FAO Irrigation Drainage Paper 29. FAO, Rome.
10. Azhane Ahmad, Asgar Ali, Improvement of postharvest quality, regulation of antioxidants capacity and softening enzymes activity of cold-stored carambola in response to polyamines application, *Postharvest Biology and Technology*. 2019; 148:208-217.
11. Balendres MA, Bengoa JC. Diseases of dragon fruit (*Hylocereus* species): Etiology and current management options, *Crop Protection*, 2021, 126.
12. Beltran, J.M. 1999. Irrigation with saline water: Benefits and environmental impact. *Agricultural Water Management* 40: 183-194.
13. Beltran, J.M. and Koo-Oshima, S. 2006. Water desalination for agricultural applications. FAO Land and Water Discussion Paper 5. FAORome.
14. Ben-Gal, A., Lazorovitch, N. and Shani, V. 2004. Subsurface drip irrigation in gravel filled cavities. *The Vadose Zone Journal* 3: 1407-1413.
15. Ben-Gal, A., Tal, A. and Tel-Zur, N., 2006. The sustainability of arid agriculture: Trends and challenges. *Annals of Arid Zone*, 45(3/4), p.227.
16. Benzioni, A. 1995. Jojoba domestication and commercialization in Israel. In *Horticulture Review* (Ed. I Janick), Vol. 17, pp. 231-264.
17. Benzioni, A., Palzkill, D.A. and Nelson, I.M. 1992. Flower bud dormancy, ABA concentration, and survival during frost of jojoba genotypes under water stress. *Journal of American Society of Horticulture Science* 117: 976-980.
18. Benzioni, A., Shiloh, E. and Ventura, M. 1999. Yield parameters in young jojoba plants and their relation to actual yield in later years. *Industrial Crop Production* 10: 85-95.

19. Bernstein, L. 1975. Effects of salinity and sodicity on plant growth. *Annual Review of Phytopathology* 13: 295-312.
20. Bonfil, DJ., Goren, Mufradi, Lihtensveig, I and Abbo, S. 2007. Development of early-flowering Kabuli chickpea with compound and simple leaves. *Plant Breeding* 126: 125-129.
21. Brown, L. 2004. World Food Security Deteriorating: Food Crunch in 2005. Now Likely. Earth Policy Institute, Washington. Brown, Lester 2006. Ethanol could leave the world hungry. *Fortune*, August 16 2006.
22. Cai, X., McKinney, D.C. and Rosegrant, M.W. 2001. Sustainability Analysis for Irrigation Water Management in the Aral Sea Region. International Food Policy Research Institute, Washington, DC. Discussion Paper, 48 p.
23. Carr, R.M., Blumenthal, V.J. and Duncan Mara, D. 2004. Guidelines for the safe use of wastewater in agriculture: Revisiting WHO guidelines. *Water Science and Technology* 50:31-38.
24. Chaojun Yan, Hafiz Muhammad Rizwan, Dangdi Liang, Michael Reichelt, Axel Mithöfer, Sandra S Scholz, Ralf Oelmüller. 2021. The effect of the root-colonizing *Piriformospora indica* on passion fruit (*Passiflora edulis*) development: Initial defense shifts to fitness benefits and higher fruit quality, *Food Chemistry*, 359.
25. Chovatia, R.S., Patel, D.S. and Patel, G.V. 1993. Performance of ber (*Ziziphus mauritiana* Lark) cultivars under arid conditions. *Annals of Arid Zone* 32: 215-217.
26. Christensen, P. 1998. Middle eastern irrigation: Legacies and lessons. In *Transformations of the Middle Eastern Environments: Legacies and Lessons* (Eds. I Coppock and Joseph Miller) New Haven, Yale School of Forestry and Environmental Studies Vol. 103, pp. 15-31.
27. Clarke, H.J., Wilson, J.G., Kuo, I., Lulsdorf, M.M., Mallikarjuna, N., Kuo, I and Siddique, K.H.M. 2006. Embryo rescue and plant regeneration in vitro of selfed chickpea (*Cicer arietinum* L.) and its wild annual relatives: *Plant Cell*
28. Concibido, V.C., La Vallee, B., McIaird, P., Pineda, N., Meyer, I., Hummel, L., Yang, J., Wu, K and Delannay, X. 2003. Introgression of a quantitative trait locus for yield from *Glycine soja* into commercial soybean cultivars. *Theoretical and Applied Genetics* 106: 575-582.

29. Croser, I.S., Ahmad, F., Clarke, H.J., Siddique, K.H.M. 2003. Utilization of wild Cicer in chickpea improvement-progress, constraints, and prospect. *Australian Journal of Agriculture Research* 54: 429-444.
30. Daily, G. and Ellison, K 2002. *The New Economy of Nature, the Quest to Make Conservation Profitable*. Washington, Island 13. Darwin, C. 1859. *The Origin of Species by Means of 'Natural Selection (Reprint)*. New York: Collier, 1962.
31. Datta, KK, Jong, C. and de Singh, O.P. 2000. Reclaiming salt-Affected land through drainage in Haryana, India: A financial analysis. *Agricultural Water Management* 46: 55-71.
32. de Villiers, M. 2000. *Shrinking Aquifers Water*. Stoddart, Toronto. Diamond, J. 1997. *Guns, Germs and Steel: The Fate of Human Societies*. New York, Norton.
33. Deen Dayal Giri, Alaa Alhazmi, Akbar Mohammad, Shafiul Haque, Neha Srivastava, Vijay Kumar Thakur, Vijai Kumar Gupta, et al. Lead removal from synthetic wastewater by bio-sorbents prepared from seeds of *Artocarpus Heterophyllus* and *Syzygium Cumini*, *Chemosphere*, 2022, 287(1),
34. Deng, Q.Y., Yuan, L.P., Liang, F.S., Li, I., Wang, L.G. and Wang, B. 2004. Studies on yield-enhancing genes from wild rice and their marked-assisted selection in hybrid rice. *Hybrid Rice* 19: 6-10.
35. Diamond, J. 2005. *Collapse, How Societies Choose to Succeed or Fail*. Viking Press, New York.
36. Diamond, J. 2002. Evolution, consequences and future of plant and animal domestication. *Nature* 418: 700-707.
37. Dudley, L., Ben-Gal, A. and Shani, U. 2006. Influence of plant, soil and water properties on the leaching
38. Estan, M.T., Martinez-Rodrigues, M.M., Perez-Alfocea, E, Flowers, T.J. and Bolarin, M.C. 2005. Grafting raises the salt tolerance of tomato through limiting the transport of sodium and chloride to the shoot. *Journal of Experimental Botany* 56: 703-712.
39. Fleischer, A.I. and Tsur, A.Y. 2000. Measuring the recreational value of agricultural landscape. *European Review of Agricultural Economics* 27: 3.

40. Fleskens, K., Ataev, A., Mamedov, B. and Spaan, W.P. 2007. Desert water harvesting from takyr surfaces: Assessing the potential of traditional and experimental technologies in the Karakum. *Land Degradation and Development* 18: 17-39.
41. Eshed, Y. and Zamir, D. 1995. An introgression line population of *Lycopersicon pennellii* in the cultivated tomato enables the identification and fine mapping of yield associated QTL. *Genetics* 141: 1147-1162.
42. Frey, K.J., Hammond, E.G. and Lawrence, P.K. 1975. Inheritance of oil percentage in interspecific crosses of hexaploid oats. *Crop Science* 15: 94-95.
43. Gal, A.B., Tal, A. and Zur, T. 2021. The Sustainability of Arid Agriculture: Trends and Challenges. *Annals of Arid Zone*, 45(3 & 4).
44. Galnoor, I. 1980. *Water Policy Making in Israel, Water Quality Management Under Conditions of Scarcity, Israel as a Case Study*, New York, Academic Press, 296 p.
45. Gardner, W.R. 1991. Modeling water uptake by roots. *Irrigation Science* 12: 109-114.
46. Garrido, A., Martinez-Santos, P. and Llama, R. 2006. Groundwater Irrigation And Its Implications For Water Policy In Semiarid Countries: The Spanish Experience. *Hydrogeology Journal* 14: 1431-1435.
47. Gelburd, D.E. 1985. Managing salinity. Lessons from the past. *Journal of Soil and Water Conservation* 40: 329-331.
48. Gibson, C.G. and Noble, P.S. 1986. *The Cactus Primer*. Harvard Univ, Press, Cambridge, MA. Gintzburger, G, Le Houerou, H. and Toderich, K. 2005. The Steppes of Middle Asia: Post-1991 agricultural and rangeland adjustment. *Arid Land Research and Management* 19: 215-239.
49. Gill, A., Fadil Aw-Hassan, R. and Aden BtiyUkalaca, O. 2005. Economic analysis of energy use in groundwater irrigation of dry areas: A case study in Syria. *Applied Energy* 82: 285-299.
50. Glennon, R. 2002. *Water Follies, Groundwater Pumpnig and the Fate of America's Fresh Waters*. Island Press, Washington. Gold, M. 1999. *Sustainable Agriculture: Definitions and Terms*. Special Reference Briefs Series No. SRB 99-02, US Department of Agriculture.
51. Goldberg, D. and Schmueli, M. 1970. Drip Irrigation: A method used under arid and desert conditions of high water and soil salinity. *Transactions of the ASAE* 13: 38-41.

52. Gupta R, Sharma, Alka, Kumar, Pradeep Vishwakarma, Rajesh, et al. Effect of blanching on thin layer drying kinetics of aonla (*Emblica officinalis*) shreds. *Journal of Food Science and Technology*, 2014, 51.
53. Gur, A. and Zamir, D. 2004. Unused natural variation can lift yield barriers in plant breeding. *Public Library of Science (PloS) Biology* 2: e245.
54. Hamoda, M.F. 2001. Desalination and water resource management in Kuwait. *Desalination* 138: 385-393.
55. Hellegers, P. 2006. The role of economics in irrigation water management. *Irrigation and Drainage* 55: 157-163.
56. Herczeg, A.L., Simpson, I.H. and Mazor, E. 1993. Transport of soluble salts in a large semiarid basin: River Murray, Australia. *Journal of Hydrology* 144: 59-84.
57. Heslop-Harrison, I.S. 2002. Exploiting novel germplasm. *Australian Journal of Agricultural Research* 53: 873-879.
58. Hodgson, J. 2002. Nutraceuticals tread business tightrope in Europe. *Nature Biotechnology* 20: 322-323.
59. Howell, T. 2001. Enhancing water use efficiency in irrigated agriculture. *Agronomy Journal* 93: 281-289.
60. Huang, S.B., Spielmeyer, W., Lagudah, E.S., James, R.A., Platten, J.D., Dennis E.S. and Munns, R. 2006. A sodium transporter (HKT1) is a candidate for Nax1, a gene for salt tolerance in Durum wheat. *Plant Physiology* 142:1719-1727.
61. Huang, X.Q., Coster, H., Ganai, M.W. and M.S. Roder. 2003. Advanced backcross QTL analysis for the identification of quantitative trait loci alleles from wild relatives of wheat (*Triticum aestivum* L.) *Theoretical and Applied Genetics* 106: 1379-1389.
62. Inbar, Y. 2002. Standards and regulations for treated wastewater reuse in Israel, Regional Initiatives for Dryland Management, Regional Workshop on TWW and Biosolids Use in Dry Areas, Granda, Spain, October 21-24, 2002. Jacobson, T. and Adams, R.M. 1958. Salt and silt in ancient Mesopotamian agriculture. *Science* 128: 1251-1258.
63. Integrating Cotton and Beef Production to Reduce Water Withdrawal from the Ogallala Aquifer in the Southern High Plains. *Agronomy Journal* 97: 556-567.

64. Jackson, W., Berry, W. and Colman, B. (Eds.) 1984. Meeting the Expectations of the Land. North Point Press, San Francisco. Janmaat, 1. 2004. Calculating the cost of irrigation induced soil salinization in the Tungabhadra project. *Agricultural Economics* 31: 81-96.
65. Jamkhande PG, Barde SR, Patwekar SL, Tidke PS. Plant profile, phytochemistry and pharmacology of *Cordia dichotoma* (Indian cherry): A review. *Asian Pacific J Tropical Medicine*. 2013;3(12):1009–1012
66. Jawanda, J.S. and Bal, J.S. 1978. The ber, highly paying and rich in food value. *Indian Horticulture* 23(3): 19-2.
67. Jnana Bharati Palai, Jagadish Jena, Sagar Maitra. Prospects of underutilized food legumes in sustaining pulse needs in India—A review. 2019, 82-88.
68. Joshi V. Some promising underutilised industrial crops for cultivation on wastelands of India. *Green page: Article. Natur. Prod. Radiance*. 2005;4(5):396-403.
69. Kahlowan, M.A. and Azam M. 2002. Individual and combined effect of waterlogging and salinity on crop yields in the Indus basin. *Irrigation and Drainage* 51: 320-338.
70. Kakade, Vijaysinha, Jinger, Dinesh, Dayal, Vishambhar, et al. Dragon fruit *Food & Scientific Reports*. 2020 Dec;1:44-48.
71. Kameswara Rao, N., Reddy, LJ. and Bramel, PJ. 2003. Potential of wild species for genetic enhancement of some semi-arid food crops. *Genetics Research Crop Evolution* 50: 707-721.
72. Kington, E.A. and Pannel, DJ. 200 1. Dryland Salinity in the Upper Kent River Catchment of Western Australia, Farmer Perceptions and Practices. SEA Working Paper, University of Western Australia.
73. Kumar, J. and Abbo, A. 2001. Genetics of flowering time in chickpea and its bearing on productivity in semiarid environments. *Advances in Agronomy* 72: 107-138.
74. Lavee, D. 2003. Cost-Benefit Analysis for National Upgrade of Treated Wastewater Quality in Israel, Tel Aviv, Pareto Engineering, Ltd. (in Hebrew).
75. Li Fu, Junnan Yang, Hongli Shang, Jianxin Song, Changes of characteristic sugar, fatty acid, organic acid and amino acid in jujubes at different dry mature stages, *Journal of Food Composition and Analysis*, 2021, 104.
76. management in arid regions irrigated agriculture.

77. Padulosi S, Hodgkin T, Williams JT, Haq N. Underutilized crops: trends, challenges and opportunities in the 21st Century, 2002, 1-23.
78. Pareek, Sunil, Yahia, Elhadi. Postharvest Biology and Technology of Ber Fruit, 2013, 201-240.
79. Pengrui AI, Yingjie MA, Ying Hai, Influence of jujube/cotton intercropping on soil temperature and crop evapotranspiration in an arid area, Agricultural Water Management, 2021, 256.
80. Rao, N.S., 2016. Arid Zone Fruit Crops. Abiotic Stress Physiology of Horticultural Crops, pp.223-234.
81. Rathore, V.S., Tanwar, S.P.S. and Yadav, O.P., 2019. Integrated farming system: Key to sustainability in arid and semi-arid regions. The Indian Journal of Agricultural Sciences, 89(2), pp.181-192.
82. Ruhollah A, Akhtar S, Abolfazl J, In-vitro Tetraploidy induction creates enhancements in morphological, physiological and phytochemical characteristics in the fig tree (*Ficus carica* L.), Plant Physiology and Biochemistry, 2021, 166.
83. Ryan, J.G. and Spencer, D.C., 2001. Future challenges and opportunities for agricultural R&D in the semi-arid tropics. International Crops Research Institute for the Semi-Arid Tropics.
84. Simrandeep Kour, Prashant Bakshi, Arti Sharma, Wali VK, Amit Jasrotia, Shilpy Kumari. Strategies on Conservation, Improvement and Utilization of Underutilized Fruit Crops, 2018, 638-650.
85. Sukirti Upadhyay, Prashant Upadhyay, Ghosh AK, Vijender Singh. *Ziziphus mauritiana*: a review on pharmacological potential of this underutilized plant, IJCRR. 2012 February;4(3):141-144.
86. Thiyagarajan, Rajasekaran G, Balamurugan S, Karthikeyan S. Studies on the effect of phytochemical and mineral analysis of kalakai (*Carissa carandas* L.) Inter. J of current research Fruits. 2005;12(10):14066-14072.
87. Tissue and Organ Culture 85: 197-204. Clemings, R. 1996. Mirage: The False Promise of Desert Agriculture. San Francisco, Sierra Club Books. 247 pp.
88. Wiley, New York, Benzioni, A. and Nerd, A. 1985. Effect of irrigation and fertilization on vegetative growth and yield of jojoba in relation to water status of the plants. In Jojoba.

Proceedings of 6th International Conference on Jojoba and its Uses (Eds. I Wisniak and I Zabisky), pp. 201-212.

UNDER PEER REVIEW