

## Review Article

# PEARL MILLET (*Pennisetum glaucum*): A CLIMATE RESILIENT AND NUTRITIONALLY SIGNIFICANT CROP FOR GLOBAL FOOD SECURITY

## ABSTRACT

The environmental balance needed for agriculture is severely strained due to changes in rainfall patterns and climate change, as well as the growing global need for food. As global temperatures rise and precipitation patterns become more erratic, millets offer an efficient solution to the trade-off between food security and water availability. Millets are rich in nutrients and possess a special ability to endure extreme conditions such as heat, drought, and various other abiotic and biotic stresses. Pearl millet is a highly climate-resilient crop capable of mitigating the adverse impacts of climate change. Antioxidant gene expression, presence of a waxy outer layer and heat shock proteins/chaperons contribute to its abiotic stress tolerance. By adjusting its flowering phenology and establishing fast-growing roots, pearl millet enhances its water and nutrient use efficiency. Stable sources of germplasm in pearl millet that exhibit resistance to diseases such as downy mildew, blast, rust, smut, and ergot have also been incorporated into disease resistance breeding programs. Pearl millet plays a crucial role in nourishing the poverty-stricken populations due to its high palatability and rich nutritional profile. Various bioactive compounds pertain to hypoglycemic, anticancerous and probiotic properties that contribute to its health benefits. Biofortification in pearl millet offers a solution to address issues of micronutrient deficiency in vulnerable populations. Continued research and investment in pearl millet production not only bolster food security but also contribute to the resilience of agricultural systems in the face of increasingly unpredictable environmental conditions. Thus, embracing pearl millet represents a forward-thinking approach to ensuring food security and promoting nutritional diversity on a global scale.

*Key words:* Abiotic stress; biotic stress; climate change; food crisis; nutrition; value addition

## 1. INTRODUCTION

Agriculture plays a crucial role in enhancing global food security, especially in the face of climate change. As global temperatures rise and precipitation patterns become more erratic, millets offer an efficient solution to the trade-off between food security and climate fluctuation. An estimated 2 to 3 billion people may experience food insecurity and hunger by 2050 as a result of declining food production rates as agricultural systems struggle to adapt to

rapidly shifting environmental conditions and the additional burden of feeding a population expected to surpass 9 billion[1].

The environmental balance necessary for agriculture is being severely strained not just by the increase in food demand but also by the altering temperature and patterns of precipitation. Events like more severe and frequent floods and droughts as well as irregular rainfall are clear indicators of this. Further deficits in the world's food supplies may result from the fragile balance between agriculture and climate.

The war in Ukraine has severely disrupted global food supplies, leading to a sharp rise in prices for staple crops like rice, wheat and barley. Many vulnerable countries that rely heavily on food imports are being hit particularly hard. Over 1.4 billion people live in highly exposed countries, mostly in Africa and Asia. They face elevated inflation, budgetary stress, and rising malnutrition levels as they struggle to protect their populations from skyrocketing food prices.

Meanwhile, trade restrictions imposed by countries trying to shore up domestic supplies are further exacerbating the global food crisis. The World Bank estimates that over 800 million people worldwide already suffer from high levels of malnutrition or undernourishment. Urgent action is needed to prevent the situation from deteriorating further. The World Bank has committed \$45 billion to tackle the crisis, including measures to expand social protection, boost agricultural productivity, and facilitate trade[2]. However, long-term solutions require transforming food systems to be more sustainable, resilient, and equitable.

Because of rising food prices, falling household incomes, and rising unemployment, economic recessions can leave communities more vulnerable. Due to its ability to diversify farmers' crop rotation and decrease reliance on a single crop, millets have the potential to play a major role in ensuring food security during difficult times. Because this diversification ensures alternate sources of food if one crop fails and it could assist boost food security[3].

Millets are highly climate-resilient crops that can thrive in arid and semi-arid regions with minimal water and input requirements, making them well-suited to withstand the effects of climate change. Millets, being highly nutritious, may help address the issue of malnutrition prevalent in many regions of the world.

**Global millet production** is estimated to be 54.75 million tonnes in 2022-23, up from 51.1 million tonnes in the previous year. India has seen a steady increase in the production of both major cereals and millets in recent years. The grains can withstand temperatures as high as **26°C to 42°C** with **70% higher water use efficiency** than rice. Since millets can withstand

high temperatures, they are a great option for tropical nations. Millets are "usually the last crops standing in droughts," according to the International Crops Research Institute for the Semi-arid Tropics[4]. However, the country faces the challenge of addressing the growing "cereal gap" and leveraging the potential of climate-resilient and nutritious crops like millets to enhance food security and sustainability.

## **2. PEARL MILLET:**

Pearl millet, also known as bajra, is one of the most widely grown and important millets globally. Pearl millet is highly nutritious, containing high levels of protein, fiber, minerals like iron, zinc, and calcium, and antioxidants. Compared to other cereals like rice and wheat, shifting to pearl millet and sorghum can increase protein (46%), iron (353%), and zinc (82%) production which makes it highly beneficial for addressing micronutrient deficiencies. Pearl millet is an efficient gluten-free grain and it is highly recommended for people who have celiac disease or gluten intolerance. It is advantageous for managing diabetes because it is low in fat and has a low glycemic index.

Pearl millet is highly resistant to abiotic stresses like drought, heat, and low soil fertility. It can thrive in harsh climatic conditions where other cereals may fail. Pearl millet cultivation is more resilient to the impacts of climate change compared to other cereals and it shows considerable tolerance to various biotic stresses through natural resilience and tolerance mechanism. Its small grains, nutritious composition, and stress resistance make it a promising crop for the future. In the Indo-Gangetic Plains, switching from rice to pearl millet and sorghum during the Kharif season can reduce water use by 32% [5]. This is crucial for sustainable water conservation in water-stressed regions.

India is the world's largest producer of pearl millet, accounting for 44% of global millet production. The area under cultivation of Pearl millet in India for the 2023-24 crop year is 7.41 million hectares with the production of around 10.3 million tonnes and productivity around 1391 kg per hectare. The major pearl millet producing states in India are Rajasthan (44% of production), Uttar Pradesh (15%), Haryana (11.4%), Maharashtra (10.8%), and Gujarat (10.5%) [6]. While pearl millet consumption has declined in recent decades, it is now gaining importance due to its climate resilient nature, nutritional status and use as food and feed products.

## **3. CLIMATE RESILIENCE OF PEARL MILLET**

Pearl millet can withstand harsh growth circumstances like drought, low soil fertility, and high temperatures and it is usually grown as a rainfed crop with little input. It can produce superior yields in stressed environments where other cereals like wheat and rice fail to provide economic returns. Its capacity to endure higher temperatures and thrive in drought-prone regions, makes it a climate-resilient crop capable of mitigating the adverse impacts of climate change[7].

### 3A. TOLERANCE TO ABIOTIC STRESS

Abiotic stresses negatively impact the growth and yield potential of all crops, including pearl millet. Therefore, identifying and utilizing genetic variations for abiotic stress tolerance is crucial for improving its adaptation to these stresses. Flowering time and photoperiodic sensitivity significantly impact the yield and yield stability of pearl millet. Pearl millet typically exhibits photoperiod sensitivity, with most landraces flowering under short-day conditions[8]. This trait allows for flexible sowing dates, ensuring that flowering and grain maturity coincide with the end of the growing season. Research indicates that pearl millet has adaptive mechanisms (morphological- root system, leaf structure, leaf angle; physiological- osmotic adjustment, stomatal regulation, heat shock proteins; anatomical- cuticle thickness, sunken stomata; biochemical- antioxidants defence, secondary metabolites) demonstrating its adaptability to varying environmental conditions[9].

Pearl millet has a deep and extensive root system that allows it to access water from deeper soil layers, which is crucial for surviving prolonged periods of drought. These root features are linked to many traits associated with abiotic stress tolerance, including crop performance, water and nutrient use efficiency. Pearl millet is largely drought-tolerant because of its capacity to retain high leaf water status and increase total root length in the face of soil drying[10]. During water scarcity, nitrogen uptake decreases because the plant's demand for nitrogen diminishes as its growth rate slows. Additionally, nitrogen absorption by roots relies on the presence of water, which transports solutes to the soil-root interface[11]. Pearl millet's capability to maintain sustained water uptake through its fast-growing roots enhances its water and nutrient use efficiency. It had been revealed according to rhizotron and micro-computed tomography (micro CT) phenotyping techniques that early-stage root development in pearl millet, characterized by rapid primary root growth and distinct lateral roots, is crucial for early-stage drought stress tolerance[12]. Pearl millet is a C4 grass that has evolved naturally to use water efficiently in high light conditions through the usage of the

NADP-dependent malic enzyme pathway [13]. By separating carbon fixation in the bundle sheath cells from the light reactions of photosynthesis in the mesophyll cells, this mechanism lessens the rivalry between oxygen and CO<sub>2</sub> molecules for the Rubisco enzyme. As a result, compared to C3 grasses like wheat or rice, this minimizes the need for extended stomatal opening, which boosts water usage efficiency.

Numerous studies on antioxidant gene expression profiling have been conducted in pearl millet under drought conditions. Genes encoding drought tolerance were significantly upregulated in drought-tolerant genotypes treated with polyethylene glycol to mimic drought stress at both early and late seedling stages. Such genes are ascorbate peroxidase (APX), glutamyl-tRNA reductase (GlutR), and superoxide dismutase (SOD)[14]. This increased gene expression in tolerant genotypes indicates their role in scavenging reactive oxygen species (ROS) during drought stress[15]. A transcriptomics analysis further supported these findings, revealing significant up-regulation of differentially expressed genes (DEGs) encoding ROS-scavenging enzymes like SOD, APX, and glutathione peroxidase (GPX) during the vegetative stage under drought conditions[16].

Another method for drought resistance involves the presence of a waxy outer layer [17]. In the pearl millet genome assembly, which includes approximately 38,579 genes, there is a significant abundance of genes related to the biosynthesis of cutin, flavonoids, suberin, and wax[18]. Genes differentially expressed in tolerant genotypes were primarily linked to secondary metabolite pathways like mevalonate, shikimate, alkaloids, phenols, flavonoids, lignin, and wax biosynthesis, as well as stress-related phytohormones such as ABA, ethylene, gibberellic acid, jasmonic acid, and salicylic acid[19].

During flowering, pearl millet can withstand temperatures as high as 26°C to 42°C, which sets it apart from other main food crops like rice, wheat, and maize. Pearl millet has greater tolerance to high temperature stress compared to maize[20]. It responds to heat stress by producing various proteins, notably heat shock proteins (HSPs), which play crucial roles in stress tolerance. These proteins aid in stress signal transduction and gene activation, with HSPs/chaperones facilitating heat shock factor (HSF) activity and defense mechanisms against reactive oxygen species (ROS) that arise from heat stress, thus preventing protein misfolding. Genes encoding calnexin and calreticulin, which process misfolded peptide chains in response to heat stress, were upregulated and differentially expressed, underscoring the ER system's critical role in heat resistance[21].

Salinity stress significantly reduces crop productivity and yield potential. Factors such as low precipitation, irrigation using saline water, and inadequate irrigation practices contribute to this stress. Proline, total soluble protein, and epicuticular wax content are among the several physiological and biochemical reactions that pearl millet cultivars that have acclimated to saline conditions show when salinity levels rise. [22].

Transcriptomic studies in pearl millet have identified several salinity stress-responsive genes including PgDHN, PgDREB2A, PgVDAC, and PgNHX1 [23]. Identified salinity-tolerant pearl millet lines (ICMB 02111, ICMB 94555, ICMB 95333, ICMB 00888, ICMB 01222, ICMP 451, IP 3732, IP 3757, ICMV 93753, and ICMV 94474) should be incorporated into breeding programs to develop locally adapted cultivars with enhanced salinity tolerance. [14].

### 3B. TOLERANCE TO BIOTIC STRESS

Crops that are subjected to abiotic stress are also more vulnerable to weeds, infections, and other insects, which greatly reduces crop yields. Occurrence of infection and disease are facilitated by moderate temperatures (20–30°C) and high relative humidity (85–90%). A number of organisms, including viruses, bacteria, fungus, insects, pests, birds, and weeds are the biotic stressors that can seriously hinder worldwide agricultural productivity. Pearl millet, like other cereals, is vulnerable to these biotic stresses. However, due to its predominant cultivation in dry climates, millets generally face lower risks of biotic stress compared to other crops. Various fungal infections such as downy mildew (*Sclerospora graminicola*), blast (*Magnaporthe grisea*), rust (*Puccinia substriata* var. *indica*), ergot (*Claviceps fusiformis*), and smut (*Tolyposporium penicillariae*) are particularly detrimental to the growth and yield of pearl millet compared to other pathogens [24].

Downy mildew, caused by *Sclerospora graminicola*, is a significant disease of pearl millet, originally traced to Africa. It poses a major biotic challenge in India and West Africa, leading to pearl millet grain yield losses estimated between 10% to 80% [25]. Blast disease, caused by *Magnaporthe grisea*, has affected nearly all pearl millet-growing states in India since 1970. Despite various control methods such as seed sanitation and chemical treatments against seed, soil, and airborne pathogens, emphasis has primarily been placed on developing host-plant resistance due to its effectiveness. Major insect pests of pearl millet include shoot fly (*Atherigona approximata*) (yield loss- 40%), grey weevil (*Mylocerus* spp.) (yield loss-

30%), and white grub (*Holotrichia consanguinea*) (yield loss- 50%). Other minor pests causing economic damage include stem borers, hairy caterpillars, cutworms, chaffer beetles, blister beetles, and grasshoppers. Presence of weeds throughout the season can result in 43.8% decrease in pearl millet grain yield. Common weeds in pearl millet include *Trianthema portulacastrum*, *Echinochloa colona*, *Digeria arvensis* and *Cyperus rotundus*. Since most of India's pearl millet-growing regions have lower rates of insect pests, official breeding efforts that expressly target insect resistance are not yet well-established. Most people agree that the best way to manage diseases in different crops is to utilize resistant cultivars. Key tactics used in resistance breeding against biotic stresses include efficient screening techniques, the use of a variety of germplasm, the identification of resistant sources, an understanding of the genetic basis of resistance, research on virulence variability, the application of efficient resistance breeding techniques, and monitoring cultivar performance in field settings. Researchers have successfully identified stable sources of germplasm in pearl millet that exhibit resistance to diseases such as downy mildew, blast, rust, smut, and ergot, some of which have been incorporated into disease resistance breeding programs. Mapping, investigating, and implementing efficient blast resistance tactics can be accomplished by leveraging the additional genetic and genomic resources that are currently available [26]. Furthermore, wild relatives of pearl millet germplasm present opportunities for strengthening pearl millet resistance to biotic stress and for germplasm enhancement [24]. Various resistant genotypes for downy mildew, blast, rust, smut, ergot, and other biotic stresses in pearl millet, preserved at ICRISAT and NBPGR had been catalogued [27].

Transcriptomic analysis using next-generation sequencing tools in pearl millet elucidated that resistant genotypes activate genes in the phenylpropanoid pathway, induce hypersensitive responses, and enhance systemic acquired resistance as effective defense mechanisms against downy mildew infection [28]. Moreover, various advanced genomic and biotechnological tools have been applied to identify resistant sources, quantitative trait loci, and genes for different biotic stresses [14]. 305 accessions of *Pennisetum violaceum*, a wild relative of pearl millet, under greenhouse conditions against multiple pathotype isolates of *Magnaporthe grisea* and a local isolate of *Pennisetum substriata* var. *indica* were screened to identify diverse sources of resistance against blast and rust diseases [24]. Integrated approaches encompassing stress biology, genomics, and bioinformatics are pivotal in developing crops resilient to stress.

**Table 1. Genes associated with abiotic and biotic stress in pearl millet**

Gene/QTL	Source	Function	Tolerance
Afp	<i>Aspergillus giganteus</i>	Formation of antifungal protein	Downy mildew
Rab 7	<i>Pennisetum glaucum</i>	Small GTP- binding protein	Abiotic stress
VDAC	<i>Pennisetum glaucum</i>	Voltage dependent anion channel	Salt stress
Hsc 70/ HSP 90	<i>Pennisetum glaucum</i>	Molecular chaperons	Abiotic stress
LEA	<i>Pennisetum glaucum</i>	Formation of late embryogenesis abundant protein	Heat and salinity stress
NPR1	<i>Brassica juncea</i>	Non-expressor pathogenesis related genes	Downy mildew

[14]

## **4. NUTRITIONAL AND ANTI-NUTRITIONAL PERSPECTIVES OF PEARL MILLET**

### **4A. NUTRITIONAL PERSPECTIVES**

Pearl millet plays a crucial role in nourishing the poverty-stricken populations due to its high palatability and rich nutritional profile [29]. It includes a high content of dietary fiber, and minerals such as calcium, iron, potassium, zinc, and magnesium, as well as B vitamins. It also serves as an excellent source of well-balanced protein, particularly abundant in threonine, although it lacks lysine and has lower leucine levels compared to sorghum. Notably, pearl millet contains ample tryptophan levels, surpassing other cereal crops. Its nutritional value stands out among millet crops, boasting high concentrations of calcium, iron, zinc, lipids, and high-quality proteins, making it a superior choice for addressing malnutrition and food security challenges in resource-constrained regions (Table 2).

Pearl millet is also low in glycemic index and gluten, which makes it a good food for people with gluten sensitivity and for controlling blood sugar levels. However, the bioavailability of these nutrients may be impacted by the presence of anti-nutritional elements such as tannins and phytic acid [30].

**Table 2. General nutritional composition of Pearl Millet**

<b>Proximate composition</b>	<b>Value per 100g</b>
Moisture (g)	7.4
Energy (kcal)	378
Protein(g)	11.0
Fat (g)	4.2
Total dietary fibre (g)	8.5
Carbohydrate (g)	72.8
Minerals (g)	2.3
Ash (g)	2.1
Crude fibre (g)	2.2
Total Phenol (mg)	7.3

<b>Vitamins</b>	<b>Value per 100g</b>
Thiamine (B <sub>1</sub> )	0.42 mg
Riboflavin (B <sub>2</sub> )	0.29 mg
Niacin (B <sub>3</sub> )	4.72 mg
Pantothenic acid (B <sub>5</sub> )	0.85 mg
Vitamin (B <sub>6</sub> )	0.38 mg
Folate (B <sub>9</sub> )	85 µg
Vitamin K	0.9 µg
<b>Minerals</b>	<b>Value per 100g</b>
Calcium (Ca)	8 mg
Iron (Fe)	3 mg
Magnesium (Mg)	114 mg
Manganese (Mn)	1.6 mg
Phosphorous (P)	285 mg
Potassium (K)	195 mg
Sodium (Na)	5 mg
Zinc (Zn)	1.7 mg
Copper (Cu)	0.8 mg
Sulphur (S)	2.7 µg
<b>Amino acids</b>	<b>g/100 g</b>

Leucine	10.7
Isoleucine	4.4
Valine	4.9
Threonine	4.0
Arginine	4.6
Lysine	3.1
Methionine	1.1
Cysteine	1.5
Tryptophan	1.4
Glutamic acid	23.0
Alanine	8.7
Proline	5.8
Aspartic acid	8.5
Phenylalanine	4.4
Tyrosine	3.0
Histidine	2.3
Glycine	2.7
Serine	5.2

Source: USDA Nutrient Database [<http://ndb.nal.usda.gov/ndb/search/list>].

#### **4B. Health enhancing properties**

The inclusion of pearl millet in the diet not only enhances nutritional security but also offers potential health benefits, making it a valuable addition to combat various health issues and promote overall well-being.

##### **1. Hypoglycemic properties**

Pearl millet exhibits significant hypoglycemic properties due to its bioactive peptides, phenolics, and phytochemicals. Proteolytic enzymes (lucalase, bromelain, and chymotrypsin) generate pearl millet protein hydrolysates with potent anti-diabetic potential, inhibiting  $\alpha$ -amylase and dipeptidyl peptidase-IV and has lesser glycemic index than many cereal grains. Additionally, pearl millet flour blended with sweet detar flour or sodium carboxymethyl cellulose shows enhanced  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibitory activities, crucial for managing type 2 diabetes [31]. Phenolics

in pearl millet play a vital role in inhibiting carbohydrate-digesting enzymes and regulating glucose uptake, showcasing its anti-hyperglycemic potential [32].

## **2. Anticancer properties**

Pearl millet has been investigated for its anticancer properties. Studies have shown that polyphenols extracted from pearl millet exhibit cytotoxic effects against HepG2 hepatic cancer cell lines, suggesting it could be a natural source of anticancer agents. Additionally, cysteine protease inhibitor (CPI) in pearl millet has demonstrated anti-fungal activity, indicating potential applications in disease prevention, including cancer[33]. Furthermore, pearl millet contains antioxidants and certain compounds (ellagic acid, gallic acid, and quercetin) which have shown anticancer effects by inhibiting matrix metalloproteinases (MMPs) and protein kinase C (PKC)[34]. Bioactive components like ferulic acid and p-coumaric acid have been found to enhance the sensitivity of drug-resistant colorectal cancer cells to chemotherapy drugs like oxaliplatin by modulating ganglioside metabolism and reducing multidrug resistance protein 1 (MDR1) expression[35]. Additionally, CPI has exhibited anti-fungal and anti-feedent activities in pearl millet, suggesting its potential in developing transgenic plants resistant to pathogens and pests, offering a novel approach to cancer treatment through genetic modification[36].

## **3. Probiotic properties**

Fermentation processes involving different microorganism combinations led to increased mineral content (calcium and iron) as observed in various studies. Fermented pearl millet showed higher levels of total phenols and oligopeptides, along with increased resistant starch content and prebiotic activity, particularly when fermented with *Saccharomyces cerevisiae* plus *Campanilactobacillus paralimentarius* [37].

## **4C. Biofortification of Pearl Millet**

Biofortification is the process of increasing the concentration of essential vitamins and minerals in food crops through genetic improvement or agronomic practices. Scientists have concentrated on boosting the bioavailability of important micronutrients like iron and zinc, which are frequently lacking in people's diet. The process of biofortification of pearl millet involves various techniques, including breeding (selective breeding, hybrid development, genome mapping, genetic engineering), agronomic practices (crop rotation, soil and fertilizer management), and post-harvest processing (dehulling, milling, thermal

processing, extrusion, soaking and germination)[30]. Recent developments in the field of biofortification have led to the creation of iron- and zinc-enriched pearl millet varieties (Table 3), which have the potential to address issues of micronutrient deficiency in vulnerable populations [38].

**Table 3. List of biofortified pearl millet varieties & hybrids**

Name of the hybrid/ Variety & Year of release	Grain yield (Kg/ha)	Micronutrients (ppm)		Institute developed	Special Character
		Fe	Zn		
NBH 4903 (2017)	4444	70	63	Nuziveedu Seeds (Balwan) Ltd, TS, India.	85 days
PROAGRO 9450 (2019)	3861	71	58	Bayer BioScience Pvt. Ltd., Hyderabad, India.	Early Maturing
Hybrid CO 9 (2011)	3728	75	41	TNAU, Coimbatore, India.	Short duration, resistant to downy mildew
Composite CO 10 (2017)	3526	81	42	TNAU, Coimbatore, India.	High Protein Content, Compact earhead and bold seed
HHB 299 (2006)	3274	73	41	CCS HAU, Jodhpur, India.	Resistant to major pest and diseases
HHB 311 (2013)	3173	83	39	CCS HAU, Jodhpur, India.	High Fe content, compact ear head
AHB 1200 Fe	3170	77	39	NARP,	High Fe content, highly

(2018)				Aurangabad, India	responsive to fertilizers
RHB 234 (2000)	3169	84	41	AICRP on PM, SKNAU Jaipur, India	Resistant to downy mildew
Moti Sakthi (2017)	3023	76	46	JAU, Jam Nagar, India	Resistant to downy mildew
Mahabeej 1005 (2007)	2994	62	37	MSSCL, Akola, India	Tolerance to stress
Gam Sakthi (2002)	2957	72	43	JAU, Jam Nagar, India	Resistant to downy mildew
Pusa 1201 (MH 1849) (2012)	2810	55	48	ICAR-IARI, New Delhi, India	highly resistant to downy mildew, blast and major pests
GHB 1231 (Sawaj Shakti) (2012)	2760	81	41	JAU, Jamnagar, India	Resistant to downy mildew
ABV 04 (2019)	2680	70	63	ANGRAU, Ananthapuram, India	Long panicle length
Phule Mahashakti (DHBH 1211) (2017)	2581	85	37	ICAR-AICRP- PM, MPKV, Dhule, India	High Fe content, compact ear head
VPMH 7 (2021)	2352	67	52	UAS, Dharwad, India	Resistant to downy mildew
Dhanshakti (2006)	2199	81	43	MPKV, Dhule, India	High Fe content

HHB 67 Imp 2  (1990)	2000	55	40	CCSHAU, Hisar, India	Resistant to downy mildew
BHB-1202	1776	47	42	ICAR-AICRP- PM, SKRAU, Bikaner, India	highly resistant to downy mildew, blast and major pests

[39]

#### 4D. Value added products of pearl millet

In the Indian subcontinent, pearl millet is typically consumed in the form of flattened bread or chapathi. Conversely, in the African subcontinent, fermented breads are more commonly used instead of flatbreads. Fura, a perishable food shaped into balls, is prepared by cooking a blend of fermented and non-fermented pearl millet flour with spices. This food is popularly consumed across Nigeria, often paired with yogurt ('nono') or mixed with water to make porridge, varying by region [40]. Porridges are breakfast cereals that can be consumed as either a thin or thick food, depending on their consistency. Dalia, a popular cereal-milk-based porridge in India, exemplifies this dish. The instant dalia mix contains 30% grain mix (40% pearl millet, 50% sorghum, and 10% maize), skim milk powder (40%) and sugar (30%). They found that this mix had superior protein content and lower fatty acid content. Additionally, the product maintained its quality for over 71 weeks when stored at 10°C [41]. As consumers' awareness of health issues grows, the frozen dessert sector is currently looking for dairy fat substitutes. At 1% and 2% concentrations, octenyl succinyl anhydride esterified pearl millet starch has been found to be a useful fat substitute in ice creams [42].

In addition to being used in salted products, pearl millet grains are also utilized in making sweets. For example, small, sweet balls known as Ladoo and Dakuwa, made from roasted pearl millet grain flours, are popular in India and Nigeria, respectively. These sweets could be enhanced by incorporating ingredients such as popped pearl millet grain and malted flour for nutritional benefits [43]. Pearl millet flour also holds promise for creating various globally popular products, including ready-to-eat snacks, weaning foods and non-dairy fermented beverages. Furthermore, it can serve as an alternative to wheat flour in a variety of food preparations such as biscuits, pastas, whole meal breads, and kibbeh.

#### **4E. Antinutritional Factors**

Pearl millet contains antinutritional factors such as phytic acid, tannins, and polyphenols, which can hinder the absorption of essential minerals like iron and zinc [44]. These antinutrients reduce the bioavailability of minerals in millets, impacting their overall nutritional value. Traditional processing methods, including roasting, blanching, and malting, have been shown to significantly decrease the levels of phytic acid in millets, thereby enhancing nutrient absorption [45]. Additionally, pearl millet germplasm exhibits variability in the levels of phytic acid and goitrogenic compounds, highlighting the potential for breeding programs to develop new varieties with improved nutritional quality and reduced antinutrient content [46]. By understanding and addressing these antinutritional factors, the nutritional value of pearl millet can be enhanced, making it a more beneficial food source for populations in developing countries where deficiencies in iron and zinc are prevalent.

#### **5.Way forward:**

Improving pearl millet for climate resilience and nutritional value involves a multi-faceted approach that includes breeding for desirable traits, effective soil and pest management, and innovative processing techniques to enhance nutrient availability. Development of resistant hybrids/varieties resistant/tolerant. Shift in focus of breeding from productivity improvement to the identification of stress specific traits. Development of efficient screening protocols and control measures against different diseases and pests of pearl millet. Biofortification in pearl millet for minor nutrients. Prolonged shelf life of pearl millet flour and overcome rancidity. This not only supports food security but also addresses nutritional deficiencies in vulnerable populations

#### **6. CONCLUSION:**

Pearl millet emerges not only as a resilient crop in the face of climate change but also as a vital contributor to global food security and nutrition. Its ability to thrive in harsh environments, characterized by heat and drought, makes it a dependable option for regions susceptible to erratic weather patterns. Furthermore, its nutritional profile, rich in essential minerals, proteins, and gluten-free properties, positions it as a valuable resource in combating malnutrition and addressing dietary deficiencies worldwide.

As we confront the challenges of a changing climate and a growing global population, promoting the cultivation and consumption of pearl millet offers a sustainable solution.

Continued research and investment in pearl millet production not only bolster food security but also contribute to the resilience of agricultural systems in the face of increasingly unpredictable environmental conditions. Embracing pearl millet represents a forward-thinking approach to ensuring food security and promoting nutritional diversity on a global scale.

### **Disclaimer (Artificial intelligence)**

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

### **REFERENCES:**

1. Wheeler T, Von Braun J. Climate change impacts on global food security. *Science*. 2013 Aug 2;341(6145):508-13.
2. Antony Ceasar S, Maharajan T. The role of millets in attaining United Nation's sustainable developmental goals. *Plants, People, Planet*. 2022 Jul;4(4):345-9.
3. Singh RP, Qidwai S, Singh O, Reddy BR, Saharan S, Kataria SK, Tiwari H, Naresh RK, Kumar L. Millets for food and nutritional security in the context of climate resilient agriculture: A Review. *International Journal of Plant & Soil Science*. 2022 Oct 31:939-53.
4. Kheya SA, Talukder SK, Datta P, Yeasmin S, Rashid MH, Hasan AK, Anwar MP, Islam AA, Islam AM. Millets: The future crops for the tropics-Status, challenges and future prospects. *Heliyon*. 2023 Nov 10.
5. Chakraborti R, Davis KF, DeFries R, Rao ND, Joseph J, Ghosh S. Crop switching for water sustainability in India's food bowl yields co-benefits for food security and farmers' profits. *Nature Water*. 2023 Oct;1(10):864-78.
6. Fatondji D, Ajeigbe HA, Ibrahim A. Modern Crop Management Practices for Pearl Millet Cultivation in Semi-Arid Africa. *Pearl Millet in the 21st Century: Food-Nutrition-Climate resilience-Improved livelihoods*. 2024 Feb 10:445-77.
7. Gupta SK, Rai KN, Singh P, Ameta VL, Gupta SK, Jayalekha AK, Mahala RS, Pareek S, Swami ML, Verma YS. Seed set variability under high temperatures during flowering period in pearl millet (*Pennisetum glaucum* L.(R.) Br.). *Field Crops Research*. 2015 Feb 1;171:41-53.
8. Dave HR. Pearl millet hybrids. In *Proceedings of the international pearl millet workshop 1986* Apr 7 (pp. 7-11).
9. Bidinger FR, Nepolean T, Hash CT, Yadav RS, Howarth CJ. Quantitative trait loci for grain yield in pearl millet under variable postflowering moisture conditions. *Crop science*. 2007 May;47(3):969-80.
10. Shrestha N, Hu H, Shrestha K, Doust AN. Pearl millet response to drought: A review. *Frontiers in Plant Science*. 2023 Feb 10;14:1059574.
11. Rostamza M, Richards RA, Watt M. Response of millet and sorghum to a varying water supply around the primary and nodal roots. *Annals of Botany*. 2013 Jul 1;112(2):439-46.

12. Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarc'h S, Ortega BM, Atkinson JA, Belko MN, Bennett MJ, Gantet P, Wells DM. Characterization of pearl millet root architecture and anatomy reveals three types of lateral roots. *Frontiers in plant science*. 2016 Jun 13;7:829.
13. Pardo J, VanBuren R. Evolutionary innovations driving abiotic stress tolerance in C4 grasses and cereals. *The Plant Cell*. 2021 Nov 1;33(11):3391-401.
14. Shivhare R, Lata C. Exploration of genetic and genomic resources for abiotic and biotic stress tolerance in pearl millet. *Frontiers in plant science*. 2017 Jan 23;7:2069.
15. Shivhare R, Lata C. Assessment of pearl millet genotypes for drought stress tolerance at early and late seedling stages. *Acta Physiologiae Plantarum*. 2019 Mar;41(3):39.
16. Sun M, Huang D, Zhang A, Khan I, Yan H, Wang X, Zhang X, Zhang J, Huang L. Transcriptome analysis of heat stress and drought stress in pearl millet based on Pacbio full-length transcriptome sequencing. *BMC Plant Biology*. 2020 Dec;20:1-5.
17. Bi H, Kovalchuk N, Langridge P, Tricker PJ, Lopato S, Borisjuk N. The impact of drought on wheat leaf cuticle properties. *BMC Plant Biology*. 2017 Dec;17:1-3.
18. Varshney RK, Shi C, Thudi M, Mariac C, Wallace J, Qi P, Zhang H, Zhao Y, Wang X, Rathore A, Srivastava RK. Pearl millet genome sequence provides a resource to improve agronomic traits in arid environments. *Nature biotechnology*. 2017 Oct;35(10):969-76.
19. Shivhare R, Asif MH, Lata C. Comparative transcriptome analysis reveals the genes and pathways involved in terminal drought tolerance in pearl millet. *Plant molecular biology*. 2020 Aug;103(6):639-52.
20. Ashraf M, Hafeez M. Thermotolerance of pearl millet and maize at early growth stages: growth and nutrient relations. *Biologia plantarum*. 2004 Mar;48:81-6.
21. Mukesh Sankar S, Tara Satyavathi C, Barthakur S, Singh SP, Bharadwaj C, Soumya SL. Differential modulation of heat-inducible genes across diverse genotypes and molecular cloning of a sHSP from pearl millet [*Pennisetum glaucum* (L.) R. Br.]. *Frontiers in Plant Science*. 2021 Jul 16;12:659893.
22. Swarna R, Jinu J, Dheeraj C, Talwar HS. Salinity Stress in Pearl Millet: From Physiological to Molecular Responses. In *Pearl Millet in the 21st Century: Food-Nutrition-Climate resilience-Improved livelihoods 2024* Feb 10 (pp. 361-394). Singapore: Springer Nature Singapore.
23. Singh J, Reddy PS, Reddy CS, Reddy MK. Molecular cloning and characterization of salt inducible dehydrin gene from the C4 plant *Pennisetum glaucum*. *Plant Gene*. 2015 Dec 1;4:55-63.
24. Sharma R, Sharma S, Gate VL. Tapping *Pennisetum violaceum*, a wild relative of pearl millet (*Pennisetum glaucum*), for resistance to blast (caused by *Magnaporthe grisea*) and rust (caused by *Puccinia substriata* var. *indica*). *Plant disease*. 2020 May 16;104(5):1487-91.
25. Kumar A, Manga VK, Gour HN, Purohit AK. Pearl millet downy mildew: challenges and prospects. *Ann Rev Plant Pathol*. 2013 Jul 1;5:139.
26. Sehgal D, Skot L, Singh R, Srivastava RK, Das SP, Taunk J, Sharma PC, Pal R, Raj B, Hash CT, Yadav RS. Exploring potential of pearl millet germplasm association panel for association mapping of drought tolerance traits. *PLoS One*. 2015 May 13;10(5):e0122165.
27. Tara Satyavathi C, Ambawat S, Singh S, Lata C, Tiwari S, Siddaiah CN. Genomic designing for biotic stress resistance in pearl millet [*Pennisetum glaucum* (L.) R. Br.]. *Genomic designing for biotic stress resistant cereal crops*. 2021:257-94.
28. Kulkarni, K.S., Zala, H.N., Bosamia, T.C., Shukla, Y.M., Kumar, S., Fougat, R.S., Patel, M.S., Narayanan, S. and Joshi, C.G., 2016. De novo transcriptome sequencing

- to dissect candidate genes associated with pearl millet-downy mildew (*Sclerospora graminicola* Sacc.) interaction. *Frontiers in Plant Science*, 7, p.847.
29. Mahfooz Ali G. Nutritional Composition and Technological Studies on Pearl Millet Grains (*Pennisetum glaucum* L.). Alexandria Science Exchange Journal. 2023 Jun 30;44(2):147-59.
  30. Boncompagni E, Orozco-Arroyo G, Cominelli E, Gangashetty PI, Grando S, Kwaku Zu TT, Daminati MG, Nielsen E, Sparvoli F. Antinutritional factors in pearl millet grains: Phytate and goitrogens content variability and molecular characterization of genes involved in their pathways. *PloS one*. 2018 Jun 1;13(6):e0198394.
  31. Ironi EA, Imam YT, Ajani EO. Physicochemical, antioxidant and starch-digesting enzymes inhibitory properties of pearl millet and sweet detar gluten-free flour blends, and sensory qualities of their breads. *Frontiers in Food Science and Technology*. 2022 Sep 20;2:974588.
  32. Krishnan V, Verma P, Saha S, Singh B, Vinutha T, Kumar RR, Kulshreshta A, Singh SP, Sathyavathi T, Sachdev A, Praveen S. Polyphenol-enriched extract from pearl millet (*Pennisetum glaucum*) inhibits key enzymes involved in post prandial hyperglycemia ( $\alpha$ -amylase,  $\alpha$ -glucosidase) and regulates hepatic glucose uptake. *Biocatalysis and Agricultural Biotechnology*. 2022 Aug 1;43:102411.
  33. Singh N, Meenu G, Sekhar A, Abraham J. Evaluation of antimicrobial and anticancer properties of finger millet (*Eleusine coracana*) and pearl millet (*Pennisetum glaucum*) extracts. *The Pharma Innovation*. 2016 Mar 1;3(11, Part B).
  34. Rathore S, Verma S, Singh K, Pandey AK. An in silico approach for the evaluation of nutraceutical properties of pearl millet as a brain food. *J In Silico In Vitro Pharmacol*. 2017;3(2):18.
  35. Zhang X, Shan S, Shi J, Li H, Li Z. Polyphenol from millet bran increases the sensitivity of colorectal cancer cells to oxaliplatin by blocking the ganglioside GM3 catabolism. *Food & function*. 2021;12(1):291-301.
  36. Joshi BN, Sainani MN, Bastawade KB, Deshpande VV, Gupta VS, Ranjekar PK. Pearl millet cysteine protease inhibitor: Evidence for the presence of two distinct sites responsible for anti-fungal and anti-feedent activities. *European journal of biochemistry*. 1999 Oct;265(2):556-63.
  37. Balli D, Cecchi L, Pieraccini G, Venturi M, Galli V, Reggio M, Di Gioia D, Furlanetto S, Orlandini S, Innocenti M, Mulinacci N. Millet Fermented by Different Combinations of Yeasts and Lactobacilli: Effects on Phenolic Composition, Starch, Mineral Content and Prebiotic Activity. *Foods*. 2023 Feb 8;12(4):748.
  38. Wrigley CW, Corke H, Seetharaman K, Faubion J, editors. *Encyclopedia of food grains*. Academic Press; 2015 Dec 17.
  39. Satyavathi CT, Khandelwal V, Rajpurohit BS, Supriya A, Beniwal BR, Kamlesh K, Sushila B, Shripal S, Mahesh CK, Yadav SL. Pearl millet-hybrids and varieties. ICAR-All India Coordinated Project on Pearl millet, Mandor, Jodhpur India. 2018;142.
  40. Filli KB, Nkama I, Jideani VA, Ibok IU. Effect of process variables on the hydration properties and acceptability of extruded millet-soybean blends for fura manufacture. *British Food Journal*. 2013 Jun 7;115(6):884-98.
  41. Singh BP, Jha A, Sharma N, Rasane P. Optimization of a process and development of a shelf life prediction model for instant multigrain dalia mix. *Journal of Food Process Engineering*. 2013 Dec;36(6):811-23.
  42. Hassan ZM, Sebola NA, Mabelebele M. The nutritional use of millet grain for food and feed: a review. *Agriculture & food security*. 2021 Dec;10:1-4.

43. Singh G, Sehgal S. Nutritional evaluation of ladoo prepared from popped pearl millet. *Nutrition & Food Science*. 2008 Jul 18;38(4):310-5.
44. Rani S, Singh R, Sehrawat R, Kaur BP, Upadhyay A. Pearl millet processing: a review. *Nutrition & Food Science*. 2018 Feb 12;48(1):30-44.
45. Singhal T, Tara Satyavathi C, Singh SP, Mallik M, Anuradha N, Sankar SM, Bharadwaj C, Singh N. Achieving nutritional security in India through iron and zinc biofortification in pearl millet (*Pennisetum glaucum* (L.) R. Br.). *Physiology and Molecular Biology of Plants*. 2022 Apr;28(4):849-69.
46. Sheethal HV, Baruah C, Subhash K, Ananthan R, Longvah T. Insights of nutritional and anti-nutritional retention in traditionally processed millets. *Frontiers in Sustainable Food Systems*. 2022 Feb 1;5:735356.