

Original Research Article

DETERMINATION OF OIL CONTENT AND FATTY ACID COMPOSITION OF TEA PLANTS AT PLANTATIONS IN THE BLACK SEA REGION

.ABSTRACT

Aims: The aim of this study was to determine the fatty acid composition and oil content of *C. sinensis* seeds collected from different locations at the Northern Black Sea region.

Study design: The study based on the collection of tea seed samples from different locations at the Northern Black Sea Region in Türkiye and comparing their fatty acid composition and oil content of analyzed samples.

Place and Duration of Study: Recep Tayyip Erdogan University, Faculty of Agronomy, Field Crops Department, Rize, Türkiye, between 2022 and July 2023.

Methodology: A total of 54 samples were taken within the scope of determining oil content and fatty acid composition of *Camellia sinensis* seeds collected from different locations. The location were chosen from different cities concerning altitude. The number of samples taken from the province of Rize is 37, the number of samples taken from the province of Trabzon is 7, the number of samples taken from the province of Giresun is 5, and the number of samples taken from the province of Artvin is 5. Fatty Acid composition was analyzed using GC-MS and oil content with an automated oil extraction device.

Results: Oil ratio and fatty acid composition were determined in these samples. As stated in the literature, oleic acid, which is the criterion of vegetable oil quality, was found to be high in all samples. With Principal Component Analysis, the oil content and fatty acid composition data of the samples were used to reveal the differences in the samples. The obtained data will shed light on future studies.

Conclusion: Knowing the fatty acid composition of tea oils will enable production of oils according to their intended use. For this purpose, it will be possible to produce oils suitable for the purpose by growing the desired types in suitable regions. We have to consider, that seeds were obtained after leaf harvest of tea plants. We can assume that the seed yield will increase if the tea plants are grown only for seed production. With governmental support tea seed oil production will be an potential additional income for local farmers.

Keywords: Camellia sinensis, oil content, fatty acid composition, Türkiye

1. INTRODUCTION

Vegetable oils represent important raw materials used in the chemical industry. The rising prices of petroleum resources, combined with their finite and non-renewable nature, have renewed researcher interest in the development and use of alternative renewable feedstocks. Vegetable oils are an alternative energy source to petrochemicals because they are renewable, biodegradable, less toxic and environmentally friendly. [1]

Tea (*Camellia sinensis* O. Kuntze) is a plant that was brought to Türkiye in the 1930s with the work of Ali Rıza Erten and Zihni Derin and quickly became an important and economically profitable plant. While we were a coffee-consuming society in the past, tea has become the most consumed beverage in Türkiye due to its suitability to our taste and the unique brewing method of Türkiye.

In Türkiye, tea is mostly known for its green leaves being collected in 2-3 harvest periods a year in order to produce green and black tea. The fact that tea seeds contain a significant amount of oil is not well known by our society.

The tea industry in Türkiye is an important sector for the regional and national economy. Although tea can grow in a limited area in the Eastern Black Sea region of our country, it is important as a beverage that we consume at every moment of our daily lives. In addition, the lack of opportunities for growing various agricultural products in the tea-growing region has increased the economic value of tea for the local people. In addition, the fact that tea cannot be grown in other regions due to the necessary climatic conditions has increased the importance of the tea region. In this context, tea is in a monopoly

position due to the lack of an alternative in the region where it is grown and the lack of possibility to grow it in other regions. [2]

Vegetable oil from oilseed plants has always been used for nutrition and is increasingly used as an industrial raw material today. Most oilseed plants characteristically accumulate oil in their seeds (rapeseed, sunflower) or fruits (palm, olive). Most of these oilseed crops grow in temperate climates, they only need adequate amounts of fertilizer and high yields can be achieved. [3]

Tea seed oil is obtained from the seeds of various tea species belonging to the Theaceae family. [4] In terms of fatty acid composition, tea seed oil is similar to quality cooking oil and olive oil. [5-6] In addition to being quite expensive for general use, olive oil is not widely consumed because there are not enough resources to meet global demand.

According to recent research, tea seed oil is emerging as a new alternative to canola oil and olive oil due to its high oleic acid, medium linoleic acid and low linolenic acid content. [6] Rich in antioxidants, tea seed oil has been shown to lower blood pressure and cholesterol levels. [7]

It has been reported that the oil content of tea seed is approximately 30-32% and has high organoleptic acceptability. [6,8] Tea seed oil is a high-quality cooking oil like olive oil and can be stored well at room temperature. It is known that tea seed oil reduces blood pressure and cholesterol levels, has high antioxidant content, is a rich source of emollients for skin care and minimizes the signs of aging. [7]

The fatty acid composition of *C. sinensis* seed oil consists of 21.5% palmitic acid, 2.9% stearic acid [9], 56% oleic acid, 22% linoleic acid and 0.3% linolenic acid. [6] The main fatty acid in *C. sinensis* seed oil (50% of the total oil) is oleic acid. [10] Therefore, in terms of oleic acid, *C. sinensis* seed oil can be ranked between sunflower and olive oil. [6]

The oil content and fatty acid composition of tea seeds collected from different villages in the Black Sea region were investigated regarding its oil content and fatty acid composition.

2. MATERIAL AND METHODS

2.1. Sample Collection

Tea seeds were collected from tea producing areas to examine the oil ratios and fatty acid composition. Seeds were collected at the appropriate time by planning according to the provinces where tea is grown and by going out to the field in sufficient numbers and from different areas in each province. In the collection of tea seeds, the most samples were collected from Rize province, which is the province where tea is grown most intensively.

Within the scope of the plan, tea seeds were collected from the field by determining the city, district, altitude, coordinates, garden owner and village or neighborhood name and accompanied by label information. was. The collected tea seeds were placed in cold storage at the end of each day to prevent possible changes in the oil content and fatty acid composition.

Information about the samples taken in the study is given in Table 1.

Table 1: Location names of collected tea samples from different cities

City	District	Village	Coordinates	Altitude	Location name
Rize	Fındıklı	Sümer	41°17'05.64" 41°14'34.64"	90	Sümer KöyüYazıcılar Mah. No.5
Rize	Fındıklı	Derbent	41°17'41.06" 41°15'49.69"	249	ArmoınMah.Sela zerGüzergahı
Rize	Fındıklı	Derbent- SelazurMevki	41°17'04.04" 41°15'54.91"	342	SelazurMah.Han e No.85
Rize	Fındıklı	Beydere	41°14'41.00" 41°13'14.00"	343	KarşıyakaMevkii

Rize	Fındıklı	Aslandere	41°14'35.19" 41°15'23.89"	420	ZiyunaMevki
Trabzon	Sürmene	Gültepe	40°52'23.81" 40°03'53.00"	120	Küçükdere Yol Üzeri
Trabzon	Sürmene	Güney Köyü	40°46'07.58" 40°03'30.22"	422	Küçükdere Oylum yoluüzeriGüneyk öy Oğuz CamiiYaoluTabel ası
Trabzon	Sürmene	Yeşilköy	40°47'28.41" 40°02'43.31"	790	KüçükdereYeşilk öyYolu
Giresun	Eynesil	Altınlı	41°04'10.42" 39°09'06.30"	7	Boztepe MahallesiMezarlı kYanı Mustafa Yüksel Cad.
Giresun	Eynesil	Boztepe Mah.	41°03'52.90" 39°10'14.26"	225	Yukarı Boztepe Camiiönü
Trabzon	Beşikdüzü	Şahmelik	41°00'12.19" 39°10'50.11"	585	Zalahnamevkii
Giresun	Tirebolu	Demircili Mah.	41°00'19.07" 38°51'27.93"	46	GüldalMah.Yolüz eryokuş
Giresun	Tirebolu	Karaahmetli	40°56'39.57" 38°52'54.69"	285	Karaahmetli Yol kenarı
Giresun	Tirebolu	Kovapınar	40°55'41.00" 38°55'02.98"	424	Kovapınarköyiçiy olçıkış
Artvin	Borçka	Muratlı	41°27'24.27" 41°42'45.62"	98	Muratlı Çay Fabrikasıönü
Artvin	Borçka	Güreşen	41°27'10.82" 41°38'48.28"	235	Güreşenmerkez amiimevkii
Rize	Hemşin	Ortaköy	41°02'15.58" 40°54'00.38"	347	Organik Çay fabrikasıkarşısı
Rize	Hemşin	Bilenköy	41°01'34.73" 40°54'15.88"	637	Bilenköyeski konak yanı
Rize	Hemşin	Nurluca	41°02'05.47" 40°54'02.85"	493	Serkümeevleri
Rize	Pazar	Yemişli	41°09'07.00" 40°55'29.00"	49	YemişliMevki
Rize	Pazar	Orta Irmak Köyü	41°07'50.00" 40°56'40.00"	247	Yüksel Mevkii

Rize	Pazar	AkbucakKöyü	41°05'24.00" 40°57'38.00"	525	YukarıMahalle
Trabzon	Of	Cumapazarı	40°50'34.16" 40°14'55.08"	528	KonofolMevkii
Trabzon	Of	Cumapazarı	40°50'07.16" 40°15'54.44"	265	Renk Mahallesi
Trabzon	Of	Balıca	40°52'53.16" 40°16'59.81"	76	
Artvin	Hopa	Koyuncular	41°23'45.73" 41°30'38.52"	506	ÇalinMevki
Artvin	Hopa	Koyuncular	41°24'08.50" 41°30'10.08"	252	ZalunaMevkii
Artvin	Hopa	Yoldere	41°23'28.25" 41°28'07.72"	69	OkulMevkii
Rize	Ardeşen	Köprüköy	41°07'18.75" 41°03'06.79"	513	KalaizeniMevki
Rize	Ardeşen	Tunca Esentepe	41°07'42.87" 41°07'00.32"	622	EsentepeMevkii
Rize	Ardeşen	Köprüköy	41°07'29.64" 41°02'34.76"	253	TaşköprüMevkii
Rize	Merkez	Hayrat	41°01'22.27" 40°30'06.97"	64	OkulCivarı
Rize	Merkez	Ortapazar	40°59'01.53" 40°28'34.96"	310	AşağıMahalle
Rize	Merkez	Ortapazar	40°57'52.56" 40°28'37.26"	520	Hudut Kaya Mevkii
Rize	Kalkandere	Fındıklı	40°54'21.60" 40°28'12.1"	237	FındıklıKöyüyolu MusluoğluMevki
Rize	İyidere	Köşklü	40°59'2.95" 40°21'32.91"	230	Köşklü- KambiyozMevki
Rize	İyidere	Yapraklar	41° 0'5.18" 40°22'49.01"	185	YapraklarMahall eiçiyol Arka Mevki
Rize	İkizdere	Ilıca	40°47'26.02" 40°35'32.89"	965	IlıcaKöyü Cami Mevkii
Rize	İkizdere	Gürdere	40°48'49.34" 40°33'14.88"	875	SırtlarMahalleMe vkii
Rize	İkizdere	Ihlamur	40°47'52.30" 40°32'7.38"	800	IhlamurKöyü Orta Mahalle

Rize	Kalkandere	Aşağı Tatlısu	40°55'42.70" 40°25'15.88"	150	Güneysu Mevkii
Rize	Kalkandere	Yolbaşı	40°57'8.30" 40°28'5.70"	449	Yolbaşı Köyü Okul Mevkii
Rize	Kalkandere	Ormanlı	40°56'53.90" 40°24'60.00"	153	Ormanlı Köyü Karana Mevkii
Rize	Kalkandere	Çağlayan	40°55'27.30" 40°28'4.90"	305	Çağlayan Köyü Sağlamlar Mevkii
Rize	Güneysu	Ulucami	40°59'23.18" 40°36'24.34"	128	Ulucamii Dere Mevkii
Rize	Güneysu	Yüksekköy	40°56'38.86" 40°34'22.79"	791	Yüksekköy Üst Mahalle
Rize	Güneysu	Dumankaya	40°58'10.62" 40°35'52.73"	426	Dumankaya Köyü Köyiçi Mevkii
Rize	Çayeli	Derecik	41° 0'3.72" 40°40'58.16"	295	Derecik Köyü Cami Mevkii
Rize	Çayeli	Yaka	41° 4'50.32" 40°42'15.44"	19	Yaka Mahallesi Yalı Civarı
Rize	Çayeli	Yalıköy	41° 5'33.68" 40°44'7.43"	45	Yalıköy Dere Mevkii
Rize	Çayeli	Eskipazar	41° 4'59.71" 40°42'59.77"	63	Sofogon Hozan
Rize	Derepazarı	Bahattinpaşa	41° 1'0.87" 40°26'44.90"	242	Aşağı Mahalle
Rize	Derepazarı	Kirazdağı	40°58'59.50" 40°24'27.60"	295	Baş Mahalle
Rize	Derepazarı	Bürücek	41° 1'37.11" 40°26'39.88"	138	Ortamahalle

The total number of samples taken within the scope of the thesis is 54. The number of samples taken from Rize province is 37, the number of samples taken from Trabzon province is 7, the number of samples taken from Giresun province is 5 and the number of samples taken from Artvin province is 5. The lowest altitude sample was collected from Giresun province Eynesil İlçesi Boztepe District (7 m) and the highest altitude sample was collected from Rize province İkizderelliçca Village (965 m).

2.2. Determination of Oil Content

After the sample collection was completed, the green outer shell parts of the tea seeds were separated and dried. It was dried at 40° C for 7 days, and the sample weights before and after drying were measured and recorded with a precision scale. Dried samples were ground with their inner shells in coffee grinder-like machines and brought to the appropriate particle size according to the extraction method for the detection of oil in tea seeds.

Fully automatic (Buchi) Soxhlet Extraction device was used to determine the tea seed oil content by extraction method. First of all, 5 g of the ground samples, which were kept in a ziplock bag with a label

number, were weighed with a precision balance and placed into the cartridges to be used for extraction. On the ground sample in each cartridge; Overflow, spillage, flying etc. that may occur during the extraction process. A small amount of pure cotton was added to eliminate drawbacks such as.

Since the fully automatic extraction device processes 6 samples in one filling, 6 cartridges; They were placed into the glass tubes where the process will be carried out, respectively, by fixing them with the help of a lid. In order for the system to work, the beakers in which the liquid resulting from the extraction will be collected are placed in their places and made ready for operation.

100 ml of Diethyl Ether, which will be used in the extraction process, was measured with a graduated measuring tape and added to the samples placed in the device with the help of a funnel. After the ground samples were processed in the system for 120 minutes at 40°C, the solution accumulated in the previously weighed beakers was taken with the help of tongs and taken to the oven with the temperature set at 40°C. It was dried in the oven at 40° C for 24 hours to allow the diethyl ether in it to evaporate. The next day, the beakers were taken from the oven again with tongs and cooled in a desiccator. The cooled samples were weighed again on a precision scale and the amount of oil was determined according to the formula below.

$$\text{Oil Content (\%)} = \frac{\text{Last weight (gr)} - \text{First Weight (gr)}}{5} \times 100$$

After weighing, the remaining oils in the beaker were removed with the help of a pipette and transferred to Eppendorf tubes. The oils in closed Eppendorf tubes were covered with stretch film and stored in the refrigerator at +4° C.

2.3. Determination of Fatty Acid Composition

A 1 molar Potassium Hydroxide (KOH) solution was prepared to determine fatty acid composition. 0.1 g of the oils in the Eppendorf tubes were weighed and dissolved with 10 ml hexane in a 10 ml falcon tube. 0.5 ml (500 microliters) KOH was added to the dissolved oil and shaken. The mixture was left in a dark environment for 45 minutes. Then it was kept in the refrigerator at +4 degrees for 30 minutes. The solution taken from the upper phase accumulated on the surface with the help of an automatic pipette was transferred to 2 ml vials and fatty acid compositions were determined by Gas Chromatography (GCMS).

A SHIMADZU GC-2010 Plus device attached a TRCN-100 column (100m x 0.25 mm x 0.20 µm) was used for fatty acid analysis. The injection temperature was 250 °C with a pressure of 250 kPa. The column temperature was increased from 140°C to 240°C. FID temperature was 250°C and injection volume was 1 µl. Split ratio was used as 1/100.

2.4. Data Analysis

Biplot Analysis were performed using the XLSTAT 2023 Statistical Program to visualize present variation in *S. tomentosa* plant parts investigated for chemical variability. Scatter plot diagrams were created using current data. [11] Based on oil content and fatty acid composition data Biplot diagram was created.

3. RESULTS AND DISCUSSION

3.1. Oil Content

Oil content values obtained from tea samples collected from Artvin, Giresun, Trabzon and Rize provinces/districts and villages are given in Table 1. Generally speaking, seed samples were collected from Altınlı village of Eynesil/Giresun, at an altitude of 7 m, and from İkizdere/RizeGüredere village, at an altitude of 875 m. 37 samples representing all districts of Rize province and 17 samples from other provinces.

The oil content of the samples collected from Artvin province was between 15.2 - 22.6%, in Giresun province between 14.2-19.4%, in Rize province between 8.4-21.8% and in Trabzon province between 15.8 - 17.8%. Significant differences were detected between provinces and in terms of oil content.

Table 2: Oil content of collected tea samples

Sample codes	Province	District	Village	Oil Content
ÇT16	Artvin	Hopa	Yoldere	19.4
ÇT15	Artvin	Borçka	Muratlı	18.2
ÇT26	Artvin	Borçka	Güreşen	15.2
ÇT28	Artvin	Hopa	Koyuncular	22.6
ÇT27	Artvin	Hopa	Koyuncular	21.2
ÇT9	Giresun	Eynesil	Altınlı	16.4
ÇT12	Giresun	Tirebolu	Demircili Mah.	19.4
ÇT10	Giresun	Eynesil	Boztepe Mah.	14.2
ÇT13	Giresun	Tirebolu	Karaahmetli	19.0
ÇT14	Giresun	Tirebolu	Kovapınar	14.4
ÇT49	Rize	Çayeli	Yaka	18.8
ÇT50	Rize	Çayeli	Yalıköy	20.6
ÇT20	Rize	Pazar	Yemişli	15.2
ÇT51	Rize	Çayeli	Eskipazar	20.6
ÇT32	Rize	Merkez	Hayrat	18.0
ÇT1	Rize	Fındıklı	Sümer	15.6
ÇT45	Rize	Güneysu	Ulucami	21.2
ÇT54	Rize	Derepazarı	Bürücek	18.8
ÇT41	Rize	Kalkandere	Aşağı Tatlısu	8.4
ÇT43	Rize	Kalkandere	Ormanlı	12.6
ÇT37	Rize	İyidere	Yapraklar	21.8
ÇT36	Rize	İyidere	Köşklü	26.0
ÇT35	Rize	Kalkandere	Fındıklı	16.4
ÇT52	Rize	Derepazarı	Bahattinpaşa	17.2
ÇT21	Rize	Pazar	Orta Irmak Köyü	16.4

ÇT2	Rize	Fındıklı	Derbent	15.4
ÇT31	Rize	Ardeşen	Köprüköy	15.8
ÇT48	Rize	Çayeli	Derecik	20.0
ÇT53	Rize	Derepazarı	Kirazdağı	17.8
ÇT44	Rize	Kalkandere	Çağlayan	17.4
ÇT33	Rize	Merkez	Ortapazar	16.6
ÇT3	Rize	Fındıklı	Derbent- SelazurMevki	17.6
ÇT4	Rize	Fındıklı	Beydere	15.6
ÇT17	Rize	Hemşin	Ortaköy	15.8
ÇT5	Rize	Fındıklı	Aslandere	16.2
ÇT47	Rize	Güneysu	Dumankaya	16.6
ÇT42	Rize	Kalkandere	Yolbaşı	18.6
ÇT19	Rize	Hemşin	Nurluca	16.6
ÇT29	Rize	Ardeşen	Köprüköy	19.2
ÇT34	Rize	Merkez	Ortapazar	15.4
ÇT22	Rize	Pazar	AkbucakKöyü	14.4
ÇT30	Rize	Ardeşen	Tunca Esentepe	20.4
ÇT18	Rize	Hemşin	Bilenköy	16.0
ÇT46	Rize	Güneysu	Yüksekköy	14.0
ÇT40	Rize	İkizdere	Ihlamur	15.4
ÇT39	Rize	İkizdere	Gürdere	19.0
ÇT38	Rize	İkizdere	Ilıca	17.6
ÇT25	Trabzon	Of	Balıca	18.0
ÇT6	Trabzon	Sürmene	Gültepe	17.8
ÇT24	Trabzon	Of	Cumapazarı	17.8
ÇT7	Trabzon	Sürmene	Güney Köyü	17.2

ÇT23	Trabzon	Of	Cumapazarı	16.6
ÇT11	Trabzon	Beşikdüzü	Şahmelik	15.8
ÇT8	Trabzon	Sürmene	Yeşilköy	16.0

In addition to the leaves of the tea (*Camellia sinensis* L.) plant (raw materials such as green tea, black tea, oolong tea, etc.), the seeds should also be considered as a product of the plant. Recently, with the widespread use of tea, the yield of tea seeds has also increased. For example, over one million tons of tea seeds are produced in China every year. [12]

The huge increase in tea seed production increases the necessity to find suitable commercial practices. Like other *Camellia* species, tea seeds, which are from the Theaceae family, are rich in oil (30-32%) [13] and are considered a high-quality edible oil type because they contain high amounts of monounsaturated fatty. [6] Therefore, it is essential to use tea seeds as a source of edible oil, otherwise it will only be used in the establishment of new tea fields.

It is known that *Camellia* genus plant seeds have high oil content, on average around 30% oil per seed, but the oil content varies between 24% and 50% depending on the species, variety and environmental conditions. [4, 6, 13-16] High seed oil content is probably the result of a number of factors, but environmental variables such as soil, altitude, light, precipitation, humidity and temperature also play a key role, and these factors have been shown to influence the oil content in the seeds of many plants. [17-20]

The oil extracted from the seeds of *Camellia* species both cultivated as well as other species is termed as tea seed oil. Though *C. sinensis* is cultivated mostly for producing tea of commerce, oil is not usually obtained from this species. Commercial production of oil is derived from species like *C. sasanqua*, *C. japonica*, *C. tenuifolia* and *C. oleifera*. [21] *C. sinensis* seeds contains 20-30 % oil in its seeds. [22]

The fat ratios obtained in this study were lower than the fat ratio values given for the *Camellia sinensis* species in the literature. One of the main reasons for this is the collection of tea seeds in Türkiye, especially after the 2nd or 3rd shoot. If certain tea farming areas are reserved for tea seed production and direct tea seed production is started without harvesting green leaves, it will be possible to increase both tea seed yield and oil rate in tea areas in Türkiye.

On the other hand, the way to be successful in growing oil *Camellia* species in different parts of the world is to select promising clones with high oil content and quality for the targeted regions. [23]

Camellia plant, which is used to obtain ornamental plants and tea, is one of the four main tree-form plants (palm tree, coconut, olive oil and tea plant) containing oil. [24] Oil obtained from *Camellia* species has become important in China and to a lesser extent in Japan.

In China, tea seed oil was also traditionally used as a medication for burn injuries and stomachaches. Rats' serum levels of low density lipoprotein, triglycerides, and cholesterol may all be lowered by the saponin in tea seeds. [25-26] Tea seed oil is said to offer several health benefits, including lowering blood pressure and cholesterol, being rich in emollients for skin care, having a high antioxidant content (polyphenols, carotenoids, and vitamin E), and minimising indications of ageing [27-28]

In recent years, increasing attention has been drawn to the health benefits of olive oil, which has led to a dramatic increase in olive oil production outside non-traditional production regions outside the Mediterranean. Similarly, *camellia* oil now has a healthy image and it is possible to say that there will be a similar increase in its use.

3.2. Fatty acid composition

The fatty acid composition of tea seed samples collected from Artvin, Giresun, Trabzon and Rize provinces is given in the Table 3.

In tea seed samples, palmitic acid content varied between 10.31-20.74%, stearic acid content 1.18-3.52%, linoleic acid content 18.98-26.36% and linolenic acid content 0-1.23%. As expected, oleic acid, an indicator of vegetable oil quality, was high in all samples. The oleic acid content in the samples collected from Artvin varied between 53.87-61.04%, 48.35-60.77% in Giresun, 34.09-62.41 in Rize and 49.94-61.28% in Trabzon.

UNDER PEER REVIEW

Table 3: Fatty acid composition of collected tea samples

Sample	Province	District	Village	Palmitic Acid	Stearic Acid	Oleic Acid	Linoleic Acid	Linolenic Acid	Eicosenic Acid
ÇT28	Artvin	Hopa	Yoldere	17,2	2,43	56,96	22,25	0,34	0.72
ÇT15	Artvin	Borçka	Muratlı	17,84	3,2	53,87	23,68	0,3	0.79
ÇT16	Artvin	Borçka	Güreşen	17,32	1,18	57,97	22,63	0,28	0.56
ÇT27	Artvin	Hopa	Koyuncular	16,08	2,15	61,04	19,79	0,23	0.61
ÇT26	Artvin	Hopa	Koyuncular	16,83	2,2	59,05	20,84	0,28	0.68
ÇT9	Giresun	Eynesil	Altınlı	19,81	2,34	48,35	27,48	0,52	1.23
ÇT12	Giresun	Tirebolu	Demircili Mah.	17,17	1,93	57,02	22,84	0,23	0.67
ÇT10	Giresun	Eynesil	Boztepe Mah.	19,41	1,79	53,82	24,13	0,23	0.61
ÇT13	Giresun	Tirebolu	Karaahmetli	17,27	1,57	59,08	21,15	0,24	0.69
ÇT14	Giresun	Tirebolu	Kovapınar	16,88	1,45	60,77	20,07	0,32	0.43
ÇT49	Rize	Çayeli	Yaka	16,52	3,52	56,56	22,38	0,23	0.71
ÇT50	Rize	Çayeli	Yalıköy	17,27	2,44	57,7	21,51	0,25	0.74
ÇT20	Rize	Pazar	Yemişli	16,76	1,61	59,69	20,62	0,25	0.55
ÇT51	Rize	Çayeli	Eskipazar	16,75	1,27	58,38	22,67	0,31	0.62

ÇT32	Rize	Merkez	Hayrat	17,82	2,02	55,92	23,17	0,27	0.80
ÇT1	Rize	Fındıklı	Sümer	17,6	2,75	55,79	21,88	0,8	0.00
ÇT45	Rize	Güneysu	Ulucami	17,33	2,22	58,03	21,65	0,22	0.53
ÇT54	Rize	Derepazarı	Bürücek	17,27	2,03	56,44	23,07	0,29	0.71
ÇT41	Rize	Kalkandere	Aşağı Tatlısu	18,21	1,42	53,66	25,56	0,34	0.57
ÇT43	Rize	Kalkandere	Ormanlı	16,77	2,52	58,47	21,45	0,2	0.59
ÇT37	Rize	İyidere	Yapraklar	17,98	2,55	53,63	24,83	0,22	0.73
ÇT36	Rize	İyidere	Köşklü	16,18	2,14	61,3	19,58	0,2	0.55
ÇT35	Rize	Kalkandere	Fındıklı	16,57	1,57	61,94	18,98	0,21	0.50
ÇT52	Rize	Derepazarı	Bahattinpaşa	16,26	2,45	57,94	21,62	0,15	0,57
ÇT21	Rize	Pazar	Orta Irmak Köyü	15,72	1,94	61,54	19,77	0,18	0,53
ÇT2	Rize	Fındıklı	Derbent	17,19	2,43	55,16	23,81	0,27	0,98
ÇT31	Rize	Ardeşen	Köprüköy	17,35	2,32	57,74	21,72	0,25	0,61
ÇT48	Rize	Çayeli	Derecik	16,85	2,37	57,02	22,77	0,28	0,64
ÇT53	Rize	Derepazarı	Kirazdağı	10,31	1,3	34,09	12,84	0,11	0.00
ÇT44	Rize	Kalkandere	Çağlayan	15,75	2,31	60,25	20,76	0,23	0.60
ÇT33	Rize	Merkez	Ortapazar	17,43	2,01	56,92	22,43	0,33	0.79

ÇT3	Rize	Fındıklı	Derbent-SelazurMevki	18,37	2,29	54,99	22,84	0,3	1.04
ÇT4	Rize	Fındıklı	Beydere	18,65	2,67	50,72	23,59	0,33	1.00
ÇT17	Rize	Hemşin	Ortaköy	16,75	1,72	58,63	21,87	0,28	0.64
ÇT47	Rize	Güneysu	Dumankaya	17,84	2,18	54,48	24,45	0,3	0.70
ÇT42	Rize	Kalkandere	Yolbaşı	17,26	2,04	57,36	22,44	0,27	0.64
ÇT19	Rize	Hemşin	Nurluca	18,27	2,84	56,68	21,23	0,22	0.69
ÇT29	Rize	Ardeşen	Köprüköy	17,47	1,53	57,53	22,44	0,24	0.61
ÇT34	Rize	Merkez	Ortapazar	15,75	1,52	61,72	19,92	0,25	0.6
ÇT22	Rize	Pazar	AkbucakKöyü	16,16	2,38	59,81	20,47	0,37	0.68
ÇT30	Rize	Ardeşen	Tunca Esentepe	16,71	1,95	60,42	19,93	0,21	0.55
ÇT18	Rize	Hemşin	Bilenköy	17,34	1,31	58,87	21,44	0,25	0.63
ÇT5	Rize	Fındıklı	Aslandere	20,74	2,15	48,75	26,36	0,45	1.23
ÇT40	Rize	İkizdere	İhlamur	15,79	1,67	62,41	19,1	0,27	0.56
ÇT46	Rize	Güneysu	Yüksekköy	17,68	2,3	54,46	24,46	0,29	0.74
ÇT39	Rize	İkizdere	Gürdere	15,33	2,85	60,49	20,13	0,36	0.70
ÇT38	Rize	İkizdere	Ilıca	16,36	1,61	58,65	22,24	0,45	0.69
ÇT25	Trabzon	Of	Balıca	16,63	1,93	60,03	20,37	0,23	0.53

ÇT6	Trabzon	Sürmene	Gültepe	18,3	2,28	57,6	20,71	0,23	0.79
ÇT24	Trabzon	Of	Cumapazarı	16,61	2,28	58,17	21,7	0,29	0.71
ÇT7	Trabzon	Sürmene	Güney Köyü	20,00	2,26	49,94	25,79	0,58	1.23
ÇT23	Trabzon	Of	Cumapazarı	15,40	2,32	61,28	19,92	0,31	0.67
ÇT11	Trabzon	Beşikdüzü	Şahmelik	15,97	1,45	55,42	18,21	0,25	0.54
ÇT8	Trabzon	Sürmene	Yeşilköy	15,37	2,76	59,76	20,78	0,38	0.75

UNDER PEER REVIEW

Principal Component Analysis of tea seed oil samples is given in Figure. 59.42% of the existing variation could be explained by the first two calculated principal components.




Fig. 1:Biplot of Tea Samples based on oil content and Fatty Acid Composition

Climate factors affect fatty acid composition. [29-30] When the climate changes, the likelihood of changes in fatty acid composition increases, with temperature strongly affecting the biosynthesis of fatty acids, especially unsaturated fatty acids. [31-32]

The climatic conditions of the cities and districts taken as a sample are different from each other. Since the tea plant is foreign pollinated, the differences obtained in this study are due to the heterogeneous structure of the tea plant and the effect of climatic factors.

We have to consider, that seeds were obtained after leaf harvest of tea plants. We can assume that the seed yield will increase if the tea plants are grown only for seed production. With governmental support tea seed oil production will be an potential additional income for local farmers.

PCA Analysis is helpful in genotype characterization and relative grouping calculated based on similarity. [33-34] PCA analysis can be used to differentiate plant materials, and the differences between various species can be visually revealed based on their chemical composition. [35-36] If these two methods are combined, characters that critically contribute to genetic variability in crops can be analyzed. [37] Biplot is another step in which one can group and identify the factors contributing to the variation obtained in PCA. [38]

Tea seed samples collected using this method could be separated from each other in terms of oil content and fatty acid composition. The data obtained provided important data for future studies on tea seeds.

4. CONCLUSION

Vegetable oils obtained by processing oilseeds; It has a special importance in human nutrition as well as great importance for our health. The physical and chemical properties of oils are determined by the ratios and types of fatty acids they contain. The fatty acid composition of oil plants is not always constant; It is under the control of genetic, ecological, morphological, physiological and cultural practices. The distribution of fatty acids varies significantly in ecological regions at different latitudes. Among environmental factors, the effect of temperature on fatty acid synthesis is especially evident.

Determining the effect of different latitudes and temperatures on the composition of fatty acids will enable regional production planning of oil crops and the production of vegetable oil for consumption purposes from different regions. Fatty acid composition is also affected by soil properties. Fatty acid compositions of oil crops are affected by genotype, as well as ecological and many other factors. In addition, fatty acids constantly change depending on the position of the seeds in the plant and during the period from the formation of the seed to its maturation. There may be differences in fatty acid composition between seeds harvested at different maturation periods; Therefore, when determining the fatty acid composition of seeds, their ripening periods should also be taken into account. As a result, fatty acid composition shows characteristic differences specific to plant species. In addition, the specific fatty acid composition of each oil plant is not constant and constantly changes depending on many factors.

Therefore, knowing under what conditions the fatty acid composition of oil plants will change is important in terms of oil quality. At the same time, the amount and type of fatty acids determine the usage method. Knowing the fatty acid composition of oils will enable production of oils according to their intended use. For this purpose, it will be possible to produce oils suitable for the purpose by growing the desired types in suitable regions.

Producing tea seeds and obtaining tea seed oil is a new topic for local tea farmers. Concerning the health issues of *C. sinensis* oil and its remarkable fatty acid composition suitable as edible oil, the support of tea seed oil production by the government will open a new additional income for local tea farmers at the Black Sea Region.

The oil content and fatty acid composition of collected tea samples varied between locations, which reveals a huge potential for oil production for tea plantations in Türkiye if governmental support is present.

ACKNOWLEDGEMENTS

There is no acknowledgement.

COMPETING INTERESTS

Authors have declared that no competing interests exist.”.

AUTHORS' CONTRIBUTIONS

Fatih Seyis designed the study. Murat Genc collected the seed samples and carried out the oil content and fatty aci composition analysis. Fatih Seyis wrote the whole manuscript.And managed the literature searches. All authors read and approved the final manuscript.”

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES

1. Van Broekhuizen P. Technical and non-technical aspects in the substitution of mineral oil based products by vegetable alternatives. *Agro-Food-Industry Hi-Tech*. 2001; 3: 39.
2. Öncü, M. The Tea Sector Turkish Economy. Ph.D.-Thesis. Bursa Uludağ University; 2019.
3. Roscoe T, Domergue F, Devic M, Lessire R. Improving Quality and Content of Oils in Seeds: Strategies, Approaches, and Applications Towards Engineering New Oilseed Crop Plants. *Seed Development: OMICS Technologies Toward Improvement of Seed Quality and Crop Yield*. 2012; 527–554.
4. Yazicioglu T, Karaali A, Goekcen J. Turkish tea oil and tea saponin. *Fett. SeifenAnstrichum*. 1997; 79: 115–120.
5. Ruter JM. Nursery production of tea oil camellia under different light levels. ASHS Press, Alexandria,VA; 2002.
6. Sahari MA, Ataii D, Hamedi M. Characteristics of tea seed oil in comparison with sunflower and olive oils and its effect as a natural antioxidant. *Am Oil Chem Soc*. 2004; 81: 585–588.
7. Fattahi-far E, Sahari MA, Barzegar M.. Interesterification of tea seed oil and its application in margarine production. *Journal of the American Oil Chemists' Society*. 2006; 83: 841-845.

8. Ravichandran R. Fat stability and amino acids in south Indian tea seeds. *International Journal of Food Science and Technology*. 1993; 28: 639-646.
9. Rajaei A, Barzegar M, Sahari MA. Comparison of antioxidative effect of tea and sesame seed oils extracted by different methods. *J Agric Sci Technol*. 2008; 10:345-350.
10. Rajaei A, Barzegar M, Yamini Y. Supercritical fluid extraction of tea seed oil and its comparison with solvent extraction. *Eur Food Res Technol*. 2005; 220:401-405.
11. Backhaus K, Erichson B, Plinke W, Weiber R. *Multivariate Analysis Methods*. Springer Verlag: Heidelberg; 1989
12. Tian HZ, Qiu AY, Shi XH. Extraction of tea seed polysaccharide. *China Oil (Chinese)*. 2004; 29:27–29.
13. Ravichandran R, Dhandapani M. Composition characteristics and potential uses of south Indian tea seeds. *J Food Sci Tech Mysore*. 1992; 29: 394–396.
14. Xia L, Zhang A, Xiao Y. An introduction to the utilization of camellia oil in China, *Am Camellia Yearbook*. 1993; 48: 12–15.
15. Wang Y, Sun D, Chen H, Qian L, Xu P. Fatty acid composition and antioxidant activity of tea (*Camellia sinensis* L.) seed oil extracted by optimized supercritical carbon dioxide. *Int J Mol Sci*. 2011; 12: 7708–7719.
16. Huang J, He J, Yin R, Wan X, Guo J, Xin C, Gong F, Li Y. Variations of fruit and seed traits of natural and artificial populations in *Camellia reticulata* L. *J Beijing Forest Univ*. 2010; 32, 94–101.
17. Kang Z, Zou D, Wang Z. Analysis on formation of tea oil camellia output by grey theory system. *J. Zhejiang Agroforest Sci Technol*. 1993; 12: 4–6.
18. Linder CR. Adaptive evolution of seed oils in plants: accounting for the biogeographic distribution of saturated and unsaturated fatty acids in seed oils. *Am Nat*. 2000; 156:442–458.
19. Debin C, Yangzhu W. The opportunity and countermeasures of Camellia industry development in Hunnan Province. *Hunan Forest Sci Technol*. 2002; 29: 3–5.
20. Junang L, Huaping P, Guoying Z. Study on model of yield and density of *Camellia oleifera*. *Forest Res Manage*. 2010; 5: 11–14.
21. Patel PK, Das B, Sarma R, Gogoi B. Tea Seed: A Review. *Annual research & Review in Biology*. 2018; 25(2): 1-14.
22. Owuor PO, Chauhanji AM, Manavu RM. Chemical studies of the Kenian tea seeds I. Physical and chemical characteristics of the Kernel oil. *Tea*. 1985; 6 (1): 23-28.
23. Zhang D, Stack L, Zhang R, Yu J, Xie B, Chen Y, Ruter JM. Tea oil camellia– eastern “olive” for the world. *ISHS Acta Hort*. 2008; 769:43–48.
24. Anon. Oils by Nature Inc. Camellia Oil (Expeller Pressed). <http://www.oilsbynature.com/products/camellia-oil.htm>; 2007.
25. Lee, CP, Yen GC. Antioxidant activity and bioactive compounds of tea seed (*Camellia oleifera* Abel.) oil. *Journal of Agriculture and Food Chemistry*. 2006; 54: 779-784.
26. Lee CP, Shih PH, Hsu CL, Yen GC. Hepatoprotection of tea seed oil (*Camellia oleifera* Abel.) against CCl₄-induced oxidative damage in rats. *Food Chem Toxicol*. 2007; 45: 888-895.

27. Fazel M, Sahari MA, Barzegar M. Determination of main tea seed oil antioxidants and their effects on common kilka oil. *International Food Research Journal*. 2008; 15: 209-217.
28. Fattahi-far E, Sahari MA, Barzegar M. Interesterification of tea seed oil and its application in margarine production. *Journal of the American Oil Chemists' Society*. 2006; 83: 841-845.
29. Munasinghe M, Wansapala J. Study on variation in seed morphology, oil content and fatty acid profile of *Madhuca longifolia* grown in different agro-climatic zones in Sri Lanka. *Sci Res*. 2015; 3: 105–109.
30. Ruraz K, Piwowarczyk R, Gajdoš P, Krasylenko Y, Certík M. Fatty acid composition in seeds of holoparasitic Orobanchaceae from the Caucasus region: Relation to species, climatic conditions and nutritional value. *Phytochemistry*. 2020; 179: 112510.
31. Li Y, Li S, Lu XH, Wang QQ, Han HY, Zhang XM, Ma YH, Gan XH. Leaf phenotypic variation of endangered plant *Tetracentron sinense* Oliv. and influence of geographical and climatic factors. *J For Res*. 2021; 32: 623–636.
32. Abdelghany AM, Zhang S, Azam M, Shaibu AS, Feng Y, Li Y, Tia, Y, Hong H, Li B, Sun, J. Profiling of fruit fatty acid composition in 1025 Chinese soybean accessions from diverse ecoregions. *Crop J*. 2022; 8: 635–644.
33. Mohammadi SA, Prasanna BN. Analysis of genetic diversity in crop plants-salient statistical tools and considerations. *Crop Science*. 2003; 43: 1235-1248.
34. Peeters JP, Martinelli JA. Hierarchical cluster analysis as a tool to manage variation in germplasm collections. *Theoretical and Applied Genetics*. 1989; 78: 42-48.
35. Smelcerovic A, Zuehlke S, Spiteller M, Raabe N, Ozen T. Phenolic Constituents of 17 *Hypericum* species from Türkiye. *Biochemical. Systematics and Ecology*. 2008; 36: 316–319.
36. Bertoli A, Cirak C, Leonardi M, Seyis F, Pistelli L. Morphogenetic changes in essential oil composition of *Hypericum perforatum* during the course of ontogenesis. *Pharmaceutical Biology*. 2011; 49: 741–751.
37. Rachovska, G., Dimova, D., Bojinov, B. (2003). Application of cluster analysis and principal component analysis for evaluation of common winter wheat genotypes. *Proceedings of the Scientific Session of Jubilee 2002-Sadovo*. pp. 68-72.
38. Aghaee M, Mohammad, R, Nabovat, S. Agro-morphological characterization of durum 16 wheat accessions using pattern analysis. *Australian Journal of Plant Sciences*. 2010; 4: 505-514.