

Original Research Article

Screening of soya bean genotypes for agronomic traits and cercospora (*Cercospora sojina*) leaf spot disease in western Tigray, Ethiopia.

Evaluation of Agronomic Performance and Resistance to Cercospora (*Cercospora sojina*) Leaf Spot of Selected Soybean Genotypes in Western Tigray

Abstract

Aim of the current study was to evaluate soya bean genotypes for ~~for agronomic performance, yield and yield traits~~ and resistance to cercospora (*Cercospora sojina*) leaf spot disease under field condition. Field experiment was conducted at Humera district during 2018 cropping season using 16 soya bean genotypes. Treatments were evaluated with RCB and replicated three times. Number of days from planting to days of flowering and days of maturity, number of pod per plant, pod shattering, seed yield and cercospora leaf spot disease were showed a significant ($P < 0.01$) variation among genotypes. All varieties were moderately resistance to pod shattering except PN13-1 (3.1) was moderately susceptible and Parc-3, Parc-5 and PN13-10 were resistance. Highest yields were recorded from genotypes PN13-2 at 20.32q/ha and PN13-9 at 19.77q/ha. On the other hand, the lowest yields were recorded from genotypes PN13-3 at 7.49q/ha and PN13-1 at 9.95q/ha. Genotypes PN13-2 and PN13-7 have been identified as displaying resistance to *Cercospora* leaf spot. Highest disease severity recorded in genotypes Awassa 04 and PN13-1. Further research is needed in future to cover a wide range of environments and genotypes to screen high yielder and cercospora leaf spot disease resistance.

Key word; Soya bean Genotypes, seed yield, pod shattering and cercospora leaf spot

1. Introduction

Soybean (*Glycine max* (L.) Merrill) is one of the most important oil grain legume crops in the world [1]. Soybean stand out as a nutritional powerhouse due to ~~its~~ their high protein and oil content, as well as the presence of functional components like flavones that offer various health benefits [2]. In Ethiopia, soybean is used for a variety of purposes including preparation of different kinds of soybean foods, animal feed and soy milk [3]. Soybean is classified into different maturity groups based on the length of

the growing period from planting to maturity. These maturity groups help choose the right genotypes that will mature within the suitable growing season in their region. The classification typically includes early, medium, and late maturing varieties. This phenological attribute is determined by two abiotic factors: photoperiod and temperature [4], and these factors can dictate the most suitable maturity groups of soybean varieties for a particular geographical location [2].

The production and productivity of soya bean is affected by abiotic and biotic factors. The biotic constraints including diseases, insect pests, and weeds are limiting the crop yield. The major economically important diseases are rust, wilts, leaf spot, rots, powdery mildew, bacterial and viral diseases [5]. Cercospora leaf spot is a common fungal disease that affects a variety of crops, including soybeans. It is caused by the fungus *Cercosporasojina* and can lead to significant yield losses if not properly managed. The disease typically manifests as small, dark spots on the leaves, which can coalesce and cause defoliation in severe cases [6]. Cercospora leaf spot cause a significant yield loss of soybean (10–60%) [7]. Management strategies for Cercospora leaf spot in soya bean include cultural practices such as crop rotation, planting resistant varieties, and maintaining proper plant spacing to improve air circulation.

Screening of early mature groups of soya bean genotypes and resistance to cercospora (*Cercosporasojina*) leaf spot disease in Western Tigray, Ethiopia, is a crucial agricultural practice aimed at identifying and selecting soya bean genotypes that exhibit early maturity traits and cercospora leaf spot disease resistance. This process involves evaluating different genotypes of soya beans to determine their performance in terms of early maturity, yield potential, disease resistance, and other agronomic characteristics.

2. Material and methods

2.1. Description of experimental site

The field experiment was conducted during 2018 in main crop growing season at Humera station. The experimental site is described as having hot to warm temperatures and high evaporation conditions. This specific environment is classified as a hot to warm semiarid lowland agro ecology. In such areas, the climate tends to be characterized by high temperatures and limited precipitation, leading to increased evaporation [8]. The site is located at 14° 00' 85" North latitude and 36° 34' 52" East longitude. The elevation of this station is about 600 meters above sea level. The mean annual temperature of the area is 29 °c and the rainy months extend from late June to the middle of September. The remaining 8-9 months are dry and hot.

2.2. Experimental treatment and field management:

A total of 16 different soya bean genotypes were used in a field experiment (Table 1). The genotypes were obtained from Federal research institute. The experiment was laid out in randomized complete block design (RCBD) with three replications. Each treatment were randomly assigned into a plot area of 15m² (5m row length and 3m width), which consisted of 5 rows of soya bean. The spacing between block and plot will be 2m and 1m, respectively. The spacing between plants and rows were 5 cm and 60 cm, respectively. Seeds are sown on June 2017 on three time's ploughed plots of land. Each experimental plot are received the same rate of NPS (100 kg/ha). The other management practices were applied equally and properly as per the recommendations.

Table 1. Description of soya bean genotypes in this experiment

S.No.	Genotypes	S.No.	Genotypes
1	PN13-2	9	Awassa 04
2	PN13-3	10	Nyala
3	PN13-4	11	parc-3
4	PN13-5	12	parc-5
5	PN13-6	13	PN13-1
6	PN13-7	14	PN13-10
7	PN13-8	15	PN13-11
8	PN13-9	16	PN13-12

2.3. Data collection:

[Agronomic traits](#) ~~Yield components~~ including plant height, branches per plant, pod per plant, hundred seed weight, seed yield and shattering were ~~measured~~ [determined](#). Plant height and branches per plant were measured from ten randomly selected plants in each plot. However, number of pods per plant was counted from ten randomly selected plants. The weight of hundred counted seeds was recorded in gram for individual plots. Similarly, seed yield (kg/ha) was measured from threshed and cleaned plots separately using a sensitive balance and converted into kilogram per hectare.

2.4. Pod shattering:

Pod shattering was measured on visual observation in the field. The number of shattered pods were counted and expressed as percentage. [Varieties were classified into five categories based on their reaction to pod shattering as described in table 2.](#)

Comment [MOU1]: Provide a citation for this scale

Table 2. Percent and reaction for recording pod shattering.

Scale	Percentage	Reaction
1	0	No shattered
2	1-10	Resistance
3	>10-25	Moderately resistance
4	>25-50	Moderately susceptible
5	>50	Susceptible

2.5. Disease data:

- Disease incidence(%): it was calculated by $(\text{total infected plant} \times 100) / \text{total observed plant}$.
- PSI(%) for easy od analysis: it was calculated by $(\text{sum of rating scales} \times 100) / (\text{No. of plants scored} * \text{maximum score of scale})$

❖ Percent severity index (PSI)

The mean per cent disease index was transformed into disease reaction as 0%=No infection/immune; 0-10%= resistance (R); 10.1-20%=moderately resistant (MR); 20.1- 30%=moderately susceptible (MS); 30.1-50%= susceptible (S) and >50%=highly susceptible (HS). The genotypes were categorized into resistance groups (0-9) accordingly to rating score description; 0: (0) Leaves apparently free from spots; 1: (0.01-1.0) Very small area of leaf covered with lesions; 3: (1.1-10) Considerable leaf area covered with spots, no spots on stem; 5: (10.1-25) One- fourth of leaf area covered with spots, no defoliation of plants; little damage; 7: (25.1-50) Some leaves dropped, death of a few plants, damage to plant is conspicuous; 9: (> 50) More than half of the leaf area covered with spots, lesions very common on all plants, defoliation common, death of plants is common[9].

2.6. Data analysis:

Analysis of variance was ~~conducted~~ for ~~all traits measured in this study~~ yield and yield components (stand count at emergence, days to 50% flowering and maturity 95%, plant height, branches per plant, number of pods per plant, and and thousand seed weight) from the field experiment, to determine know the main effects and their interactions using Genstat version 18 software. Least Significant Difference (LSD) values were used to separate differences among treatment means at 5% probability level.

3. Result and discussion

3.1. Agronomic traits

3.1.1. Day of flowering:

There was a significant variation observed in the number of days from planting to flowering among tested soya bean genotypes underscores the importance of genetic diversity in crop improvement programs. In the given data, the lowest days of flowering was measured from genotypes Parc-3 (34), Parc-5 (34), and PN13-12 (37). On the other hand, the highest days of flowering was observed in genotypes PN13-7 (50.77), PN13-1 (50.67), and PN13-5 (49.67) (Table 3). This result suggests that there is a significant variation in the flowering dates among different genotypes. The differences in flowering times can be influenced by various genetic factors, environmental conditions, and interactions between genes and the environment.

Comment [MOU2]: Provide a P-value

3.1.2. Date of maturity:

It was found that there was a significant variation in the number of days from planting to maturity among the tested genotypes. The statistical analysis revealed a p-value of less than 0.01, indicating a high level of significance in the differences observed. The overall average of maturity ranged from 64 to 92.67 days. The lower end of the range at 64 days implies instances where maturity is achieved relatively quickly, while the upper end at 92.67 days indicates cases where a longer period is required for maturity to be reached. Maximum day of maturity was recorded from genotypes PN13-7 (92.67), PN13-3 (92), PN13-5 (90.67) and PN13-6 (90.67). However, minimum date of maturity was recorded from Parc-5 (64), Parc-3 (69) PN13-12 (74.33) and PN13-4 (77). Intermediate days of maturity were recorded from genotypes PN13-10 (80.67), PN13-2 (82.33) and PN13-9 (82.67) (Table 3). The result was in similar [10] who stated that the range of days to maturity from 107.33 to 79.33 shows variability among the different soya bean varieties in terms of how long they take to mature. This result was in agreement with the report of [11] who reported that days to maturity were significantly affected by soybean varieties.

Comment [MOU3]: Whenever you state that there were significant differences, please, provide a P-value

3.1.3. Branches per plant and Plant height:

In the study of soya bean genotypes, it was found that there wasn't significant variation among most of the genotypes except for the comparison between Parc-5 and PN13-8 genotypes. This suggests that the branches per plant in soya bean genotypes did not differ significantly, except for these two specific genotypes. In the study comparing different genotypes of soybean for plant height, it was found that there was a highly significant difference in plant height among the tested genotypes. The significant difference in plant height indicates that certain genotypes may have genes that promote taller growth, while others may have genes that restrict vertical growth. Highest plant height was measured from genotypes Awassa 04 (64), Parc-3 (63.33), PN13-6 (61.33) and PN13-7 (61). However, lowest plant height was measured from genotypes PN13-3 (33.33) and PN13-1 (37.33) (Table 3). These findings were in line with [12] who

reported that plant height significantly difference among soybean genotypes. Similarly, [13] also evaluated soybean varieties resulted a significantly difference at plant height in soybean varieties.

3.1.4. Pod per plant:

Pod per plant was showed that significant ($P<0.01$) difference among the soya bean genotypes. Maximum number of pod counted from genotypes PN13-2 (72.67) followed by PN13-9 (62.67) and PN13-6 (54.33). While, minimum number of pod was counted from PN13-3(24) and PN13-1 (28.67). The data provided indicates that the genotypes PN13-2 and PN13-9 had significantly higher yields in terms of the number of pods produced compared to other genotypes in the study. Conversely, genotypes PN13-3 and PN13-1 had lower yields with fewer pods produced overall (Table 3). It could be due to; water deficit poses a significant threat to soybean production by reducing pod formation, seed quantity, and seed quality. The current findings align with a study by [14] which highlighted the impact of water deficiency on yield and yield traits. Water is a critical factor in plant growth and development, influencing various aspects of crop productivity.

Table 3. Mean values of yield and yield component of soya bean genotypes

Genotypes	DF (50%)	DM (95%)	NBP	PH (cm)	PPP
Awassa 04	44.67bcd	82.67bcd	1.67ab	64a	39.67b-e
Nyala	48.33abc	88.33abc	1.33ab	59.67ab	39b-e
parc-3	34f	69.67ef	1.33ab	63.33a	38b-e
parc-5	34f	64f	1b	44.67cd	33.67cde
PN13-1	50.67ab	91ab	2ab	37.33de	28.67de
PN13-10	43cde	80.67cd	1.67ab	54.67ab	41.33b-e
PN13-11	40def	83bcd	2ab	42cd	35b-e
PN13-12	37ef	74.33de	1.33ab	56.67ab	43.33b-e
PN13-2	45.67a	82.33bcd	2ab	59.67ab	72.67a
PN13-3	47.67abc	92a	1.67ab	33.33f	24e
PN13-4	48.67abc	77de	1.67ab	51bc	45bcd
PN13-5	49.67abc	90.67ab	2ab	55.33ab	41.33b-e
PN13-6	49.33abc	90.67ab	1.67ab	61.33ab	54.33b
PN13-7	50.77ab	92.67a	1.67ab	61ab	46.67bcd
PN13-8	49abc	89abc	2.333a	38.67ef	31de
PN13-9	43.67cd	82.67bcd	1.67ab	57.67ab	62.67bc

Lsd (5%)	5.82	7.65	0.92	9.11	17.44
CV (%)	7.7	5.5	32.8	10.7	24.5

Note; DF: Date of flowering; DM: Date of maturity; PPP: pod per plant; PH: Plant height;

3.1.5. Hundred seed weight:

There was a significant variation in hundred seed weight between the genotypes. The heaviest seed weights were measured from three different soya bean genotypes; PN13-12, PN13-11, and Parc-5. The specific weights for these seeds were 12.4 grams, 11 grams, and 10.13 grams, respectively. On the other hand, the smallest seed weights were recorded from two other soya bean genotypes; PN13-2 and Nyala. The specific weights for these seeds were 6.1 grams and 6.7 grams, respectively.

3.1.6. Seed yield:

In the study conducted on soya bean genotypes, there was a significant difference in seed yield among the tested genotypes. Overall result of seed yield ranged from 7.49q/ha to 20.32q/ha. The highest yields were recorded from genotypes PN13-2 at 20.32q/ha and PN13-9 at 19.77q/ha. On the other hand, the lowest yields were recorded from genotypes PN13-3 at 7.49q/ha and PN13-1 at 9.95q/ha. Early maturing soybean varieties allow farmers to harvest their crops sooner, reducing the risk of yield losses due to adverse weather conditions or pest infestations. The result was supported by [10] early-maturing varieties (Coker and Gonzela) are suitable for short rainfall areas.

3.1.7. Pod shattering:

Pre-harvest soybean pod shatter refers to the phenomenon where soybean pods split open prematurely before harvest, leading to yield losses. This issue is particularly concerning as it can significantly affect the overall yield potential of a soybean crop, especially if it occurs while the plants are still in their green stages. Pod shattering based on the shattering level shows there was significant ($P < 0.05$) variation among the soya bean genotypes. All genotypes were moderately resistance to pod shattering except PN13-1 (3.1) was moderately susceptible and Parc-3, Parc-5 and PN13-10 were resistance (Table 4). The result was supported by [15, 16] who reported that five soya bean lines was resistance (7-10%) to pod shattering. In addition also [17] stated that genotypes SB-8, Gazelle, SB-74, SB-4 and Nyala were the most resistant to pod shattering and high grain yield, while Genotypes SB-90 and SB-25 were highly susceptible.

Table 4. Mean values of hundred seed weight, yield and Pod shattering character of soya bean genotypes

Genotypes	Hundred seed weight	Seed yield(q/ha)	Shattering	
			scale	Reaction
Awassa 04	8.77cde	13.37abc	2.67a	MR
Nyala	6.7f	12.2abc	2.33ab	MR
parc-3	9.47cd	12.49abc	1.33ab	R
parc-5	10.13bc	10.6abc	1b	R
PN13-1	7.73e	9.95bc	3.1ab	MS
PN13-10	9.8bcd	13.92abc	2ab	R
PN13-11	11b	16.68abc	1.33ab	MR
PN13-12	12.4a	13.85abc	1.67ab	MR
PN13-2	6.1f	20.32a	1.67ab	MR
PN13-3	7.67e	7.49c	1.67ab	MR
PN13-4	8.4de	13.11abc	2.33ab	MR
PN13-5	8.4de	14.09abc	1.67ab	MR
PN13-6	8.67cde	15.9abc	1.67ab	MR
PN13-7	7.43ef	11.54abc	2ab	MR
PN13-8	7.4ef	12.92abc	1.67ab	MR
PN13-9	8.37de	19.77ab	1b	
Lsd (5%)	1.36	8.49	1.158	
CV (%)	9.4	37.3	39.7	

Note; q/ha: Quintal per hectare

3.2. Incidence and severity of *Cercospora*(*Cercosporasojina*)leaf spot disease:

In the study conducted on soybean genotypes to evaluate the incidence of *Cercospora* leaf spot, it was found that there wasn't a significant variation among the tested soybean genotypes. This indicates that the different genotypes of soybeans showed similar levels of susceptibility or resistance to *Cercospora* leaf spot. While, severity shows there was a significant variation among the tested soya bean genotypes. Genotypes PN13-2 and PN13-7 have been identified as displaying resistance to *Cercospora* leaf spot, with low disease severity recorded in comparison to other genotypes. The disease severity ratings for these genotypes were 1.03 for PN13-2 and 1.4 for PN13-7. Highest disease severity recorded in genotypes Awassa 04 and PN13-1. Both genotypes exhibited a disease severity rating of 2.7 and 2.67, respectively. These ratings indicate that these genotypes are moderately susceptible to the disease (Table 5). *Cercospora* leaf spot is a common fungal disease which is recorded with a severity on an average of 2.50 to 15.75% [18]. The result supported [7]; who identified three genes, Rcs1, Rcs2, and cs3, for frog eye

leaf spot resistance in soybeans. These genes have been officially named by the Soybean Genetics Committee.

Table 5. Incidence and severity of *Cercospora (Cercosporasojina)* leaf spot on soya bean genotypes

Genotypes	Leaf spot			
	Incidence (%)	Severity (PSI)	Percentage (%)	reaction
Awassa 04	100	2.7a	29.5	MS
Nyala	100	2.47ab	24.7	MS
parc-3	100	2.4 7ab	24.2	MR
parc-5	100	1.7ab	13.2	MR
PN13-1	100	2.67ab	25.4	MS
PN13-10	100	2.47ab	24.2	MR
PN13-11	100	2.47ab	24.2	MR
PN13-12	100	2.03ab	18.9	MR
PN13-2	100	1.03b	8.9	R
PN13-3	100	2.03ab	18.6	MR
PN13-4	100	2.03ab	18.6	MR
PN13-5	100	2.47ab	24.2	MR
PN13-6	100	2.03ab	17.9	MR
PN13-7	100	1.4ab	9.2	R
PN13-8	100	2.47ab	24.5	MR
PN13-9	100	2.03ab	17.5	MR
Lsd (5%)	ns	2.46		
CV (%)	*	29.3		

Note; R: Resistance; MR: Moderately resistant, MS: Moderately susceptible, ns: Non-significant, PSI: Percentage severity index

4. Conclusion

Soybean crop plays a crucial role in Ethiopia's agriculture sector and contributes significantly to food security and nutrition. Host resistance is the most effective and economical management practice for frogeye leaf spot. Overall result of seed yield ranged from 7.49q/ha to 20.32q/ha. The highest recorded

yields were from genotypes PN13-2 at 20.32q/ha and PN13-9 at 19.77q/ha. On the other hand, the lowest yields were recorded from genotypes PN13-3 at 7.49q/ha and PN13-1 at 9.95q/ha. Genotypes PN13-2 and PN13-7 have been identified as displaying resistance to *Cercospora* leaf spot. Highest disease severity recorded in genotypes Awassa 04 and PN13-1. Both genotypes exhibited a disease severity rating of 2.7 and 2.67, respectively. Further research is needed in future to cover a wide range of environments and genotypes to screen high yielder and disease resistance.

Reference

1. Laswai HS, Mpanalile JJ, Silayo VC, Ballegu WR. Use of soybeans in food formulation in Tanzania. In Proceedings of the first national soybean stakeholders workshop 2005 Nov 11 (pp. 52-59). Morogoro, Tanzania: Sokoine University of Agriculture.
2. Liu X, Wu JA, Ren H, Qi Y, Li C, Cao J, Zhang X, Zhang Z, Cai Z, Gai J. Genetic variation of world soybean maturity date and geographic distribution of maturity groups. *Breeding science*. 2017;67(3):221-32.
3. Hailu M, Kelemu K. Trends in soy bean trade in Ethiopia. *Research Journal of Agriculture and Environmental Management*. 2014 Sep;3(9):477-84.
4. Funga, A., Bekele, D., Monyo, E., Tadesse, M, Mohamed, R., Gaur, P., Eshete, M, Ojiewo, C, Bishaw, Z, Fikre, A, Rao, G., Korbu, L., Girma, N. and Siambi, M., Genotype by environment interaction on yield stability of desi type chickpea (*Cicer arietinum* L.) at major chickpea producing areas of [77] Ethiopia. *Australian Journal of Crop Science*, 2017; 11(02):212-219.
5. Jahagirdar S. Present status and future research dimensions of soybean diseases for sustainable productivity of soybean in India. *Indian Phytopathology*. 2019 Mar 13;72(1):3-14.
6. Borah, M., Identification of soybean diseases in assam. *International Journal of Recent Scientific Rese*. 2019.
7. Lin B, Kelly H. Frogeye leaf spot of soybean. *The Plant Health Instructor*. 2018.
8. GeremewTerefe, AdugnaWakjira, MuezBerhe and Hagos Tadesse. *Sesame Production Manual*. EIAR and Embassy of the Kingdom of the Netherlands.pp. 2012; 134.
9. Robinson EH, Li MH. Channel catfish, *Ictalurus punctatus*, nutrition in the United States: A historical perspective. *Journal of the World Aquaculture Society*. 2020 Feb;51(1):93-118.
10. Tariku A, Begna T, Asrat Z, Gojam M. Evaluation of early and medium maturity released soybean varieties for agronomic performance and adaptation at Meiso District of West Harerghea zone, Eastern Ethiopia. "(2023): 31-35.

11. Hunde D, Tefera G. Participatory varietal selection and evaluation of twelve soybeans *Glycine max* (L.) Merrill] varieties for lowland areas of North Western Ethiopia. *International Journal of Plant Breeding and Crop Science*. 2018;5(2):403.
12. Yechalew S, Masresha Y, Mesfin H, Bahailu A. Performance of Released Soybean Varieties at Jimma, South Western Ethiopia. *Journal of Biology, Agriculture and Healthcare*. 2020;10(4):12-5.
13. Awoke T. Performance evaluation of soybean (*Glycine max* L.) varieties for growth, yield and yield components under irrigation at lowland area of south Omo zone, Southern Ethiopia. *Journal of Agriculture and Aquaculture*. 2022;4(2):5-10.
14. Basal O, Szabó A. The combined effect of drought stress and nitrogen fertilization on soybean. *Agronomy*. 2020 Mar 11;10(3):384.
15. Krisnawati A, Soegianto A, Waluyo B, Kuswanto, The pod shattering resistance of soybean lines based on the shattering incidence and severity. *Czech J. Genet. Plant Breed*. 2020 Jun 22; 56: 111-22.
16. Adie MM, Sundari T, Wijanarko A, Purwaningrahayu RD, Krisnawati A. Identification of Pod Shattering Resistance and Associations between Agronomic Characters in Soybean using Genotype by Trait Biplot. *Legume Research-An International Journal*. 2022; 45 (1):18-24.
17. Katembo KR. Genetic analyses of pod shattering and agronomic traits of soybean (*glycine max* (L.) merr.) genotypes (Doctoral dissertation, University of Nairobi). 2018.
18. Huilgol SN, Hiremath I, Hegde YR, Basavaraja GT. First record of target leaf spot disease of soybean and progression of other fungal foliar spots in Karnataka. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(4):3203-5.